


A Solution Manual For

**Elementary differential equations with
boundary value problems. William F.
Trench. Brooks/Cole 2001**



Nasser M. Abbasi

May 16, 2024

Contents

1	Chapter 1, Introduction. Section 1.2 Page 14	3
2	Chapter 2, First order equations. Linear first order. Section 2.1 Page 41	22
3	Chapter 2, First order equations. separable equations. Section 2.2 Page 52	65
4	Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60	98
5	Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68	116
6	Chapter 2, First order equations. Exact equations. Section 2.5 Page 79	181
7	Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6 Page 91	219
8	Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203	251
9	Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253	272
10	Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262	322
11	Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318	359
12	Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329	375
13	Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338	414

14	Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358	468
15	Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374	527
16	Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389	611
17	Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471	657
18	Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483	666
19	Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495700	700
20	Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503	777
21	Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540	795
22	Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555	813
23	Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566	847

1 Chapter 1, Introduction. Section 1.2 Page 14

1.1	problem 2(a)	4
1.2	problem 2(b)	5
1.3	problem 2(c)	6
1.4	problem 2(d)	7
1.5	problem 2(e)	8
1.6	problem 3(a)	9
1.7	problem 3(b)	10
1.8	problem 3(c)	11
1.9	problem 4(a)	12
1.10	problem 4(b)	13
1.11	problem 4(c)	14
1.12	problem 5(a)	15
1.13	problem 5(b)	16
1.14	problem 5(c)	17
1.15	problem 5(d)	18
1.16	problem 8(a)	19
1.17	problem 9	20
1.18	problem 10(a)	21

1.1 problem 2(a)

Internal problem ID [869]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x) = 2*y(x),y(x), singsol=all)
```

$$y(x) = e^{2x} c_1$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 16

```
DSolve[y'[x]== y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.2 problem 2(b)

Internal problem ID [870]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x + y = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x) +y(x)= x^2,y(x), singsol=all)
```

$$y(x) = \frac{x^3 + 3c_1}{3x}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 19

```
DSolve[x*y'[x] +y[x]== x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{3} + \frac{c_1}{x}$$

1.3 problem 2(c)

Internal problem ID [871]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2yx + y' = x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve(diff(y(x),x) +2*x*y(x)= x,y(x), singsol=all)
```

$$y(x) = \frac{1}{2} + e^{-x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 26

```
DSolve[y'[x] +2*x*y[x]== x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} + c_1 e^{-x^2}$$
$$y(x) \rightarrow \frac{1}{2}$$

1.4 problem 2(d)

Internal problem ID [872]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$2y' + x(-1 + y^2) = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(2*diff(y(x),x) +x*(y(x)^2-1)= 0,y(x), singsol=all)
```

$$y(x) = \tanh\left(\frac{x^2}{4} + \frac{c_1}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 52

```
DSolve[2*y'[x] +x*(y[x]^2-1)== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{\frac{x^2}{2}} - e^{2c_1}}{e^{\frac{x^2}{2}} + e^{2c_1}}$$
$$y(x) \rightarrow -1$$
$$y(x) \rightarrow 1$$

1.5 problem 2(e)

Internal problem ID [873]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - (1 + y^2)x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x) = x^2*(1+y(x)^2),y(x), singsol=all)
```

$$y(x) = \tan\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.171 (sec). Leaf size: 30

```
DSolve[y'[x] == x^2*(1+y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan\left(\frac{x^3}{3} + c_1\right)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

1.6 problem 3(a)

Internal problem ID [874]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = -x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x) = -x,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 15

```
DSolve[y'[x] == -x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{2} + c_1$$

1.7 problem 3(b)

Internal problem ID [875]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = -\sin(x)x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x) = -x*sin(x),y(x), singsol=all)
```

$$y(x) = -\sin(x) + \cos(x)x + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 16

```
DSolve[y'[x] == -x*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sin(x) + x \cos(x) + c_1$$

1.8 problem 3(c)

Internal problem ID [876]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = x \ln(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x) = x*ln(x),y(x), singsol=all)
```

$$y(x) = \frac{\ln(x) x^2}{2} - \frac{x^2}{4} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 24

```
DSolve[y'[x] == x*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x) + c_1$$

1.9 problem 4(a)

Internal problem ID [877]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = -x e^x$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x) = -x*exp(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = -(x - 1) e^x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 13

```
DSolve[{y'[x] == -x*Exp[x],y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^x(x - 1)$$

1.10 problem 4(b)

Internal problem ID [878]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = x \sin(x^2)$$

With initial conditions

$$\left[y\left(\frac{\sqrt{2}\sqrt{\pi}}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 12

```
dsolve([diff(y(x),x) = x*sin(x^2),y(1/2*2^(1/2)*Pi^(1/2)) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x^2)}{2} + 1$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 15

```
DSolve[{y'[x] == x*Sin[x^2],y[Sqrt[Pi/2]]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 - \frac{\cos(x^2)}{2}$$

1.11 problem 4(c)

Internal problem ID [879]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' = \tan(x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = 3 \right]$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 15

```
dsolve([diff(y(x),x) = tan(x),y(1/4*Pi) = 3],y(x), singsol=all)
```

$$y(x) = -\ln(\cos(x)) + 3 - \frac{\ln(2)}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[{y'[x] == Tan[x],y[Pi/4]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(\cos(x)) + 3 - \frac{\log(2)}{2}$$

1.12 problem 5(a)

Internal problem ID [880]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \tan(x)y = \cos(x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}\pi}{8} \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(x),x) = cos(x)-y(x)*tan(x),y(1/4*Pi) = 1/8*2^(1/2)*Pi],y(x), singsol=all)
```

$$y(x) = \cos(x)x$$

✓ Solution by Mathematica

Time used: 0.07 (sec). Leaf size: 9

```
DSolve[{y'[x] ==Cos[x]-y[x]*Tan[x],y[Pi/4]==Pi/(4*Sqrt[2])},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow x \cos(x)$$

1.13 problem 5(b)

Internal problem ID [881]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 5(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - \frac{x^2 - 2x^2y + 2}{x^3} = 0$$

With initial conditions

$$\left[y(1) = \frac{3}{2} \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([diff(y(x),x) = (x^2-2*x^2*y(x)+2)/x^3,y(1) = 3/2],y(x), singsol=all)
```

$$y(x) = \frac{\frac{x^2}{2} + 2 \ln(x) + 1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 19

```
DSolve[{y'[x] ==(x^2-2*x^2*y[x]+2)/x^3,y[1]==3/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{x^2} + \frac{2 \log(x)}{x^2} + \frac{1}{2}$$

1.14 problem 5(c)

Internal problem ID [882]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 5(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - x(1 + y^2) = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 10

```
dsolve([diff(y(x),x) = x*(1+y(x)^2),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan\left(\frac{x^2}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.161 (sec). Leaf size: 13

```
DSolve[{y'[x] ==x*(1+y[x]^2),y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan\left(\frac{x^2}{2}\right)$$

1.15 problem 5(d)

Internal problem ID [883]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 5(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \frac{y(1+y)}{x} = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 11

```
dsolve([diff(y(x),x) = (- y(x)*(y(x)+1))/x,y(1) = -2],y(x), singsol=all)
```

$$y(x) = \frac{2}{-2+x}$$

✓ Solution by Mathematica

Time used: 0.224 (sec). Leaf size: 12

```
DSolve[{y'[x] ==(- y[x]*(y[x]+1))/x,y[1]==-2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{x-2}$$

1.16 problem 8(a)

Internal problem ID [884]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 8(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - ay^{\frac{a-1}{a}} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 9

```
dsolve(diff(y(x),x) = a*y(x)^( (a-1)/a),y(x), singsol=all)
```

$$y(x) = (c_1 + x)^a$$

✓ Solution by Mathematica

Time used: 0.843 (sec). Leaf size: 28

```
DSolve[y'[x] ==a*y[x]^( (a-1)/a),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(x + \frac{c_1}{a}\right)^a$$

$$y(x) \rightarrow 0^{\frac{a}{a-1}}$$

1.17 problem 9

Internal problem ID [885]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - |y| = 1$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 19

```
dsolve([diff(y(x),x) = abs(y(x))+1,y(0) = 0],y(x), singsol=all)
```

$$y(x) = e^x - 1$$

$$y(x) = 1 - e^{-x}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x] ==Abs[y[x]]+1,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

```
{}
```

1.18 problem 10(a)

Internal problem ID [886]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 1, Introduction. Section 1.2 Page 14

Problem number: 10(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Clairaut]`

$$y' - \frac{\sqrt{x^2 + 4x + 4y}}{2} = -\frac{x}{2} - 1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x) = 1/2*(-(x+2)+sqrt(x^2+4*x+4*y(x))),y(x), singsol=all)
```

$$x - \sqrt{x^2 + 4x + 4y(x)} - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.801 (sec). Leaf size: 47

```
DSolve[y'[x] == 1/2*(-(x+2)+Sqrt[x^2+4*x+4*y[x]]),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow \frac{1}{4}(-2x + 2e^{c_1}(x + 1) + 1 + e^{2c_1}) \\y(x) &\rightarrow 1 \\y(x) &\rightarrow \frac{1}{4}(1 - 2x)\end{aligned}$$

2 Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

2.1	problem 1	24
2.2	problem 2	25
2.3	problem 3	26
2.4	problem 4	27
2.5	problem 5	28
2.6	problem 6	29
2.7	problem 7	30
2.8	problem 8	31
2.9	problem 9	32
2.10	problem 10	33
2.11	problem 11	34
2.12	problem 12	35
2.13	problem 13	36
2.14	problem 14	37
2.15	problem 15	38
2.16	problem 16	39
2.17	problem 17	40
2.18	problem 18	41
2.19	problem 19	42
2.20	problem 20	43
2.21	problem 21	44
2.22	problem 22	45
2.23	problem 23	46
2.24	problem 24	47
2.25	problem 25	48
2.26	problem 26	49
2.27	problem 27	50
2.28	problem 28	51
2.29	problem 29	52
2.30	problem 30	53
2.31	problem 31	54
2.32	problem 32	55
2.33	problem 33	56
2.34	problem 34	57
2.35	problem 35	58
2.36	problem 36	59

2.37	problem 44	60
2.38	problem 48(a)	61
2.39	problem 48(b)	62
2.40	problem 48(c)	63
2.41	problem 48(d)	64

2.1 problem 1

Internal problem ID [887]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' + ay = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x) + a*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-ax}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 19

```
DSolve[y'[x] + a*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-ax}$$

$$y(x) \rightarrow 0$$

2.2 problem 2

Internal problem ID [888]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + 3x^2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x) + 3*x^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x^3}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 20

```
DSolve[y'[x] + 3*x^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x^3}$$

$$y(x) \rightarrow 0$$

2.3 problem 3

Internal problem ID [889]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y'x + \ln(x)y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x) + ln(x)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{\ln(x)^2}{2}}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 23

```
DSolve[x*y'[x] +Log[x]*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\frac{1}{2} \log^2(x)}$$
$$y(x) \rightarrow 0$$

2.4 problem 4

Internal problem ID [890]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$3y + y'x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*diff(y(x),x) + 3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x^3}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 16

```
DSolve[x*y'[x] +3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^3}$$

$$y(x) \rightarrow 0$$

2.5 problem 5

Internal problem ID [891]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y'x^2 + y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve(x^2*diff(y(x),x) + y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{1}{x}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 18

```
DSolve[x^2*y'[x] + y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{1}{x}}$$

$$y(x) \rightarrow 0$$

2.6 problem 6

Internal problem ID [892]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \frac{(x+1)y}{x} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x) + ((1+x)/x)*y(x)=0,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e^{1-x}}{x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 16

```
DSolve[{y'[x] + ((1+x)/x)*y[x]==0,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{1-x}}{x}$$

2.7 problem 7

Internal problem ID [893]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'x + \left(1 + \frac{1}{\ln(x)}\right)y = 0$$

With initial conditions

$$[y(e) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x) + (1+1/ln(x))*y(x)=0,y(exp(1)) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e}{x \ln(x)}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 18

```
DSolve[{y'[x] + (1+1/Log[x])*y[x]==0,y[Exp[1]]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\text{LogIntegral}(x)+\text{LogIntegral}(e)-x+e}$$

2.8 problem 8

Internal problem ID [894]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y'x + (1 + x \cot(x))y = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 2 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([x*diff(y(x),x) + (1+x*cot(x))*y(x)=0,y(1/2*Pi) = 2],y(x), singsol=all)
```

$$y(x) = \frac{\csc(x) \pi}{x}$$

✓ Solution by Mathematica

Time used: 0.099 (sec). Leaf size: 66

```
DSolve[{y'[x] + (1+x*Cot[x])*y[x]==0,y[Pi/2]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2^{1+\frac{\pi}{2}} (1 - e^{2ix})^{-x} \exp\left(-\frac{1}{12}i(-6 \text{PolyLog}(2, e^{2ix}) - 6x(x + 2i) + \pi^2 + 6i\pi)\right)$$

2.9 problem 9

Internal problem ID [895]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2xy}{x^2 + 1} = 0$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([diff(y(x),x) - (2*x)/(1+x^2)*y(x)=0,y(0) = 2],y(x), singsol=all)
```

$$y(x) = 2x^2 + 2$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 12

```
DSolve[{y'[x] - (2*x)/(1+x^2)*y[x]==0,y[0]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2(x^2 + 1)$$

2.10 problem 10

Internal problem ID [896]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' + \frac{ky}{x} = 0$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x) +k/x*y(x)=0,y(1) = 3],y(x), singsol=all)
```

$$y(x) = 3x^{-k}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 12

```
DSolve[{y'[x] +k/x*y[x]==0,y[1]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3x^{-k}$$

2.11 problem 11

Internal problem ID [897]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \tan(kx)y = 0$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 18

```
dsolve([diff(y(x),x) +tan(k*x)*y(x)=0,y(0) = 2],y(x), singsol=all)
```

$$y(x) = 2(\sec(kx)^2)^{-\frac{1}{2k}}$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 15

```
DSolve[{y'[x] +Tan[k*x]*y[x]==0,y[0]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2\sqrt[k]{\cos(kx)}$$

2.12 problem 12

Internal problem ID [898]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$3y + y' = 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x) +3*y(x)=1,y(x), singsol=all)
```

$$y(x) = \frac{1}{3} + c_1 e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 24

```
DSolve[y'[x] +3*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} + c_1 e^{-3x}$$
$$y(x) \rightarrow \frac{1}{3}$$

2.13 problem 13

Internal problem ID [899]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \left(\frac{1}{x} - 1\right)y = -\frac{2}{x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x) +(1/x-1)*y(x)=-2/x,y(x), singsol=all)
```

$$y(x) = \frac{e^x c_1 + 2}{x}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 17

```
DSolve[y'[x] +(1/x-1)*y[x]==-2/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 + c_1 e^x}{x}$$

2.14 problem 14

Internal problem ID [900]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2yx + y' = x e^{-x^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x) +2*x*y(x)=x*exp(-x^2),y(x), singsol=all)
```

$$y(x) = \frac{(x^2 + 2c_1) e^{-x^2}}{2}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 24

```
DSolve[y' [x] +2*x*y[x]==x*Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x^2} (x^2 + 2c_1)$$

2.15 problem 15

Internal problem ID [901]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y' + \frac{2xy}{x^2 + 1} = \frac{e^{-x^2}}{x^2 + 1}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x), x) + (2*x)/(1+x^2)*y(x)=exp(-x^2)/(1+x^2), y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\pi} \operatorname{erf}(x) + 2c_1}{2x^2 + 2}$$

✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 28

```
DSolve[y'[x] + (2*x)/(1+x^2)*y[x]==Exp[-x^2]/(1+x^2), y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{\pi} \operatorname{erf}(x) + 2c_1}{2x^2 + 2}$$

2.16 problem 16

Internal problem ID [902]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{y}{x} = \frac{7}{x^2} + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x) +1/x*y(x)=7/x^2+3,y(x), singsol=all)
```

$$y(x) = \frac{\frac{3x^2}{2} + 7 \ln(x) + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 24

```
DSolve[y'[x] +1/x*y[x]==7/x^2+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x}{2} + \frac{7 \log(x)}{x} + \frac{c_1}{x}$$

2.17 problem 17

Internal problem ID [903]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{4y}{x-1} = \frac{1}{(x-1)^5} + \frac{\sin(x)}{(x-1)^4}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x) +4/(x-1)*y(x)=1/(x-1)^5+sin(x)/(x-1)^4,y(x), singsol=all)
```

$$y(x) = \frac{-\cos(x) + \ln(x-1) + c_1}{(x-1)^4}$$

✓ Solution by Mathematica

Time used: 0.07 (sec). Leaf size: 22

```
DSolve[y'[x] +4/(x-1)*y[x]==1/(x-1)^5+Sin[x]/(x-1)^4,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{\log(x-1) - \cos(x) + c_1}{(x-1)^4}$$

2.18 problem 18

Internal problem ID [904]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x + y(2x^2 + 1) = x^3e^{-x^2}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(x*diff(y(x),x) +(1+2*x^2)*y(x)=x^3*exp(-x^2),y(x), singsol=all)
```

$$y(x) = \frac{(x^4 + 4c_1)e^{-x^2}}{4x}$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 27

```
DSolve[x*y'[x] +(1+2*x^2)*y[x]==x^3*Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x^2}(x^4 + 4c_1)}{4x}$$

2.19 problem 19

Internal problem ID [905]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + y'x = \frac{2}{x^2} + 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x*diff(y(x),x) +2*y(x)=2/x^2+1,y(x), singsol=all)
```

$$y(x) = \frac{\frac{x^2}{2} + 2 \ln(x) + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 22

```
DSolve[x*y'[x] +2*y[x]==2/x^2+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 \log(x)}{x^2} + \frac{c_1}{x^2} + \frac{1}{2}$$

2.20 problem 20

Internal problem ID [906]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \tan(x)y = \cos(x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve(diff(y(x),x) +tan(x)*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = (c_1 + x) \cos(x)$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 12

```
DSolve[y'[x] +Tan[x]*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_1) \cos(x)$$

2.21 problem 21

Internal problem ID [907]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(x + 1)y' + 2y = \frac{\sin(x)}{x + 1}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve((1+x)*diff(y(x),x) +2*y(x)=sin(x)/(1+x),y(x), singsol=all)
```

$$y(x) = \frac{-\cos(x) + c_1}{(x + 1)^2}$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 18

```
DSolve[(1+x)*y'[x] +2*y[x]==Sin[x]/(1+x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-\cos(x) + c_1}{(x + 1)^2}$$

2.22 problem 22

Internal problem ID [908]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(-2 + x)(x - 1)y' - (4x - 3)y = (-2 + x)^3$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve((x-2)*(x-1)*diff(y(x),x) -(4*x-3)*y(x)=(x-2)^3,y(x), singsol=all)
```

$$y(x) = \frac{2\left(-\frac{1}{2} + (-2 + x)^2 c_1\right) (-2 + x)^3}{2x - 2}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 30

```
DSolve[(x-2)*(x-1)*y'[x] -(4*x-3)*y[x]==(x-2)^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(x - 2)^3 (-1 + 2c_1(x - 2)^2)}{2(x - 1)}$$

2.23 problem 23

Internal problem ID [909]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + 2 \cos(x) \sin(x) y = e^{-\sin(x)^2}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x) +(2*sin(x)*cos(x))*y(x)=exp(-sin(x)^2),y(x), singsol=all)
```

$$y(x) = (c_1 + x) e^{-\frac{1}{2} + \frac{\cos(2x)}{2}}$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 24

```
DSolve[y'[x] +(2*Sin[x]*Cos[x])*y[x]==Exp[-Sin[x]^2],y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow (x + \sqrt{e}c_1) e^{-\sin^2(x)}$$

2.24 problem 24

Internal problem ID [910]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y'x^2 + 3yx = e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x) +3*x*y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = \frac{(x - 1)e^x + c_1}{x^3}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 19

```
DSolve[x^2*y'[x] +3*x*y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x(x - 1) + c_1}{x^3}$$

2.25 problem 25

Internal problem ID [911]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_linear, 'class A']]`

$$y' + 7y = e^{3x}$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(y(x),x) +7*y(x)=exp(3*x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(e^{10x} - 1)e^{-7x}}{10}$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 21

```
DSolve[{y'[x] +7*y[x]==Exp[3*x],y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10}e^{-7x}(e^{10x} - 1)$$

2.26 problem 26

Internal problem ID [912]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$(x^2 + 1)y' + 4yx = \frac{2}{x^2 + 1}$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([(1+x^2)*diff(y(x),x)+4*x*y(x)=2/(1+x^2),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1 + 2x}{(x^2 + 1)^2}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 18

```
DSolve[{(1+x^2)*y'[x]+4*x*y[x]==2/(1+x^2),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x + 1}{(x^2 + 1)^2}$$

2.27 problem 27

Internal problem ID [913]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$3y + y'x = \frac{2}{x(x^2 + 1)}$$

With initial conditions

$$[y(-1) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve([x*diff(y(x),x)+3*y(x)=2/(x*(1+x^2)),y(-1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\ln(x^2 + 1) - \ln(2)}{x^3}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 19

```
DSolve[{x*y'[x]+3*y[x]==2/(x*(1+x^2)),y[-1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log\left(\frac{1}{2}(x^2 + 1)\right)}{x^3}$$

2.28 problem 28

Internal problem ID [914]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + y \cot(x) = \cos(x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 1 \right]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)+cot(x)*y(x)=cos(x),y(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x) \cot(x)}{2} + \csc(x)$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 16

```
DSolve[{y'[x]+Cot[x]*y[x]==Cos[x],y[Pi/2]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \csc(x) - \frac{1}{2} \cos(x) \cot(x)$$

2.29 problem 29

Internal problem ID [915]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' + \frac{y}{x} = \frac{2}{x^2} + 1$$

With initial conditions

$$[y(-1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve([diff(y(x),x)+y(x)/x=2/x^2+1,y(-1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{-4i\pi + x^2 + 4 \ln(x) - 1}{2x}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 26

```
DSolve[{y'[x]+y[x]/x==2/x^2+1,y[-1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 + 4 \log(x) - 4i\pi - 1}{2x}$$

2.30 problem 30

Internal problem ID [916]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(x - 1)y' + 3y = \frac{1}{(x - 1)^3} + \frac{\sin(x)}{(x - 1)^2}$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve([(x-1)*diff(y(x),x)+3*y(x)=1/(x-1)^3+sin(x)/(x-1)^2,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{-\cos(x) + \ln(x - 1) - i\pi}{(x - 1)^3}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 25

```
DSolve[{(x-1)*y'[x]+3*y[x]==1/(x-1)^3+Sin[x]/(x-1)^2,y[0]==1},y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow \frac{\log(x - 1) - \cos(x) - i\pi}{(x - 1)^3}$$

2.31 problem 31

Internal problem ID [917]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$2y + y'x = 8x^2$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x*diff(y(x),x)+2*y(x)=8*x^2,y(1) = 3],y(x), singsol=all)
```

$$y(x) = \frac{2x^4 + 1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 14

```
DSolve[{x*y'[x]+2*y[x]==8*x^2,y[1]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^2 + \frac{1}{x^2}$$

2.32 problem 32

Internal problem ID [918]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$y'x - 2y = -x^2$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x)-2*y(x)=-x^2,y(1) = 1],y(x), singsol=all)
```

$$y(x) = (1 - \ln(x))x^2$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 14

```
DSolve[{x*y'[x]-2*y[x]==-x^2,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^2(\log(x) - 1)$$

2.33 problem 33

Internal problem ID [919]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 33.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2yx + y' = x$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+2*x*y(x)=x,y(0) = 3],y(x), singsol=all)
```

$$y(x) = \frac{1}{2} + \frac{5e^{-x^2}}{2}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 20

```
DSolve[{y'[x]+2*x*y[x]==x,y[0]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5e^{-x^2}}{2} + \frac{1}{2}$$

2.34 problem 34

Internal problem ID [920]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 34.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(x - 1)y' + 3y = \frac{1 + (x - 1)\sec(x)^2}{(x - 1)^3}$$

With initial conditions

$$[y(0) = -1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 22

```
dsolve([(x-1)*diff(y(x),x)+3*y(x)=(1+(x-1)*sec(x)^2)/(x-1)^3,y(0) = -1],y(x), singsol=all)
```

$$y(x) = \frac{\ln(x - 1) + \tan(x) + 1 - i\pi}{(x - 1)^3}$$

✓ Solution by Mathematica

Time used: 0.171 (sec). Leaf size: 21

```
DSolve[{(x-1)*y'[x]+3*y[x]==(1+(x-1)*Sec[x]^2)/(x-1)^3,y[0]==-1},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{\log(1 - x) + \tan(x) + 1}{(x - 1)^3}$$

2.35 problem 35

Internal problem ID [921]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 35.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(2+x)y' + 4y = \frac{2x^2 + 1}{x(2+x)^3}$$

With initial conditions

$$[y(-1) = 2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve([(x+2)*diff(y(x),x)+4*y(x) = (1+2*x^2)/(x*(x+2)^3), y(-1) = 2], y(x), singsol=all)
```

$$y(x) = \frac{x^2 + \ln(x) + 1 - i\pi}{(2+x)^4}$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 23

```
DSolve[{(x+2)*y'[x]+4*y[x]==(1+2*x^2)/(x*(x+2)^3), y[-1]==2}, y[x], x, IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{x^2 + \log(x) - i\pi + 1}{(x+2)^4}$$

2.36 problem 36

Internal problem ID [922]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 36.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$(x^2 - 1) y' - 2yx = x(x^2 - 1)$$

With initial conditions

$$[y(0) = 4]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 29

```
dsolve([(x^2-1)*diff(y(x),x)-2*x*y(x)= x*(x^2-1),y(0) = 4],y(x), singsol=all)
```

$$y(x) = -\frac{(i\pi - \ln(x+1) - \ln(x-1) + 8)(x^2 - 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 27

```
DSolve[{(x^2-1)*y'[x]-2*x*y[x]== x*(x^2-1),y[0]==4},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x^2 - 1) (\log(x^2 - 1) - i\pi - 8)$$

2.37 problem 44

Internal problem ID [923]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 44.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'x - 2y = -1$$

With initial conditions

$$\left[y(0) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 11

```
dsolve([x*diff(y(x),x)-2*y(x)=-1,y(0) = 1/2],y(x), singsol=all)
```

$$y(x) = \frac{1}{2} + c_1x^2$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 8

```
DSolve[{x*y'[x]-2*y[x]== -1,y[0]==1/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}$$

2.38 problem 48(a)

Internal problem ID [924]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 48(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$\sec(y)^2 y' - 3 \tan(y) = -1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(sec(y(x))^2*diff(y(x),x)-3*tan(y(x))=-1,y(x), singsol=all)
```

$$y(x) = \arctan\left(\frac{c_1 e^{3x}}{3} + \frac{1}{3}\right)$$

✓ Solution by Mathematica

Time used: 60.217 (sec). Leaf size: 177

```
DSolve[Sec[y[x]]^2*y'[x]-3*Tan[y[x]]== -1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(\frac{3e^{6c_1}}{\sqrt{e^{6x} - 2e^{3x+6c_1} + 10e^{12c_1}}}\right)$$

$$y(x) \rightarrow \arccos\left(\frac{3e^{6c_1}}{\sqrt{e^{6x} - 2e^{3x+6c_1} + 10e^{12c_1}}}\right)$$

$$y(x) \rightarrow -\arccos\left(\frac{3e^{6c_1}}{\sqrt{e^{6x} - 2e^{3x+6c_1} + 10e^{12c_1}}}\right)$$

$$y(x) \rightarrow \arccos\left(\frac{3e^{6c_1}}{\sqrt{e^{6x} - 2e^{3x+6c_1} + 10e^{12c_1}}}\right)$$

2.39 problem 48(b)

Internal problem ID [925]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 48(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(y)]']]`

$$e^{y^2} \left(2yy' + \frac{2}{x} \right) = \frac{1}{x^2}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve(exp(y(x)^2)*(2*y(x)*diff(y(x),x)+2/x)= 1/x^2,y(x), singsol=all)
```

$$y(x) = \sqrt{\ln\left(\frac{-c_1 + x}{x^2}\right)}$$
$$y(x) = -\sqrt{\ln\left(\frac{-c_1 + x}{x^2}\right)}$$

✓ Solution by Mathematica

Time used: 7.286 (sec). Leaf size: 37

```
DSolve[Exp[y[x]^2]*(2*y[x]*y'[x]+2/x)== 1/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\log\left(\frac{x + c_1}{x^2}\right)}$$
$$y(x) \rightarrow \sqrt{\log\left(\frac{x + c_1}{x^2}\right)}$$

2.40 problem 48(c)

Internal problem ID [926]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 48(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)*y+H(x)]']]`

$$2 \ln(y) = -\frac{xy'}{y} + 4x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)/y(x)+2*ln(y(x))= 4*x^2,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^4 - c_1}{x^2}}$$

✓ Solution by Mathematica

Time used: 0.248 (sec). Leaf size: 17

```
DSolve[x*y'[x]/y[x]+2*Log[y[x]]== 4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x^2 + \frac{c_1}{x^2}}$$

2.41 problem 48(d)

Internal problem ID [927]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Linear first order. Section 2.1 Page 41

Problem number: 48(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, _Riccati]`

$$\frac{y'}{(1+y)^2} - \frac{1}{x(1+y)} = -\frac{3}{x^2}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)/(1+y(x))^2-1/(x*(1+y(x))))=-3/x^2,y(x), singsol=all)
```

$$y(x) = -1 + \frac{x}{3 \ln(x) + 3c_1}$$

✓ Solution by Mathematica

Time used: 0.252 (sec). Leaf size: 31

```
DSolve[y'[x]/(1+y[x])^2-1/(x*(1+y[x]))== -3/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x - 3 \log(x) - 3c_1}{3(\log(x) + c_1)}$$
$$y(x) \rightarrow -1$$

3 Chapter 2, First order equations. separable equations. Section 2.2 Page 52

3.1	problem 1	66
3.2	problem 2	67
3.3	problem 3	68
3.4	problem 5	69
3.5	problem 6	70
3.6	problem 7	71
3.7	problem 8	72
3.8	problem 9	73
3.9	problem 10	74
3.10	problem 11	75
3.11	problem 12	76
3.12	problem 13	77
3.13	problem 14	79
3.14	problem 15	81
3.15	problem 16	82
3.16	problem 17	83
3.17	problem 18	84
3.18	problem 19	85
3.19	problem 20	86
3.20	problem 21	87
3.21	problem 22	88
3.22	problem 23	89
3.23	problem 24	90
3.24	problem 25	91
3.25	problem 26	92
3.26	problem 27	93
3.27	problem 35	94
3.28	problem 36	95
3.29	problem 37	96
3.30	problem 38	97

3.1 problem 1

Internal problem ID [928]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{3x^2 + 2x + 1}{y - 2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 53

```
dsolve(diff(y(x),x)= (3*x^2+2*x+1)/(y(x)-2),y(x), singsol=all)
```

$$y(x) = 2 - \sqrt{2x^3 + 2x^2 + 2c_1 + 2x + 4}$$
$$y(x) = 2 + \sqrt{2x^3 + 2x^2 + 2c_1 + 2x + 4}$$

✓ Solution by Mathematica

Time used: 0.158 (sec). Leaf size: 56

```
DSolve[y'[x]== (3*x^2+2*x+1)/(y[x]-2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 - \sqrt{2}\sqrt{x^3 + x^2 + x + 2 + c_1}$$
$$y(x) \rightarrow 2 + \sqrt{2}\sqrt{x^3 + x^2 + x + 2 + c_1}$$

3.2 problem 2

Internal problem ID [929]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\sin(y) \sin(x) + \cos(y) y' = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 12

```
dsolve(sin(x)*sin(y(x))+cos(y(x))*diff(y(x),x)= 0,y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{e^{\cos(x)}}{c_1}\right)$$

✓ Solution by Mathematica

Time used: 29.953 (sec). Leaf size: 22

```
DSolve[Sin[x]*Sin[y[x]]+Cos[y[x]]*y'[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(e^{\cos(x)+\frac{c_1}{2}}\right)$$
$$y(x) \rightarrow 0$$

3.3 problem 3

Internal problem ID [930]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'x + y^2 + y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve(x*diff(y(x),x)+y(x)^2+y(x)= 0,y(x), singsol=all)
```

$$y(x) = \frac{1}{c_1x - 1}$$

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 31

```
DSolve[x*y'[x]+y[x]^2+y[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow -\frac{e^{c_1}}{-x + e^{c_1}} \\y(x) &\rightarrow -1 \\y(x) &\rightarrow 0\end{aligned}$$

3.4 problem 5

Internal problem ID [931]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(3y^3 + 3y \cos(y) + 1) y' + \frac{(1 + 2x)y}{x^2 + 1} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve((3*y(x)^3+3*y(x)*cos(y(x))+1)*diff(y(x),x)+((2*x+1)*y(x))/(1+x^2)= 0,y(x), singsol=al
```

$$\ln(x^2 + 1) + \arctan(x) + y(x)^3 + 3 \sin(y(x)) + \ln(y(x)) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.379 (sec). Leaf size: 40

```
DSolve[(3*y[x]^3+3*y[x]*Cos[y[x]]+1)*y'[x]+((2*x+1)*y[x])/(1+x^2)== 0,y[x],x,IncludeSingular
```

$$y(x) \rightarrow \text{InverseFunction}[\#1^3 + \log(\#1) + 3 \sin(\#1)\&] [-\arctan(x) - \log(x^2 + 1) + c_1]$$

$$y(x) \rightarrow 0$$

3.5 problem 6

Internal problem ID [932]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y'x^2y - (-1 + y^2)^{\frac{3}{2}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(x^2*y(x)*diff(y(x),x)= (y(x)^2-1)^(3/2),y(x), singsol=all)
```

$$-\frac{1}{x} + \frac{(y(x) - 1)(y(x) + 1)}{(y(x)^2 - 1)^{\frac{3}{2}}} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.707 (sec). Leaf size: 111

```
DSolve[x^2*y[x]*y'[x]== (y[x]^2-1)^(3/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{(1 + c_1^2)x^2 - 2c_1x + 1}}{1 - c_1x}$$

$$y(x) \rightarrow \frac{\sqrt{(1 + c_1^2)x^2 - 2c_1x + 1}}{-1 + c_1x}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow -\frac{\sqrt{x^2}}{x}$$

$$y(x) \rightarrow \frac{\sqrt{x^2}}{x}$$

3.6 problem 7

Internal problem ID [933]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - (1 + y^2) x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)= x^2*(1+y(x)^2),y(x), singsol=all)
```

$$y(x) = \tan\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.169 (sec). Leaf size: 30

```
DSolve[y'[x]== x^2*(1+y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan\left(\frac{x^3}{3} + c_1\right)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

3.7 problem 8

Internal problem ID [934]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^2 + 1) y' + yx = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)*(1+x^2)+x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{\sqrt{x^2 + 1}}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 22

```
DSolve[y'[x]*(1+x^2)+x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{\sqrt{x^2 + 1}}$$
$$y(x) \rightarrow 0$$

3.8 problem 9

Internal problem ID [935]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - (x - 1)(y - 1)(y - 2) = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)=(x-1)*(y(x)-1)*(y(x)-2),y(x), singsol=all)
```

$$y(x) = \frac{-2 + c_1 e^{\frac{x(-2+x)}{2}}}{c_1 e^{\frac{x(-2+x)}{2}} - 1}$$

✓ Solution by Mathematica

Time used: 0.308 (sec). Leaf size: 56

```
DSolve[y'[x]==(x-1)*(y[x]-1)*(y[x]-2),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow \frac{2e^x - e^{\frac{x^2}{2}+c_1}}{e^x - e^{\frac{x^2}{2}+c_1}} \\y(x) &\rightarrow 1 \\y(x) &\rightarrow 2\end{aligned}$$

3.9 problem 10

Internal problem ID [936]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(y - 1)^2 y' = 2x + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 102

```
dsolve((y(x)-1)^2*diff(y(x),x)=2*x+3,y(x), singsol=all)
```

$$y(x) = (3x^2 + 3c_1 + 9x)^{\frac{1}{3}} + 1$$
$$y(x) = -\frac{(3x^2 + 3c_1 + 9x)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3}(3x^2 + 3c_1 + 9x)^{\frac{1}{3}}}{2} + 1$$
$$y(x) = -\frac{(3x^2 + 3c_1 + 9x)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3}(3x^2 + 3c_1 + 9x)^{\frac{1}{3}}}{2} + 1$$

✓ Solution by Mathematica

Time used: 0.482 (sec). Leaf size: 103

```
DSolve[(y[x]-1)^2*y'[x]==2*x+3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 + \sqrt[3]{3x^2 + 9x - 1 + 3c_1}$$
$$y(x) \rightarrow 1 + \frac{1}{2}i(\sqrt{3} + i)\sqrt[3]{3x^2 + 9x - 1 + 3c_1}$$
$$y(x) \rightarrow 1 - \frac{1}{2}(1 + i\sqrt{3})\sqrt[3]{3x^2 + 9x - 1 + 3c_1}$$

3.10 problem 11

Internal problem ID [937]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{x^2 + 3x + 2}{y - 2} = 0$$

With initial conditions

$$[y(1) = 4]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 25

```
dsolve([diff(y(x),x)=(x^2+3*x+2)/(y(x)-2),y(1) = 4],y(x), singsol=all)
```

$$y(x) = 2 + \frac{\sqrt{6x^3 + 27x^2 + 36x - 33}}{3}$$

✓ Solution by Mathematica

Time used: 0.167 (sec). Leaf size: 30

```
DSolve[{y'[x]==(x^2+3*x+2)/(y[x]-2),y[1]==4},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{\frac{2x^3}{3} + 3x^2 + 4x - \frac{11}{3}} + 2$$

3.11 problem 12

Internal problem ID [938]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + x(y^2 + y) = 0$$

With initial conditions

$$[y(2) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve([diff(y(x),x)+x*(y(x)^2+y(x))=0,y(2) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{-1 + 2e^{\frac{(2+x)(-2+x)}{2}}}$$

✓ Solution by Mathematica

Time used: 0.241 (sec). Leaf size: 27

```
DSolve[{y'[x]+x*(y[x]^2+y[x])==0,y[2]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^2}{e^2 - 2e^{\frac{x^2}{2}}}$$

3.12 problem 13

Internal problem ID [939]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$(3y^2 + 4y) y' = -2x - \cos(x)$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.422 (sec). Leaf size: 102

```
dsolve([(3*y(x)^2+4*y(x))*diff(y(x),x)+2*x+cos(x)=0,y(0) = 1],y(x), singsol=all)
```

$y(x)$

$$= \frac{\left(260 - 108x^2 - 108 \sin(x) + 12\sqrt{441 - 390x^2 - 390 \sin(x) + 81x^4 + 162 \sin(x)x^2 + 81 \sin(x)^2}\right)^{\frac{1}{3}}}{6} + \frac{3 \left(260 - 108x^2 - 108 \sin(x) + 12\sqrt{441 - 390x^2 - 390 \sin(x) + 81x^4 + 162 \sin(x)x^2 + 81 \sin(x)^2}\right)^{\frac{1}{3}}}{8} - \frac{2}{3}$$

✓ Solution by Mathematica

Time used: 2.549 (sec). Leaf size: 127

```
DSolve[{(3*y[x]^2+4*y[x])*y'[x]+2*x+Cos[x]==0,y[0]==1},y[x],x,IncludeSingularSolutions -> Tr
```

$y(x)$

$$\rightarrow \frac{2^{2/3} \left(-27x^2 + \sqrt{(27x^2 + 27 \sin(x) - 65)^2 - 256 - 27 \sin(x) + 65} \right)^{2/3} - 4 \sqrt[3]{-27x^2 + \sqrt{(27x^2 + 27 \sin(x) - 65)^2 - 256 - 27 \sin(x) + 65}}}{6 \sqrt[3]{-27x^2 + \sqrt{(27x^2 + 27 \sin(x) - 65)^2 - 256 - 27 \sin(x) + 65}}}$$

3.13 problem 14

Internal problem ID [940]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + \frac{(1+y)(y-1)(y-2)}{x+1} = 0$$

With initial conditions

$$[y(1) = 0]$$

✓ Solution by Maple

Time used: 6.86 (sec). Leaf size: 633

```
dsolve([diff(y(x),x)+((y(x)+1)*(y(x)-1)*(y(x)-2))/(x+1)=0,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{8 \operatorname{RootOf}\left(512_Z^6 + \left(96 \cdot 2^{\frac{1}{3}} x^2 + 192 x 2^{\frac{1}{3}} + 96 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}{8 \operatorname{RootOf}\left(512_Z^6 + \left(96 \cdot 2^{\frac{1}{3}} x^2 + 192 x 2^{\frac{1}{3}} + 96 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}$$

$$y(x) = \frac{16 \operatorname{RootOf}\left(512_Z^6 + \left(48ix^2\sqrt{3} \cdot 2^{\frac{1}{3}} + 96ix\sqrt{3} \cdot 2^{\frac{1}{3}} + 48i\sqrt{3} \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}} x^2 - 96x \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}{16 \operatorname{RootOf}\left(512_Z^6 + \left(48ix^2\sqrt{3} \cdot 2^{\frac{1}{3}} + 96ix\sqrt{3} \cdot 2^{\frac{1}{3}} + 48i\sqrt{3} \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}} x^2 - 96x \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}$$

$$y(x) = \frac{16 \operatorname{RootOf}\left(512_Z^6 + \left(-48ix^2\sqrt{3} \cdot 2^{\frac{1}{3}} - 96ix\sqrt{3} \cdot 2^{\frac{1}{3}} - 48i\sqrt{3} \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}} x^2 - 96x \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}{16 \operatorname{RootOf}\left(512_Z^6 + \left(-48ix^2\sqrt{3} \cdot 2^{\frac{1}{3}} - 96ix\sqrt{3} \cdot 2^{\frac{1}{3}} - 48i\sqrt{3} \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}} x^2 - 96x \cdot 2^{\frac{1}{3}} - 48 \cdot 2^{\frac{1}{3}}\right) _Z^4 - x^6 - 6x^5 - 15x^4 - 20x^3 - 15x^2 - 6x - 257\right)}$$

✓ Solution by Mathematica

Time used: 60.912 (sec). Leaf size: 1618

```
DSolve[{y'[x]+(y[x]+1)*(y[x]-1)*(y[x]-2)/(x+1)==0,y[1]==0},y[x],x,IncludeSingularSolutions
```

$y(x)$

$$(-1 - i\sqrt{3})x^{12} - 12i(-i + \sqrt{3})x^{11} - 66i(-i + \sqrt{3})x^{10} - 220i(-i + \sqrt{3})x^9 - 495i(-i + \sqrt{3})x^8 - 7$$

→

3.14 problem 15

Internal problem ID [941]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + 2x(1 + y) = 0$$

With initial conditions

$$[y(0) = 2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+2*x*(y(x)+1)=0,y(0) = 2],y(x), singsol=all)
```

$$y(x) = -1 + 3e^{-x^2}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 16

```
DSolve[{y'[x]+2*x*(y[x]+1)==0,y[0]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3e^{-x^2} - 1$$

3.15 problem 16

Internal problem ID [942]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 2xy(1 + y^2) = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)=2*x*y(x)*(1+y(x)^2),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{2e^{-2x^2} - 1}}$$

✓ Solution by Mathematica

Time used: 60.086 (sec). Leaf size: 27

```
DSolve[{y'[x]==2*x*y[x]*(1+y[x]^2),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{ie^{x^2}}{\sqrt{e^{2x^2} - 2}}$$

3.16 problem 17

Internal problem ID [943]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x^2 + 2)y' - 4x(y^2 + 2y + 1) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve(diff(y(x),x)*(x^2+2)=4*x*(y(x)^2+2*y(x)+1),y(x), singsol=all)
```

$$y(x) = \frac{-2 \ln(x^2 + 2) - 4c_1 - 1}{2 \ln(x^2 + 2) + 4c_1}$$

✓ Solution by Mathematica

Time used: 0.195 (sec). Leaf size: 37

```
DSolve[y'[x]*(x^2+2)==4*x*(y[x]^2+2*y[x]+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2 \log(x^2 + 2) + 1 + c_1}{2 \log(x^2 + 2) + c_1}$$
$$y(x) \rightarrow -1$$

3.17 problem 18

Internal problem ID [944]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + 2x(y^3 - 3y + 2) = 0$$

With initial conditions

$$[y(0) = 3]$$

✓ Solution by Maple

Time used: 0.375 (sec). Leaf size: 70

```
dsolve([diff(y(x),x)=-2*x*(y(x)^3-3*y(x)+2),y(0) = 3],y(x), singsol=all)
```

$y(x)$

$$= e^{\text{RootOf}(18x^2e^{-Z}-2\ln(e^{-Z}-3)e^{-Z}+2e^{-Z}\ln(2)-2e^{-Z}\ln(5)+2_Ze^{-Z}-54x^2+3e^{-Z}+6\ln(e^{-Z}-3)-6\ln(2)+6\ln(5)-6_Z-15)} - 2$$

✓ Solution by Mathematica

Time used: 1.071 (sec). Leaf size: 49

```
DSolve[{y'[x]==-2*x*(y[x]^3-3*y[x]+2),y[0]==3},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{1}{9} \left(-\frac{3}{\#1-1} - \log(\#1-1) + \log(\#1+2) \right) \&t \right] \left[-x^2 - \frac{1}{6} + \frac{1}{9} \log \left(\frac{5}{2} \right) \right]$$

3.18 problem 19

Internal problem ID [945]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{2x}{1+2y} = 0$$

With initial conditions

$$[y(2) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)=2*x/(1+2*y(x)),y(2) = 0],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2} + \frac{\sqrt{4x^2 - 15}}{2}$$

✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 22

```
DSolve[{y'[x]==2*x/(1+2*y[x]),y[2]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{4x^2 - 15} - 1 \right)$$

3.19 problem 20

Internal problem ID [946]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 2y + y^2 = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)=2*y(x)-y(x)^2,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{2}{1 + e^{-2x}}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 21

```
DSolve[{y'[x]==2*y[x]-y[x]^2,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{2x}}{e^{2x} + 1}$$

3.20 problem 21

Internal problem ID [947]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$yy' = -x$$

With initial conditions

$$[y(3) = -4]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 15

```
dsolve([x+y(x)*diff(y(x),x)=0,y(3) = -4],y(x), singsol=all)
```

$$y(x) = -\sqrt{-x^2 + 25}$$

✓ Solution by Mathematica

Time used: 0.083 (sec). Leaf size: 18

```
DSolve[{x+y[x]*y'[x]==0,y[3]==-4},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{25 - x^2}$$

3.21 problem 22

Internal problem ID [948]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' + x^2(1 + y)(y - 2)^2 = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 35

```
dsolve(diff(y(x),x)+x^2*(y(x)+1)*(y(x)-2)^2=0,y(x), singsol=all)
```

$$y(x) = e^{\text{RootOf}(3x^3e^{-Z} + \ln(e^{-Z} + 3)e^{-Z} + 9c_1e^{-Z} - Ze^{-Z} - 3)} + 2$$

✓ Solution by Mathematica

Time used: 0.482 (sec). Leaf size: 52

```
DSolve[y'[x]+x^2*(y[x]+1)*(y[x]-2)^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{1}{9} \left(-\frac{3}{\#1 - 2} - \log(\#1 - 2) + \log(\#1 + 1) \right) \& \right] \left[-\frac{x^3}{3} + c_1 \right]$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 2$$

3.22 problem 23

Internal problem ID [949]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x + 1)(-2 + x)y' + y = 0$$

With initial conditions

$$[y(1) = -3]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 22

```
dsolve([(x+1)*(x-2)*diff(y(x),x)+y(x)=0,y(1) = -3],y(x), singsol=all)
```

$$y(x) = -\frac{3 \cdot 2^{\frac{2}{3}} (1 + i\sqrt{3}) (x + 1)^{\frac{1}{3}}}{4 (-2 + x)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 23

```
DSolve[{(x+1)*(x-2)*y'[x]+y[x]==0,y[1]==-3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3\sqrt[3]{x+1}}{\sqrt[3]{4-2x}}$$

3.23 problem 24

Internal problem ID [950]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{1 + y^2}{x^2 + 1} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=(1+y(x)^2)/(1+x^2),y(x), singsol=all)
```

$$y(x) = \tan(\arctan(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.227 (sec). Leaf size: 25

```
DSolve[y'[x]==(1+y[x]^2)/(1+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(\arctan(x) + c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

3.24 problem 25

Internal problem ID [951]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' \sqrt{-x^2 + 1} + \sqrt{1 - y^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)*sqrt(1-x^2)+sqrt(1-y(x)^2)=0,y(x), singsol=all)
```

$$y(x) = -\sin(\arcsin(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.317 (sec). Leaf size: 47

```
DSolve[y'[x]*Sqrt[1-x^2]+Sqrt[1-y[x]^2]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos\left(2 \arctan\left(\frac{\sqrt{1-x^2}}{x+1}\right) + c_1\right)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

3.25 problem 26

Internal problem ID [952]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - \frac{\cos(x)}{\sin(y)} = 0$$

With initial conditions

$$\left[y(\pi) = \frac{\pi}{2} \right]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)=cos(x)/sin(y(x)),y(Pi) = 1/2*Pi],y(x), singsol=all)
```

$$y(x) = \frac{\pi}{2} + \arcsin(\sin(x))$$

✓ Solution by Mathematica

Time used: 0.439 (sec). Leaf size: 10

```
DSolve[{y'[x]==Cos[x]/Sin[y[x]],y[Pi]==Pi/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arccos(-\sin(x))$$

3.26 problem 27

Internal problem ID [953]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - ay + by^2 = 0$$

With initial conditions

$$[y(0) = y_0]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 25

```
dsolve([diff(y(x),x)=a*y(x)-b*y(x)^2,y(0) = y0],y(x), singsol=all)
```

$$y(x) = \frac{a y_0}{(-y_0 b + a) e^{-ax} + y_0 b}$$

✓ Solution by Mathematica

Time used: 0.854 (sec). Leaf size: 27

```
DSolve[{y'[x]==a*y[x]-b*y[x]^2,y[0]==y0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{a y_0 e^{ax}}{b y_0 (e^{ax} - 1) + a}$$

3.27 problem 35

Internal problem ID [954]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 35.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Abel, '2nd type', 'class B']]`

$$y' + y - \frac{2x e^{-x}}{1 + e^x y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
dsolve(diff(y(x),x)+y(x)=(2*x*exp(-x))/(1+y(x)*exp(x)),y(x), singsol=all)
```

$$y(x) = \left(-1 - \sqrt{2x^2 - 2c_1 + 1}\right) e^{-x}$$

$$y(x) = \left(-1 + \sqrt{2x^2 - 2c_1 + 1}\right) e^{-x}$$

✓ Solution by Mathematica

Time used: 32.427 (sec). Leaf size: 70

```
DSolve[y'[x]+y[x]==(2*x*Exp[-x])/(1+y[x]*Exp[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^{-2x} \left(e^x + \sqrt{e^{2x} (2x^2 + 1 + c_1)} \right)$$

$$y(x) \rightarrow e^{-2x} \left(-e^x + \sqrt{e^{2x} (2x^2 + 1 + c_1)} \right)$$

3.28 problem 36

Internal problem ID [955]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 36.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$y'x - 2y - \frac{x^6}{x^2 + y} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 41

```
dsolve(x*diff(y(x),x)-2*y(x)=x^6/(y(x)+x^2),y(x), singsol=all)
```

$$y(x) = \left(-1 - \sqrt{x^2 - 2c_1 + 1}\right) x^2$$

$$y(x) = \left(-1 + \sqrt{x^2 - 2c_1 + 1}\right) x^2$$

✓ Solution by Mathematica

Time used: 0.635 (sec). Leaf size: 70

```
DSolve[x*y'[x]-2*y[x]==x^6/(y[x]+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^2 \left(1 + \sqrt{\frac{1}{x^5} x^2 \sqrt{x(x^2 + 1 + c_1)}}\right)$$

$$y(x) \rightarrow -x^2 + \sqrt{\frac{1}{x^5} x^4 \sqrt{x(x^2 + 1 + c_1)}}$$

3.29 problem 37

Internal problem ID [956]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 37.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(x)*y+H(x)]']]`

$$y' - y - \frac{(x+1)e^{4x}}{(y+e^x)^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 92

```
dsolve(diff(y(x),x)-y(x)=((x+1)*exp(4*x))/(y(x)+exp(x))^2,y(x), singsol=all)
```

$$y(x) = e^x \left(-1 + (3x e^x - 3c_1 + 1)^{\frac{1}{3}} \right)$$
$$y(x) = - \frac{e^x \left(i\sqrt{3} (3x e^x - 3c_1 + 1)^{\frac{1}{3}} + (3x e^x - 3c_1 + 1)^{\frac{1}{3}} + 2 \right)}{2}$$
$$y(x) = \frac{e^x (i\sqrt{3} - 1) (3x e^x - 3c_1 + 1)^{\frac{1}{3}}}{2} - e^x$$

✓ Solution by Mathematica

Time used: 19.018 (sec). Leaf size: 143

```
DSolve[y'[x]-y[x]==((x+1)*Exp[4*x])/(y[x]+Exp[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -e^x + e^{3x} \sqrt[3]{e^{-6x} (3e^x x + 1 + 3c_1)}$$
$$y(x) \rightarrow -e^x + \frac{1}{2} i \left(\sqrt{3} + i \right) e^{3x} \sqrt[3]{e^{-6x} (3e^x x + 1 + 3c_1)}$$
$$y(x) \rightarrow -e^x - \frac{1}{2} \left(1 + i\sqrt{3} \right) e^{3x} \sqrt[3]{e^{-6x} (3e^x x + 1 + 3c_1)}$$

3.30 problem 38

Internal problem ID [957]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. separable equations. Section 2.2 Page 52

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Abel, '2nd type', 'class A']]`

$$y' - 2y - \frac{x e^{2x}}{1 - y e^{-2x}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 60

```
dsolve(diff(y(x),x)-2*y(x)=x*exp(2*x)/(1-y(x)*exp(-2*x)),y(x), singsol=all)
```

$$y(x) = e^{4x} \sqrt{-e^{-4x} (x^2 + 2c_1 - 1)} + e^{2x}$$
$$y(x) = -e^{4x} \sqrt{-e^{-4x} (x^2 + 2c_1 - 1)} + e^{2x}$$

✓ Solution by Mathematica

Time used: 0.777 (sec). Leaf size: 72

```
DSolve[y'[x]-2*y[x]==x*Exp[2*x]/(1-y[x]*Exp[-2*x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} - \frac{\sqrt{x^2 - 1 - c_1}}{\sqrt{-e^{-4x}}}$$
$$y(x) \rightarrow e^{2x} + \frac{\sqrt{x^2 - 1 - c_1}}{\sqrt{-e^{-4x}}}$$

4 Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations.

Section 2.3 Page 60

4.1	problem 1	99
4.2	problem 2	100
4.3	problem 3	101
4.4	problem 4	102
4.5	problem 5	103
4.6	problem 6	104
4.7	problem 7	105
4.8	problem 8	106
4.9	problem 9	107
4.10	problem 10	108
4.11	problem 11	109
4.12	problem 12	110
4.13	problem 13	111
4.14	problem 16	112
4.15	problem 18	113
4.16	problem 19	114
4.17	problem 20(a)	115

4.1 problem 1

Internal problem ID [958]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Riccati]

$$y' - \frac{x^2 + y^2}{\sin(x)} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)/sin(x),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x^2+y[x]^2)/Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.2 problem 2

Internal problem ID [959]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type ['y=_G(x,y)']

$$y' - \frac{y + e^x}{x^2 + y^2} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(exp(x)+y(x))/(x^2+y(x)^2),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(Exp[x]+y[x])/(x^2+y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.3 problem 3

Internal problem ID [960]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type ['y=_G(x,y)']

$$y' - \tan(yx) = 0$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 44

```
dsolve(diff(y(x),x)=tan(x*y(x)),y(x), singsol=all)
```

$$y(x) = -i \operatorname{RootOf} \left(-\operatorname{erf} \left(\frac{(-x + _Z) \sqrt{2}}{2} \right) \sqrt{\pi} - \sqrt{\pi} \operatorname{erf} \left(\frac{\sqrt{2}(x + _Z)}{2} \right) + \sqrt{2} c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.311 (sec). Leaf size: 69

```
DSolve[y'[x]==Tan[x*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$\operatorname{Solve} \left[\frac{1}{2} \sqrt{\frac{\pi}{2}} e^{\frac{x^2}{2}} \left(\operatorname{erfi} \left(\frac{y(x) - ix}{\sqrt{2}} \right) + \operatorname{erfi} \left(\frac{y(x) + ix}{\sqrt{2}} \right) \right) = c_1 e^{\frac{x^2}{2}}, y(x) \right]$$

4.4 problem 4

Internal problem ID [961]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type ['y=_G(x,y)']

$$y' - \frac{x^2 + y^2}{\ln(yx)} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)/ln(x*y(x)),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x^2+y[x]^2)/Log[x*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.5 problem 5

Internal problem ID [962]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [$y' = G(x, y)$]

$$y' - (x^2 + y^2) y^{\frac{1}{3}} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)*y(x)^(1/3),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x^2+y[x]^2)*y[x]^(1/3),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.6 problem 6

Internal problem ID [963]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$-2yx + y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=2*x*y(x),y(x), singsol=all)
```

$$y(x) = e^{x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 18

```
DSolve[y'[x]==2*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{x^2}$$

$$y(x) \rightarrow 0$$

4.7 problem 7

Internal problem ID [964]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [$y = G(x, y')$]

$$y' - \ln(1 + x^2 + y^2) = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=ln(1+x^2+y(x)^2),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==Log[1+x^2+y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.8 problem 8

Internal problem ID [965]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{2x + 3y}{x - 4y} = 0$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 51

```
dsolve(diff(y(x),x)=(2*x+3*y(x))/(x-4*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{x(\sqrt{7} \tan(\text{RootOf}(\sqrt{7} \ln(\sec(_Z)^2 x^2) + \sqrt{7} \ln(7) - 3\sqrt{7} \ln(2) + 2\sqrt{7} c_1 - 4_Z)) - 1)}{4}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 53

```
DSolve[y'[x]==(2*x+3*y[x])/(x-4*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\log \left(\frac{2y(x)^2}{x^2} + \frac{y(x)}{x} + 1 \right) - \frac{4 \arctan \left(\frac{\frac{4y(x)}{x} + 1}{\sqrt{7}} \right)}{\sqrt{7}} = -2 \log(x) + c_1, y(x) \right]$$

4.9 problem 9

Internal problem ID [966]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [$y'_x = G(x, y)$]

$$y' - \sqrt{x^2 + y^2} = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)^(1/2),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x^2+y[x]^2)^(1/2),y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.10 problem 10

Internal problem ID [967]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - x(-1 + y^2)^{\frac{2}{3}} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(diff(y(x),x)=x*(y(x)^2-1)^(2/3),y(x), singsol=all)
```

$$\frac{x^2}{2} - \frac{(-\operatorname{signum}(y(x)^2 - 1))^{\frac{2}{3}} y(x) \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{2}{3}\right], \left[\frac{3}{2}\right], y(x)^2\right)}{\operatorname{signum}(y(x)^2 - 1)^{\frac{2}{3}}} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 6.768 (sec). Leaf size: 66

```
DSolve[y'[x]==x*(y[x]^2-1)^(2/3),y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \operatorname{InverseFunction}\left[\frac{\#1(1 - \#1^2)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{2}, \#1^2\right)}{(\#1^2 - 1)^{2/3}} \&\right]\left[\frac{x^2}{2} + c_1\right]$$

$y(x) \rightarrow -1$

$y(x) \rightarrow 1$

4.11 problem 11

Internal problem ID [968]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [$y' = G(x, y)$]

$$y' - (x^2 + y^2)^2 = 0$$

X Solution by Maple

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)^2,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x^2+y[x]^2)^2,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.12 problem 12

Internal problem ID [969]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _dAlembert]`

$$y' - \sqrt{x+y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 42

```
dsolve(diff(y(x),x)=(x+y(x))^(1/2),y(x), singsol=all)
```

$$\begin{aligned} x - 2\sqrt{x+y(x)} - \ln\left(-1 + \sqrt{x+y(x)}\right) \\ + \ln\left(1 + \sqrt{x+y(x)}\right) + \ln(x+y(x)-1) - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 9.242 (sec). Leaf size: 59

```
DSolve[y'[x]==(x+y[x])^(1/2),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) &\rightarrow W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right)^2 + 2W\left(-e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right) - x + 1 \\ y(x) &\rightarrow 1 - x \end{aligned}$$

4.13 problem 13

Internal problem ID [970]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - \frac{\tan(y)}{x-1} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=tan(y(x))/(x-1),y(x), singsol=all)
```

$$y(x) = \arcsin(c_1(x-1))$$

✓ Solution by Mathematica

Time used: 7.817 (sec). Leaf size: 19

```
DSolve[y'[x]==Tan[y[x]]/(x-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(e^{c_1}(x-1))$$

$$y(x) \rightarrow 0$$

4.14 problem 16

Internal problem ID [971]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y^{\frac{2}{5}} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)=y(x)^(2/5),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(3x + 5) \left(\frac{3x}{5} + 1\right)^{\frac{2}{3}}}{5}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 23

```
DSolve[{y'[x]==y[x]^(2/5),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(3x + 5)^{5/3}}{5 \cdot 5^{2/3}}$$

4.15 problem 18

Internal problem ID [972]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - 3x(y - 1)^{\frac{1}{3}} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=3*x*(y(x)-1)^(1/3),y(0) = 1],y(x), singsol=all)
```

$$y(x) = 1$$

✓ Solution by Mathematica

Time used: 0.296 (sec). Leaf size: 19

```
DSolve[{y'[x]==3*x*(y[x]-1)^(1/3),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$
$$y(x) \rightarrow (x^2)^{3/2} + 1$$

4.16 problem 19

Internal problem ID [973]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$y' - 3x(y - 1)^{\frac{1}{3}} = 0$$

With initial conditions

$$[y(0) = 9]$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)=3*x*(y(x)-1)^(1/3),y(0) = 9],y(x), singsol=all)
```

$$y(x) = x^2\sqrt{x^2 + 4} + 4\sqrt{x^2 + 4} + 1$$

✓ Solution by Mathematica

Time used: 0.125 (sec). Leaf size: 16

```
DSolve[{y'[x]==3*x*(y[x]-1)^(1/3),y[0]==9},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x^2 + 4)^{3/2} + 1$$

4.17 problem 20(a)

Internal problem ID [974]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Existence and Uniqueness of Solutions of Nonlinear Equations. Section 2.3 Page 60

Problem number: 20(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y' - 3x(y - 1)^{\frac{1}{3}} = 0$$

With initial conditions

$$[y(3) = -7]$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 19

```
dsolve([diff(y(x),x)=3*x*(y(x)-1)^(1/3),y(3) = -7],y(x), singsol=all)
```

$$y(x) = 1 + \left(-11 + 2i\sqrt{3} + x^2\right)^{\frac{3}{2}}$$

✓ Solution by Mathematica

Time used: 0.134 (sec). Leaf size: 49

```
DSolve[{y'[x]==3*x*(y[x]-1)^(1/3),y[3]==-7},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 + \left(x^2 - 2i\sqrt{3} - 11\right)^{3/2}$$

$$y(x) \rightarrow 1 + \left(x^2 + 2i\sqrt{3} - 11\right)^{3/2}$$

5 Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

5.1	problem Example 1	118
5.2	problem Example 2	119
5.3	problem Example 3(a) (As Riccati)	120
5.4	problem Example 3(b)	121
5.5	problem 1	122
5.6	problem 2	123
5.7	problem 3	125
5.8	problem 4	126
5.9	problem 5	127
5.10	problem 6	128
5.11	problem 7	130
5.12	problem 8	131
5.13	problem 9	132
5.14	problem 10	133
5.15	problem 11	134
5.16	problem 12	135
5.17	problem 13	136
5.18	problem 15	137
5.19	problem 16	138
5.20	problem 17	139
5.21	problem 18	140
5.22	problem 19	141
5.23	problem 20	142
5.24	problem 21	143
5.25	problem 22	144
5.26	problem 23	145
5.27	problem 24	146
5.28	problem 25	147
5.29	problem 26	148
5.30	problem 27	149
5.31	problem 28	150
5.32	problem 29	151
5.33	problem 30	152
5.34	problem 31	153

5.35	problem 32	155
5.36	problem 33	158
5.37	problem 34	160
5.38	problem 35(a)	162
5.39	problem 36(a)	163
5.40	problem 37(a)	164
5.41	problem 38	166
5.42	problem 41	167
5.43	problem 42	168
5.44	problem 43	169
5.45	problem 44	170
5.46	problem 45	171
5.47	problem 46	172
5.48	problem 47	173
5.49	problem 48	174
5.50	problem 49	175
5.51	problem 50	176
5.52	problem 51	177
5.53	problem 52	178
5.54	problem 53	179
5.55	problem 56	180

5.1 problem Example 1

Internal problem ID [975]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: Example 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' - y - xy^2 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)-y(x)=x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{1 + e^{-x}c_1 - x}$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 27

```
DSolve[y'[x]-y[x]==x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{-e^x(x-1) + c_1}$$
$$y(x) \rightarrow 0$$

5.2 problem Example 2

Internal problem ID [976]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: Example 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _dAlembert]`

$$y' - \frac{y + x e^{-\frac{y}{x}}}{x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=(y(x)+x*exp(-y(x)/x))/x,y(x), singsol=all)
```

$$y(x) = \ln(\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.368 (sec). Leaf size: 13

```
DSolve[y'[x]==(y[x]+x*Exp[-y[x]/x])/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \log(\log(x) + c_1)$$

5.3 problem Example 3(a) (As Riccati)

Internal problem ID [977]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: Example 3(a) (As Riccati).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$y'x^2 - y^2 - yx = -x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(x^2*diff(y(x),x)=y(x)^2+x*y(x)-x^2,y(x), singsol=all)
```

$$y(x) = -\tanh(\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.298 (sec). Leaf size: 298

```
DSolve[y'[x]==y[x]^2+x*y[x]-x^2,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{5(\sqrt{5}-1)x \left(c_1 \operatorname{HermiteH} \left(\frac{1}{10}(-5+\sqrt{5}), \frac{\sqrt[4]{5}x}{\sqrt{2}} \right) - \operatorname{Hypergeometric1F1} \left(\frac{5}{4} - \frac{1}{4\sqrt{5}}, \frac{3}{2}, \frac{\sqrt{5}x^2}{2} \right) + \operatorname{Hypergeometric1F1} \left(\frac{5}{4} + \frac{1}{4\sqrt{5}}, \frac{3}{2}, \frac{\sqrt{5}x^2}{2} \right) \right)}{10 \left(\operatorname{Hypergeometric1F1} \left(\frac{1}{20}(5-\sqrt{5}), \frac{1}{2}, \frac{\sqrt{5}x^2}{2} \right) + \operatorname{Hypergeometric1F1} \left(\frac{1}{20}(5+\sqrt{5}), \frac{1}{2}, \frac{\sqrt{5}x^2}{2} \right) \right)}$$

$$y(x) \rightarrow \frac{1}{2}(\sqrt{5}-1)x - \frac{(\sqrt{5}-5) \operatorname{HermiteH} \left(\frac{1}{10}(-15+\sqrt{5}), \frac{\sqrt[4]{5}x}{\sqrt{2}} \right)}{\sqrt{2}5^{3/4} \operatorname{HermiteH} \left(\frac{1}{10}(-5+\sqrt{5}), \frac{\sqrt[4]{5}x}{\sqrt{2}} \right)}$$

5.4 problem Example 3(b)

Internal problem ID [978]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: Example 3(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y'x^2 - y^2 - yx = -x^2$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x)=y(x)^2+x*y(x)-x^2,y(1) = 2],y(x), singsol=all)
```

$$y(x) = -\frac{x(x^2 + 3)}{x^2 - 3}$$

✓ Solution by Mathematica

Time used: 0.576 (sec). Leaf size: 20

```
DSolve[{x^2*y'[x]==y[x]^2+x*y[x]-x^2,y[1]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x(x^2 + 3)}{x^2 - 3}$$

5.5 problem 1

Internal problem ID [979]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' + y - y^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+y(x)=y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{1 + e^{x+c_1}}$$

✓ Solution by Mathematica

Time used: 0.777 (sec). Leaf size: 54

```
DSolve[y'[x]+y[x]==y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{1 + e^{2(x+c_1)}}}$$

$$y(x) \rightarrow \frac{1}{\sqrt{1 + e^{2(x+c_1)}}}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 1$$

5.6 problem 2

Internal problem ID [980]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational, _Bernoulli]`

$$7y'x - 2y + \frac{x^2}{y^6} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 134

```
dsolve(7*x*diff(y(x),x)-2*y(x)=-x^2/y(x)^6,y(x), singsol=all)
```

$$\begin{aligned}y(x) &= (-x^2(\ln(x) - c_1))^{\frac{1}{7}} \\y(x) &= -((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{1}{7}} \\y(x) &= ((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{6}{7}} \\y(x) &= -((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{5}{7}} \\y(x) &= ((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{2}{7}} \\y(x) &= -((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{3}{7}} \\y(x) &= ((c_1 - \ln(x)) x^2)^{\frac{1}{7}} (-1)^{\frac{4}{7}}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.238 (sec). Leaf size: 181

```
DSolve[7*x*y'[x]-2*y[x]==-x^2/y[x]^6,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow -\sqrt[7]{-1} x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow (-1)^{2/7} x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow -(-1)^{3/7} x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow (-1)^{4/7} x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow -(-1)^{5/7} x^{2/7} \sqrt[7]{-\log(x) + c_1} \\y(x) &\rightarrow (-1)^{6/7} x^{2/7} \sqrt[7]{-\log(x) + c_1}\end{aligned}$$

5.7 problem 3

Internal problem ID [981]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y'x^2 + 2y - 2e^{\frac{1}{x}}\sqrt{y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x)+2*y(x)=2*exp(1/x)*y(x)^(1/2),y(x), singsol=all)
```

$$\frac{-c_1 x e^{\frac{1}{x}} + x \sqrt{y(x)} + e^{\frac{1}{x}}}{x} = 0$$

✓ Solution by Mathematica

Time used: 0.269 (sec). Leaf size: 39

```
DSolve[y'[x]+2*y[x]==2*Exp[1/x]*y[x]^(1/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} e^{-2x} \left(\int_1^x 2e^{K[1] + \frac{1}{K[1]}} dK[1] + 2c_1 \right)^2$$

5.8 problem 4

Internal problem ID [982]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, _Bernoulli]`

$$(x^2 + 1) y' + 2yx - \frac{1}{(x^2 + 1)y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve((1+x^2)*diff(y(x),x)+2*x*y(x)=1/((1+x^2)*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{2x + c_1}}{x^2 + 1}$$
$$y(x) = -\frac{\sqrt{2x + c_1}}{x^2 + 1}$$

✓ Solution by Mathematica

Time used: 0.264 (sec). Leaf size: 46

```
DSolve[(1+x^2)*y'[x]+2*x*y[x]==1/((1+x^2)*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{2x + c_1}}{x^2 + 1}$$
$$y(x) \rightarrow \frac{\sqrt{2x + c_1}}{x^2 + 1}$$

5.9 problem 5

Internal problem ID [983]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$-yx + y' - x^3y^3 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)-x*y(x)=x^3*y(x)^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{e^{-x^2}c_1 - x^2 + 1}}$$
$$y(x) = -\frac{1}{\sqrt{e^{-x^2}c_1 - x^2 + 1}}$$

✓ Solution by Mathematica

Time used: 1.866 (sec). Leaf size: 80

```
DSolve[y'[x]-x*y[x]==x^3*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{ie^{\frac{x^2}{2}}}{\sqrt{e^{x^2}(x^2 - 1) - c_1}}$$
$$y(x) \rightarrow \frac{ie^{\frac{x^2}{2}}}{\sqrt{e^{x^2}(x^2 - 1) - c_1}}$$
$$y(x) \rightarrow 0$$

5.10 problem 6

Internal problem ID [984]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_rational, _Bernoulli]

$$y' - \frac{(x+1)y}{3x} - y^4 = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 121

```
dsolve(diff(y(x),x)-(1+x)/(3*x)*y(x)=y(x)^4,y(x), singsol=all)
```

$$y(x) = \frac{\left(x(e^{-x}c_1 - 3x + 3)^2\right)^{\frac{1}{3}}}{e^{-x}c_1 - 3x + 3}$$
$$y(x) = \frac{(1 + i\sqrt{3}) \left(x(e^{-x}c_1 - 3x + 3)^2\right)^{\frac{1}{3}}}{-2e^{-x}c_1 + 6x - 6}$$
$$y(x) = \frac{(i\sqrt{3} - 1) \left(x(e^{-x}c_1 - 3x + 3)^2\right)^{\frac{1}{3}}}{2e^{-x}c_1 - 6x + 6}$$

✓ Solution by Mathematica

Time used: 1.88 (sec). Leaf size: 120

```
DSolve[y'[x]-(1+x)/(3*x)*y[x]==y[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{x/3} \sqrt[3]{x}}{\sqrt[3]{3e^x(x-1) - c_1}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-1} e^{x/3} \sqrt[3]{x}}{\sqrt[3]{3e^x(x-1) - c_1}}$$

$$y(x) \rightarrow -\frac{(-1)^{2/3} e^{x/3} \sqrt[3]{x}}{\sqrt[3]{3e^x(x-1) - c_1}}$$

$$y(x) \rightarrow 0$$

5.11 problem 7

Internal problem ID [985]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' - 2y - xy^3 = 0$$

With initial conditions

$$[y(0) = 2\sqrt{2}]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 13

```
dsolve([diff(y(x),x)-2*y(x)=x*y(x)^3,y(0) = 2*2^(1/2)],y(x), singsol=all)
```

$$y(x) = \frac{4}{\sqrt{-8x + 2}}$$

✓ Solution by Mathematica

Time used: 1.869 (sec). Leaf size: 34

```
DSolve[{y'[x]-2*y[x]==x*y[x]^3,y[0]==2*Sqrt[2]},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2i\sqrt{2}e^{2x}}{\sqrt{e^{4x}(4x - 1)}}$$

5.12 problem 8

Internal problem ID [986]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$-yx + y' - xy^{\frac{3}{2}} = 0$$

With initial conditions

$$[y(1) = 4]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 21

```
dsolve([diff(y(x),x)-x*y(x)=x*y(x)^(3/2),y(1) = 4],y(x), singsol=all)
```

$$y(x) = \frac{4}{\left(-2 + 3e^{-\frac{(x-1)(x+1)}{4}}\right)^2}$$

✓ Solution by Mathematica

Time used: 0.405 (sec). Leaf size: 55

```
DSolve[{y'[x]-x*y[x]==x*y[x]^(3/2),y[1]==4},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4} \left(\tanh \left(\frac{1}{8} (-8 \operatorname{arctanh}(3) - x^2 + 1) \right) - 1 \right)^2$$
$$y(x) \rightarrow \frac{1}{4} \left(\tanh \left(\operatorname{arctanh}(5) - \frac{x^2}{8} + \frac{1}{8} \right) - 1 \right)^2$$

5.13 problem 9

Internal problem ID [987]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$y'x + y - x^4y^4 = 0$$

With initial conditions

$$\left[y(1) = \frac{1}{2} \right]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 35

```
dsolve([x*diff(y(x),x)+y(x)=x^4*y(x)^4,y(1) = 1/2],y(x), singsol=all)
```

$$y(x) = \frac{(-(3x - 11)^2)^{\frac{1}{3}} (i\sqrt{3} - 1)}{6x^2 - 22x}$$

✓ Solution by Mathematica

Time used: 0.415 (sec). Leaf size: 19

```
DSolve[{x*y'[x]+y[x]==x^4*y[x]^4,y[1]==1/2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\sqrt[3]{-x^3(3x - 11)}}$$

5.14 problem 10

Internal problem ID [988]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - 2y - 2\sqrt{y} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)-2*y(x)=2*y(x)^(1/2),y(0) = 1],y(x), singsol=all)
```

$$y(x) = 4e^{2x} - 4e^x + 1$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 14

```
DSolve[{y'[x]-2*y[x]==2*y[x]^(1/2),y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (1 - 2e^x)^2$$

5.15 problem 11

Internal problem ID [989]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_rational, _Bernoulli]

$$y' - 4y - \frac{48x}{y^2} = 0$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 17

```
dsolve([diff(y(x),x)-4*y(x)=48*x/y(x)^2,y(0) = 1],y(x), singsol=all)
```

$$y(x) = (2e^{12x} - 12x - 1)^{\frac{1}{3}}$$

✓ Solution by Mathematica

Time used: 4.195 (sec). Leaf size: 21

```
DSolve[{y'[x]-4*y[x]==48*x/y[x]^2,y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{-12x + 2e^{12x} - 1}$$

5.16 problem 12

Internal problem ID [990]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$y'x^2 + 2yx - y^3 = 0$$

With initial conditions

$$\left[y(1) = \frac{\sqrt{2}}{2} \right]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 26

```
dsolve([x^2*diff(y(x),x)+2*x*y(x)=y(x)^3,y(1) = 1/2*2^(1/2)],y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{10} \sqrt{4x^6 + x}}{8x^5 + 2}$$

✓ Solution by Mathematica

Time used: 0.462 (sec). Leaf size: 29

```
DSolve[{x^2*y'[x]+2*x*y[x]==y[x]^3,y[1]==1/Sqrt[2]},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{\frac{5}{2}} \sqrt{x}}{\sqrt{4x^5 + 1}}$$

5.17 problem 13

Internal problem ID [991]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Bernoulli]

$$y' - y - \sqrt{y}x = 0$$

With initial conditions

$$[y(0) = 4]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 26

```
dsolve([diff(y(x),x)-y(x)=x*y(x)^(1/2),y(0) = 4],y(x), singsol=all)
```

$$y(x) = (-8x - 16)e^{\frac{x}{2}} + x^2 + 4x + 16e^x + 4$$

✓ Solution by Mathematica

Time used: 0.146 (sec). Leaf size: 28

```
DSolve[{y'[x]-y[x]==x*y[x]^(1/2),y[0]==4},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x - 4e^{x/2} + 2)^2$$
$$y(x) \rightarrow (x + 2)^2$$

5.18 problem 15

Internal problem ID [992]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$y' - \frac{x+y}{x} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=(y(x)+x)/x,y(x), singsol=all)
```

$$y(x) = (\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 12

```
DSolve[y'[x]==(y[x]+x)/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\log(x) + c_1)$$

5.19 problem 16

Internal problem ID [993]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{y^2 + 2yx}{x^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=(y(x)^2+2*x*y(x))/x^2,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{c_1 - x}$$

✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 23

```
DSolve[y'[x]==(y[x]^2+2*x*y[x])/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{x - c_1}$$
$$y(x) \rightarrow 0$$

5.20 problem 17

Internal problem ID [994]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$xy^3y' - y^4 = x^4$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 58

```
dsolve(x*y(x)^3*diff(y(x),x)=y(x)^4+x^4,y(x), singsol=all)
```

$$y(x) = (4 \ln(x) + c_1)^{\frac{1}{4}} x$$

$$y(x) = -(4 \ln(x) + c_1)^{\frac{1}{4}} x$$

$$y(x) = -i(4 \ln(x) + c_1)^{\frac{1}{4}} x$$

$$y(x) = i(4 \ln(x) + c_1)^{\frac{1}{4}} x$$

✓ Solution by Mathematica

Time used: 0.2 (sec). Leaf size: 76

```
DSolve[x*y[x]^3*y'[x]==y[x]^4+x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \sqrt[4]{4 \log(x) + c_1}$$

$$y(x) \rightarrow -ix \sqrt[4]{4 \log(x) + c_1}$$

$$y(x) \rightarrow ix \sqrt[4]{4 \log(x) + c_1}$$

$$y(x) \rightarrow x \sqrt[4]{4 \log(x) + c_1}$$

5.21 problem 18

Internal problem ID [995]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _dAlembert]`

$$y' - \frac{y}{x} - \sec\left(\frac{y}{x}\right) = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=y(x)/x+sec(y(x)/x),y(x), singsol=all)
```

$$y(x) = \arcsin(\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.409 (sec). Leaf size: 13

```
DSolve[y'[x]==y[x]/x+Sec[y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(\log(x) + c_1)$$

5.22 problem 19

Internal problem ID [996]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y'x^2 - y^2 - yx = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*diff(y(x),x)=x*y(x)+x^2+y(x)^2,y(x), singsol=all)
```

$$y(x) = \tan(\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.174 (sec). Leaf size: 13

```
DSolve[x^2*y'[x]==x*y[x]+x^2+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(\log(x) + c_1)$$

5.23 problem 20

Internal problem ID [997]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y'xy - 2y^2 = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x*y(x)*diff(y(x),x)=x^2+2*y(x)^2,y(x), singsol=all)
```

$$y(x) = \sqrt{c_1x^2 - 1} x$$
$$y(x) = -\sqrt{c_1x^2 - 1} x$$

✓ Solution by Mathematica

Time used: 0.441 (sec). Leaf size: 38

```
DSolve[x*y[x]*y'[x]==x^2+2*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x\sqrt{-1 + c_1x^2}$$
$$y(x) \rightarrow x\sqrt{-1 + c_1x^2}$$

5.24 problem 21

Internal problem ID [998]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A']]`

$$y' - \frac{2y^2 + x^2 e^{-\frac{y^2}{x^2}}}{2yx} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x)=(2*y(x)^2+x^2*exp(-(y(x)/x)^2))/(2*x*y(x)),y(x), singsol=all)
```

$$y(x) = \sqrt{\ln(\ln(x) + c_1)} x$$
$$y(x) = -\sqrt{\ln(\ln(x) + c_1)} x$$

✓ Solution by Mathematica

Time used: 2.166 (sec). Leaf size: 38

```
DSolve[y'[x]==(2*y[x]^2+x^2*Exp[-(y[x]/x)^2 ])/(2*x*y[x]),y[x],x,IncludeSingularSolutions -
```

$$y(x) \rightarrow -x\sqrt{\log(\log(x) + 2c_1)}$$
$$y(x) \rightarrow x\sqrt{\log(\log(x) + 2c_1)}$$

5.25 problem 22

Internal problem ID [999]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{yx + y^2}{x^2} = 0$$

With initial conditions

$$[y(-1) = 2]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 19

```
dsolve([diff(y(x),x)=(x*y(x)+y(x)^2)/x^2,y(-1) = 2],y(x), singsol=all)
```

$$y(x) = \frac{2x}{-2 \ln(x) - 1 + 2i\pi}$$

✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 25

```
DSolve[{y'[x]==(x*y[x]+y[x]^2)/x^2,y[-1]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2ix}{2i \log(x) + 2\pi + i}$$

5.26 problem 23

Internal problem ID [1000]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y' - \frac{x^3 + y^3}{xy^2} = 0$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)=(x^3+y(x)^3)/(x*y(x)^2),y(1) = 3],y(x), singsol=all)
```

$$y(x) = (3 \ln(x) + 27)^{\frac{1}{3}} x$$

✓ Solution by Mathematica

Time used: 0.21 (sec). Leaf size: 20

```
DSolve[{y'[x]==(x^3+y[x]^3)/(x*y[x]^2),y[1]==3},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{3x^3 \sqrt{\log(x) + 9}}$$

5.27 problem 24

Internal problem ID [1001]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y'xy + y^2 = -x^2$$

With initial conditions

$$[y(1) = 2]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 18

```
dsolve([x*y(x)*diff(y(x),x)+x^2+y(x)^2=0,y(1) = 2],y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-2x^4 + 18}}{2x}$$

✓ Solution by Mathematica

Time used: 0.216 (sec). Leaf size: 25

```
DSolve[{x*y[x]*y'[x]+x^2+y[x]^2==0,y[1]==2},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{9 - x^4}}{\sqrt{2}x}$$

5.28 problem 25

Internal problem ID [1002]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y' - \frac{y^2 - 3yx - 5x^2}{x^2} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.313 (sec). Leaf size: 23

```
dsolve([diff(y(x),x)=(y(x)^2-3*x*y(x)-5*x^2)/x^2,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{-2x^7 + 5x}{2x^6 + 1}$$

✓ Solution by Mathematica

Time used: 1.947 (sec). Leaf size: 24

```
DSolve[{y'[x]==(y[x]^2-3*x*y[x]-5*x^2)/x^2,y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5x - 2x^7}{2x^6 + 1}$$

5.29 problem 26

Internal problem ID [1003]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Riccati]`

$$y'x^2 - y^2 - 4yx = 2x^2$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve([x^2*diff(y(x),x)=2*x^2+y(x)^2+4*x*y(x),y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{-4x^2 + 3x}{2x - 3}$$

✓ Solution by Mathematica

Time used: 0.492 (sec). Leaf size: 20

```
DSolve[{x^2*y'[x]==2*x^2+y[x]^2+4*x*y[x],y[1]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x(4x - 3)}{2x - 3}$$

5.30 problem 27

Internal problem ID [1004]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _Bernoulli]`

$$y'xy - 4y^2 = 3x^2$$

With initial conditions

$$[y(1) = \sqrt{3}]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 15

```
dsolve([x*y(x)*diff(y(x),x)=3*x^2+4*y(x)^2,y(1) = 3^(1/2)],y(x), singsol=all)
```

$$y(x) = \sqrt{4x^6 - 1} x$$

✓ Solution by Mathematica

Time used: 0.603 (sec). Leaf size: 18

```
DSolve[{x*y[x]*y'[x]==3*x^2+4*y[x]^2,y[1]==Sqrt[3]},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x\sqrt{4x^6 - 1}$$

5.31 problem 28

Internal problem ID [1005]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 28.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x+y}{x-y} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=(x+y(x))/(x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-2_Z + \ln \left(\sec \left(_Z \right)^2 \right) + 2 \ln(x) + 2c_1 \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 36

```
DSolve[y'[x]==(x+y[x])/(x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + 1 \right) - \arctan \left(\frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

5.32 problem 29

Internal problem ID [1006]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 29.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A']`

$$(-y + y'x)(\ln(y) - \ln(x)) = x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 24

```
dsolve((diff(y(x),x)*x-y(x))*(ln(y(x))-ln(x))=x,y(x), singsol=all)
```

$$y(x) = \frac{x \ln\left(\frac{x}{c_1}\right)}{\text{LambertW}\left(\ln\left(\frac{x}{c_1}\right) e^{-1}\right)}$$

✓ Solution by Mathematica

Time used: 60.147 (sec). Leaf size: 24

```
DSolve[(y'[x]*x-y[x])*(Log[y[x]]-Log[x])==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x(\log(x) + c_1)}{W\left(\frac{\log(x)+c_1}{e}\right)}$$

5.33 problem 30

Internal problem ID [1007]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 30.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$y' - \frac{y^3 + 2xy^2 + x^2y + x^3}{x(x+y)^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 80

```
dsolve(diff(y(x),x)=(y(x)^3+2*x*y(x)^2+x^2*y(x)+x^3)/(x*(y(x)+x)^2),y(x), singsol=all)
```

$$y(x) = x \left(-1 + (3 \ln(x) + 3c_1)^{\frac{1}{3}} \right)$$
$$y(x) = -\frac{x \left(i\sqrt{3} (3 \ln(x) + 3c_1)^{\frac{1}{3}} + (3 \ln(x) + 3c_1)^{\frac{1}{3}} + 2 \right)}{2}$$
$$y(x) = \frac{x(i\sqrt{3} - 1)(3 \ln(x) + 3c_1)^{\frac{1}{3}}}{2} - x$$

✓ Solution by Mathematica

Time used: 1.409 (sec). Leaf size: 109

```
DSolve[y'[x]==(y[x]^3+2*x*y[x]^2+x^2*y[x]+x^3)/(x*(y[x]+x)^2),y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow -x + \sqrt[3]{x^3(3 \log(x) + 1 + 3c_1)}$$
$$y(x) \rightarrow -x + \frac{1}{2}i(\sqrt{3} + i) \sqrt[3]{x^3(3 \log(x) + 1 + 3c_1)}$$
$$y(x) \rightarrow -x - \frac{1}{2}(1 + i\sqrt{3}) \sqrt[3]{x^3(3 \log(x) + 1 + 3c_1)}$$

5.34 problem 31

Internal problem ID [1008]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 31.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x + 2y}{2x + y} = 0$$

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 271

```
dsolve(diff(y(x),x)=(x+2*y(x))/(2*x+y(x)),y(x), singsol=all)
```

$$y(x) = \frac{x \left(\frac{c_1 \left(\left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{2}{3}} + 3 \right)}{3x \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{1}{3}}} + c_1^2 \right)}{c_1^2}$$

$$y(x) = \frac{(1 + i\sqrt{3}) \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{2}{3}} - 6xc_1 \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{1}{3}} - 3i\sqrt{3} + 3}{6 \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{1}{3}} c_1}$$

$$y(x) = \frac{(i\sqrt{3} - 1) \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{2}{3}} + 6xc_1 \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{1}{3}} - 3i\sqrt{3} - 3}{6 \left(27c_1x + 3\sqrt{3} \sqrt{27c_1^2x^2 - 1} \right)^{\frac{1}{3}} c_1}$$

✓ Solution by Mathematica

Time used: 30.082 (sec). Leaf size: 382

`DSolve[y'[x]==(x+2*y[x])/(2*x+y[x]),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}}{3^{2/3}} - \frac{e^{2c_1}}{\sqrt[3]{3}\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}} + x$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i)\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}}{2 \cdot 3^{2/3}} + \frac{(1 + i\sqrt{3})e^{2c_1}}{2\sqrt[3]{3}\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}} + x$$

$$y(x) \rightarrow -\frac{(1 + i\sqrt{3})\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}}{2 \cdot 3^{2/3}} + \frac{(1 - i\sqrt{3})e^{2c_1}}{2\sqrt[3]{3}\sqrt[3]{\sqrt{3}\sqrt{27e^{4c_1}x^2 + e^{6c_1}} - 9e^{2c_1}x}} + x$$

5.35 problem 32

Internal problem ID [1009]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 32.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A'], _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{y}{-2x + y} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 306

```
dsolve(diff(y(x),x)=y(x)/(y(x)-2*x),y(x), singsol=all)
```

$$y(x) = \frac{\left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{c_1(-4x^3 + 3c_1)}\right)^{\frac{1}{3}}}{2} + \frac{\left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{c_1(-4x^3 + 3c_1)}\right)^{\frac{1}{3}} + x}{2x^2} + x$$

$$y(x) = \frac{(-1 - i\sqrt{3}) \left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}}{4} + \frac{x \left(ix\sqrt{3} - x + \left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}\right)}{\left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{(i\sqrt{3} - 1) \left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}}{4} - \frac{x \left(ix\sqrt{3} + x - \left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}\right)}{\left(-12c_1 + 8x^3 + 4\sqrt{3} \sqrt{-4c_1x^3 + 3c_1^2}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 37.127 (sec). Leaf size: 479

`DSolve[y'[x]==y[x]/(y[x]-2*x),y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{\sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}}{\sqrt[3]{2}} + \frac{\sqrt[3]{2}x^2}{\sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}} + x$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i) \sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}}{2\sqrt[3]{2}} - \frac{(1 + i\sqrt{3})x^2}{2^{2/3} \sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}} + x$$

$$y(x) \rightarrow -\frac{(1 + i\sqrt{3}) \sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}}{2\sqrt[3]{2}} + \frac{i(\sqrt{3} + i)x^2}{2^{2/3} \sqrt[3]{2x^3 + \sqrt{e^{6c_1} - 4e^{3c_1}x^3} - e^{3c_1}}} + x$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\frac{i(\sqrt[3]{x^3} - x) \left((\sqrt{3} - i) \sqrt[3]{x^3} - 2ix \right)}{2x}$$

$$y(x) \rightarrow \frac{i(\sqrt[3]{x^3} - x) \left((\sqrt{3} + i) \sqrt[3]{x^3} + 2ix \right)}{2x}$$

$$y(x) \rightarrow \sqrt[3]{x^3} + \frac{(x^3)^{2/3}}{x} + x$$

5.36 problem 33

Internal problem ID [1010]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 33.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, _dAlembert]`

$$y' - \frac{xy^2 + 2y^3}{x^3 + x^2y + xy^2} = 0$$

✓ Solution by Maple

Time used: 2.969 (sec). Leaf size: 129

```
dsolve(diff(y(x),x)=(x*y(x)^2+2*y(x)^3)/(x^3+x^2*y(x)+x*y(x)^2),y(x), singsol=all)
```

$$y(x) = x \left(\text{RootOf} \left(_Z^8 c_1 x^2 + 2_Z^6 c_1 x^2 + _Z^4 c_1 x^2 - 2_Z^2 - 1 \right)^6 c_1 x^2 \right. \\ \left. + 2 \text{RootOf} \left(_Z^8 c_1 x^2 + 2_Z^6 c_1 x^2 + _Z^4 c_1 x^2 - 2_Z^2 - 1 \right)^4 c_1 x^2 \right. \\ \left. + \text{RootOf} \left(_Z^8 c_1 x^2 + 2_Z^6 c_1 x^2 + _Z^4 c_1 x^2 - 2_Z^2 - 1 \right)^2 c_1 x^2 - 1 \right)$$

✓ Solution by Mathematica

Time used: 60.157 (sec). Leaf size: 1989

`DSolve[y'[x]==(x*y[x]^2+2*y[x]^3)/(x^3+x^2*y[x]+x*y[x]^2),y[x],x,IncludeSingularSolutions ->`

$$y(x) \rightarrow \frac{1}{6} \left(-\sqrt{3} \sqrt{-2e^{2c_1}x^4 + 3x^2 + \frac{e^{4c_1}x^8}{\sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8 + 6\sqrt{3}\sqrt{e^{4c_1}x^{16}(27 + e^{4c_1}x^4)}}}} + \sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8} - \sqrt{3} \sqrt{-4e^{2c_1}x^4 + 6x^2 - \frac{e^{4c_1}x^8}{\sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8 + 6\sqrt{3}\sqrt{e^{4c_1}x^{16}(27 + e^{4c_1}x^4)}}}} - \sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8} + 3x \right)$$

$$y(x) \rightarrow \frac{1}{6} \left(-\sqrt{3} \sqrt{-2e^{2c_1}x^4 + 3x^2 + \frac{e^{4c_1}x^8}{\sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8 + 6\sqrt{3}\sqrt{e^{4c_1}x^{16}(27 + e^{4c_1}x^4)}}}} + \sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8} + \sqrt{3} \sqrt{-4e^{2c_1}x^4 + 6x^2 - \frac{e^{4c_1}x^8}{\sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8 + 6\sqrt{3}\sqrt{e^{4c_1}x^{16}(27 + e^{4c_1}x^4)}}}} - \sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8} + 3x \right)$$

$$y(x) \rightarrow \frac{1}{6} \left(-\sqrt{3} \sqrt{-2e^{2c_1}x^4 + 3x^2 + \frac{e^{4c_1}x^8}{\sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8 + 6\sqrt{3}\sqrt{e^{4c_1}x^{16}(27 + e^{4c_1}x^4)}}}} + \sqrt[3]{e^{6c_1}x^{12} + 54e^{2c_1}x^8} \right)$$

5.37 problem 34

Internal problem ID [1011]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 34.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _dAlembert]`

$$y' - \frac{x^3 + x^2y + 3y^3}{x^3 + 3xy^2} = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 276

```
dsolve(diff(y(x),x)=(x^3+x^2*y(x)+3*y(x)^3)/(x^3+3*x*y(x)^2),y(x), singsol=all)
```

$$y(x) = \frac{\left(\left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{2}{3}} - 12 \right) x}{6 \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{1}{3}}}$$

$$y(x) = \frac{\left(i\sqrt{3} \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{2}{3}} + 12i\sqrt{3} + \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{2}{3}} \right) x}{12 \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{1}{3}}}$$

$$y(x) = \frac{x \left((i\sqrt{3} - 1) \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{2}{3}} + 12i\sqrt{3} + 12 \right)}{12 \left(108 \ln(x) + 108c_1 + 12\sqrt{12 + 81 \ln(x)^2 + 162 \ln(x) c_1 + 81c_1^2} \right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 27.532 (sec). Leaf size: 398

`DSolve[y'[x]==(x^3+x^2*y[x]+3*y[x]^3)/(x^3+3*x*y[x]^2),y[x],x,IncludeSingularSolutions -> Tr`

$$y(x) \rightarrow \frac{-2\sqrt[3]{3}x^2 + \sqrt[3]{2}\left(\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3\right)^{2/3}}{6^{2/3}\sqrt[3]{\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3}}$$

$$y(x) \rightarrow \frac{2\sqrt[3]{3}(1+i\sqrt{3})x^2 + i\sqrt[3]{2}(\sqrt{3}+i)\left(\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3\right)^{2/3}}{2\cdot 6^{2/3}\sqrt[3]{\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3}}$$

$$y(x) \rightarrow \frac{2\sqrt[3]{3}(1-i\sqrt{3})x^2 - \sqrt[3]{2}(1+i\sqrt{3})\left(\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3\right)^{2/3}}{2\cdot 6^{2/3}\sqrt[3]{\sqrt{3}\sqrt{x^6(4+27(\log(x)+c_1)^2)} + 9x^3\log(x) + 9c_1x^3}}$$

5.38 problem 35(a)

Internal problem ID [1012]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 35(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class A'], _rational, _Riccati]`

$$y'x^2 - y^2 - yx = -4x^2$$

With initial conditions

$$[y(-1) = 0]$$

✓ Solution by Maple

Time used: 0.312 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x)=y(x)^2+x*y(x)-4*x^2,y(-1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{-2x^5 + 2x}{x^4 + 1}$$

✓ Solution by Mathematica

Time used: 2.186 (sec). Leaf size: 20

```
DSolve[{x^2*y'[x]==y[x]^2+x*y[x]-4*x^2,y[-1]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x(x^4 - 1)}{x^4 + 1}$$

5.39 problem 36(a)

Internal problem ID [1013]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 36(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y'xy - y^2 + yx = x^2$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve(x*y(x)*diff(y(x),x)=x^2-x*y(x)+y(x)^2,y(x), singsol=all)
```

$$y(x) = x \left(1 + \text{LambertW} \left(\frac{e^{-1-c_1}}{x} \right) \right)$$

✓ Solution by Mathematica

Time used: 3.649 (sec). Leaf size: 25

```
DSolve[x*y[x]*y'[x]==x^2-x*y[x]+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \left(1 + W \left(\frac{e^{-1+c_1}}{x} \right) \right)$$

$$y(x) \rightarrow x$$

5.40 problem 37(a)

Internal problem ID [1014]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 37(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{2y^2 - yx + 2x^2}{yx + 2x^2} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)=(2*y(x)^2-x*y(x)+2*x^2)/(x*y(x)+2*x^2),y(x), singsol=all)
```

$y(x) = \text{RootOf}(_Z^4 + c_1x + 16 + (-3c_1x - 32)_Z + (3c_1x + 24)_Z^2 + (-c_1x - 8)_Z^3) x$

✓ Solution by Mathematica

Time used: 60.166 (sec). Leaf size: 1913

`DSolve[y'[x]==(2*y[x]^2-x*y[x]+2*x^2)/(x*y[x]+2*x^2),y[x],x,IncludeSingularSolutions -> True`

$$y(x) \rightarrow \frac{1}{12} \left(- \sqrt{9e^{2c_1}x^4 + 6 \cdot 2^{2/3} \sqrt[3]{3\sqrt{3}\sqrt{e^{3c_1}x^{15}(-256 + 27e^{c_1}x)} + 27e^{2c_1}x^8 + 24e^{c_1}x^3} \left(-3 + \frac{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}} \right) \right. \right.$$

$$-6 \left. \sqrt{\frac{1}{2}x^2(-8 + e^{c_1}x)^2 + 4x^2(-8 + e^{c_1}x) - \frac{\sqrt[3]{27e^{2c_1}x^8 + \sqrt{729e^{4c_1}x^{16} - 6912e^{3c_1}x^{15}}}}{3\sqrt[3]{2}} - \frac{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}} \right.}$$

$$\left. - 3x(-8 + e^{c_1}x) \right)$$

$$y(x) \rightarrow \frac{1}{12} \left(- \sqrt{9e^{2c_1}x^4 + 6 \cdot 2^{2/3} \sqrt[3]{3\sqrt{3}\sqrt{e^{3c_1}x^{15}(-256 + 27e^{c_1}x)} + 27e^{2c_1}x^8 + 24e^{c_1}x^3} \left(-3 + \frac{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}} \right) \right. \right.$$

$$+6 \left. \sqrt{\frac{1}{2}x^2(-8 + e^{c_1}x)^2 + 4x^2(-8 + e^{c_1}x) - \frac{\sqrt[3]{27e^{2c_1}x^8 + \sqrt{729e^{4c_1}x^{16} - 6912e^{3c_1}x^{15}}}}{3\sqrt[3]{2}} - \frac{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}{\sqrt[3]{\sqrt{3}\sqrt{e^{3c_1}x^{15}}}}} \right.}$$

5.41 problem 38

Internal problem ID [1015]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{x^2 + yx + y^2}{yx} = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)=(x*y(x)+x^2+y(x)^2)/(x*y(x)),y(x), singsol=all)
```

$$y(x) = x \left(-\text{LambertW} \left(-\frac{e^{-1-c_1}}{x} \right) - 1 \right)$$

✓ Solution by Mathematica

Time used: 4.422 (sec). Leaf size: 31

```
DSolve[y'[x]==(x*y[x]+x^2+y[x]^2)/(x*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \left(1 + W \left(-\frac{e^{-1-c_1}}{x} \right) \right)$$
$$y(x) \rightarrow -x$$

5.42 problem 41

Internal problem ID [1016]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 41.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{-6x + y - 3}{2x - y - 1} = 0$$

✓ Solution by Maple

Time used: 1.422 (sec). Leaf size: 99

```
dsolve(diff(y(x),x)=(-6*x+y(x)-3)/(2*x-y(x)-1),y(x), singsol=all)
```

$$y(x) = \frac{-\text{RootOf}(_Z^{25} + (-5c_1x^5 - 25c_1x^4 - 50c_1x^3 - 50c_1x^2 - 25c_1x - 5c_1)_Z^5 - c_1x^5 - 5c_1x^4 - 10c_1x^3 - \dots)}{c_1(x+1)^4}$$

✓ Solution by Mathematica

Time used: 60.095 (sec). Leaf size: 3011

```
DSolve[y'[x]==(-6*x+y[x]-3)/(2*x-y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

5.43 problem 42

Internal problem ID [1017]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 42.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{2x + y + 1}{x + 2y - 4} = 0$$

✓ Solution by Maple

Time used: 4.359 (sec). Leaf size: 138

```
dsolve(diff(y(x),x)=(2*x+y(x)+1)/(x+2*y(x)-4),y(x), singsol=all)
```

$$y(x) = \frac{(x + 5) \text{RootOf}(_Z^{16} + (2c_1x^4 + 16c_1x^3 + 48c_1x^2 + 64c_1x + 32c_1)_Z^4 - c_1x^4 - 8c_1x^3 - 24c_1x^2 - 32c_1x - 16c_1), \text{RootOf}(_Z^{16} + (2c_1x^4 + 16c_1x^3 + 48c_1x^2 + 64c_1x + 32c_1)_Z^4 - c_1x^4 - 8c_1x^3 - 24c_1x^2 - 32c_1x - 16c_1)}{\text{RootOf}(_Z^{16} + (2c_1x^4 + 16c_1x^3 + 48c_1x^2 + 64c_1x + 32c_1)_Z^4 - c_1x^4 - 8c_1x^3 - 24c_1x^2 - 32c_1x - 16c_1)}$$

✓ Solution by Mathematica

Time used: 60.312 (sec). Leaf size: 8077

```
DSolve[y'[x]==(2*x+y[x]+1)/(x+2*y[x]-4),y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

5.44 problem 43

Internal problem ID [1018]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 43.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class C', _rational, [_Abel, '2nd type', 'cl`

$$y' - \frac{-x + 3y - 14}{x + y - 2} = 0$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)=(-x+3*y(x)-14)/(x+y(x)-2),y(x), singsol=all)
```

$$y(x) = \frac{(x + 6) \text{LambertW}(-2c_1(2 + x)) + 2x + 4}{\text{LambertW}(-2c_1(2 + x))}$$

✓ Solution by Mathematica

Time used: 1.05 (sec). Leaf size: 144

```
DSolve[y'[x]==(-x+3*y[x]-14)/(x+y[x]-2),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{2^{2/3} \left(x \log \left(\frac{y(x)-x-6}{y(x)+x-2} \right) - (x+6) \log \left(\frac{x+2}{y(x)+x-2} \right) + 6 \log \left(\frac{y(x)-x-6}{y(x)+x-2} \right) + y(x) \left(\log \left(\frac{x+2}{y(x)+x-2} \right) - \log \left(\frac{y(x)-x-6}{y(x)+x-2} \right) \right) \right)}{9(-y(x) + x + 6)} \right]$$

5.45 problem 44

Internal problem ID [1019]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 44.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$3y'y^2x - y^3 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
dsolve(3*x*y(x)^2*diff(y(x),x)=y(x)^3+x,y(x), singsol=all)
```

$$y(x) = ((\ln(x) + c_1)x)^{\frac{1}{3}}$$
$$y(x) = -\frac{((\ln(x) + c_1)x)^{\frac{1}{3}}(1 + i\sqrt{3})}{2}$$
$$y(x) = \frac{((\ln(x) + c_1)x)^{\frac{1}{3}}(i\sqrt{3} - 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.196 (sec). Leaf size: 69

```
DSolve[3*x*y[x]^2*y'[x]==y[x]^3+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{x} \sqrt[3]{\log(x) + c_1}$$
$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{x} \sqrt[3]{\log(x) + c_1}$$
$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{x} \sqrt[3]{\log(x) + c_1}$$

5.46 problem 45

Internal problem ID [1020]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 45.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$y'xy - 6y^2 = 3x^6$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(x*y(x)*diff(y(x),x)=3*x^6+6*y(x)^2,y(x), singsol=all)
```

$$y(x) = \sqrt{c_1 x^6 - 1} x^3$$

$$y(x) = -\sqrt{c_1 x^6 - 1} x^3$$

✓ Solution by Mathematica

Time used: 0.672 (sec). Leaf size: 42

```
DSolve[x*y[x]*y'[x]==3*x^6+6*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^3 \sqrt{-1 + c_1 x^6}$$

$$y(x) \rightarrow x^3 \sqrt{-1 + c_1 x^6}$$

5.47 problem 46

Internal problem ID [1021]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 46.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Riccati]`

$$x^3 y' - 2y^2 - 2x^2 y = -2x^4$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^3*diff(y(x),x)=2*(y(x)^2+x^2*y(x)-x^4),y(x), singsol=all)
```

$$y(x) = \tanh(-2 \ln(x) + 2c_1) x^2$$

✓ Solution by Mathematica

Time used: 1.137 (sec). Leaf size: 62

```
DSolve[x^3*y'[x]==2*(y[x]^2+x^2*y[x]-x^4),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow ix^2 \tan(2i \log(x) + c_1)$$
$$y(x) \rightarrow \frac{x^2(-x^4 + e^{2i \text{Interval}[\{0,\pi\}]})}{x^4 + e^{2i \text{Interval}[\{0,\pi\}]}}$$

5.48 problem 47

Internal problem ID [1022]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 47.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, _with_linear_symmetries], _Riccati]`

$$y' - y^2 e^{-x} - 4y = 2e^x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)=y(x)^2*exp(-x)+4*y(x)+2*exp(x),y(x), singsol=all)
```

$$y(x) = -\frac{2e^x(e^x c_1 - 1)}{-2 + e^x c_1}$$

✓ Solution by Mathematica

Time used: 0.265 (sec). Leaf size: 30

```
DSolve[y'[x]==y[x]^2*Exp[-x]+4*y[x]+2*Exp[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -2e^x + \frac{1}{e^{-x} + c_1}$$
$$y(x) \rightarrow -2e^x$$

5.49 problem 48

Internal problem ID [1023]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 48.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_Riccati]

$$y' - \frac{y^2 + y \tan(x) + \tan(x)^2}{\sin(x)^2} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 102

```
dsolve(diff(y(x),x)=(y(x)^2+y(x)*tan(x)+tan(x)^2)/sin(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\tan(x) \left(c_1 \sin \left(\frac{\ln(\sin(x)-1)}{2} + \frac{\ln(\sin(x)+1)}{2} - \ln(\sin(x)) \right) - \cos \left(\frac{\ln(\sin(x)-1)}{2} + \frac{\ln(\sin(x)+1)}{2} - \ln(\sin(x)) \right) \right)}{c_1 \cos \left(\frac{\ln(\sin(x)-1)}{2} + \frac{\ln(\sin(x)+1)}{2} - \ln(\sin(x)) \right) + \sin \left(\frac{\ln(\sin(x)-1)}{2} + \frac{\ln(\sin(x)+1)}{2} - \ln(\sin(x)) \right)}$$

✓ Solution by Mathematica

Time used: 0.678 (sec). Leaf size: 20

```
DSolve[y'[x]==(y[x]^2+y[x]*Tan[x]+Tan[x]^2)/Sin[x]^2,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \tan(x) \tan(\log(\sin(x)) - \log(\cos(x))) + c_1$$

5.50 problem 49

Internal problem ID [1024]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 49.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(y)]'], _Riccati]`

$$x \ln(x)^2 y' - \ln(x) y - y^2 = -4 \ln(x)^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x*(ln(x))^2*diff(y(x),x)=-4*(ln(x))^2+y(x)*ln(x)+y(x)^2,y(x), singsol=all)
```

$$y(x) = 2i \tan(2i \ln(\ln(x)) + c_1) \ln(x)$$

✓ Solution by Mathematica

Time used: 1.259 (sec). Leaf size: 64

```
DSolve[x*(Log[x])^2*y'[x]==-4*(Log[x])^2+y[x]*Log[x]+y[x]^2,y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow 2i \log(x) \tan(2i \log(\log(x)) + c_1)$$
$$y(x) \rightarrow \frac{2 \log(x) (-\log^4(x) + e^{2i \text{Interval}\{0,\pi\}})}{\log^4(x) + e^{2i \text{Interval}\{0,\pi\}}}$$

5.51 problem 50

Internal problem ID [1025]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 50.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$2x(y + 2\sqrt{x})y' - (y + \sqrt{x})^2 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(2*x*(y(x)+2*sqrt(x))*diff(y(x),x)=(y(x)+sqrt(x))^2,y(x), singsol=all)
```

$$y(x) = \frac{-2x + \sqrt{x^2 (\ln(x) - c_1 + 4)}}{\sqrt{x}}$$
$$y(x) = -\frac{2x + \sqrt{x^2 (\ln(x) - c_1 + 4)}}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.61 (sec). Leaf size: 68

```
DSolve[2*x*(y[x]+2*Sqrt[x])*y'[x]==(y[x]+Sqrt[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2\sqrt{x} - \sqrt{\frac{1}{x^2}x\sqrt{x(\log(x) + 4 + c_1)}}$$
$$y(x) \rightarrow -2\sqrt{x} + \sqrt{\frac{1}{x^2}x\sqrt{x(\log(x) + 4 + c_1)}}$$

5.52 problem 51

Internal problem ID [1026]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 51.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x),G(y)]'], [_Abel, '2nd type`

$$(y + e^{x^2}) y' - 2x(y^2 + e^{x^2} y + e^{2x^2}) = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
dsolve((y(x)+exp(x^2))*diff(y(x),x)=2*x*(y(x)^2+y(x)*exp(x^2)+exp(2*x^2)),y(x), singsol=all)
```

$$y(x) = \left(-1 - \sqrt{2x^2 - 2c_1 + 1}\right) e^{x^2}$$

$$y(x) = \left(-1 + \sqrt{2x^2 - 2c_1 + 1}\right) e^{x^2}$$

✓ Solution by Mathematica

Time used: 0.744 (sec). Leaf size: 76

```
DSolve[(y[x]+Exp[x^2])*y'[x]==2*x*(y[x]^2+y[x]*Exp[x^2]+Exp[2*x^2]),y[x],x,IncludeSingularSo
```

$$y(x) \rightarrow -e^{x^2} - \frac{\sqrt{2x^2 + 1 + c_1}}{\sqrt{e^{-2x^2}}}$$

$$y(x) \rightarrow -e^{x^2} + \frac{\sqrt{2x^2 + 1 + c_1}}{\sqrt{e^{-2x^2}}}$$

5.53 problem 52

Internal problem ID [1027]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 52.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$y' + \frac{2y}{x} - \frac{3x^2y^2 + 6yx + 2}{x^2(2yx + 3)} = 0$$

With initial conditions

$$[y(2) = 2]$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)+2/x*y(x)=(3*x^2*y(x)^2+6*x*y(x)+2)/(x^2*(2*x*y(x)+3)),y(2) = 2],y(x), s
```

$$y(x) = \frac{-3 + \sqrt{60x + 1}}{2x}$$

✓ Solution by Mathematica

Time used: 0.725 (sec). Leaf size: 35

```
DSolve[{y'[x]+2/x*y[x]==(3*x^2*y[x]^2+6*x*y[x]+2)/(x^2*(2*x*y[x]+3)),y[2]==2},y[x],x,Include
```

$$y(x) \rightarrow \frac{\sqrt{\frac{1}{x^2} \sqrt{x^2(60x + 1)} - 3}}{2x}$$

5.54 problem 53

Internal problem ID [1028]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 53.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$y' + \frac{3y}{x} - \frac{3x^4y^2 + 10x^2y + 6}{x^3(2x^2y + 5)} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)+3/x*y(x)=(3*x^4*y(x)^2+10*x^2*y(x)+6)/(x^3*(2*x^2*y(x)+5)),y(1) = 1],y(x))
```

$$y(x) = \frac{-5 + \sqrt{48x + 1}}{2x^2}$$

✓ Solution by Mathematica

Time used: 0.815 (sec). Leaf size: 37

```
DSolve[{y'[x]+3/x*y[x]==(3*x^4*y[x]^2+10*x^2*y[x]+6)/(x^3*(2*x^2*y[x]+5)),y[1]==1},y[x],x,Integrate]
```

$$y(x) \rightarrow \frac{\sqrt{\frac{1}{x^2}} \sqrt{x^4(48x + 1)} - 5x}{2x^3}$$

5.55 problem 56

Internal problem ID [1029]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Transformation of Nonlinear Equations into Separable Equations. Section 2.4 Page 68

Problem number: 56.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [Riccati]

$$y' + (1 + 2x)y - xy^2 = x + 1$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x)=1+x-(1+2*x)*y(x)+x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{(2x + 4)e^{-x} - c_1}{(2 + 2x)e^{-x} - c_1}$$

✓ Solution by Mathematica

Time used: 0.149 (sec). Leaf size: 31

```
DSolve[y'[x]==1+x-(1+2*x)*y[x]+x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x + c_1 e^x + 2}{x + c_1 e^x + 1}$$
$$y(x) \rightarrow 1$$

6 Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

6.1	problem 1	182
6.2	problem 2	183
6.3	problem 3	184
6.4	problem 4	185
6.5	problem 5	187
6.6	problem 6	188
6.7	problem 7	190
6.8	problem 8	192
6.9	problem 9	193
6.10	problem 10	194
6.11	problem 11	195
6.12	problem 12	196
6.13	problem 13	197
6.14	problem 14	198
6.15	problem 15	199
6.16	problem 16	200
6.17	problem 17	201
6.18	problem 18	202
6.19	problem 19	203
6.20	problem 20	204
6.21	problem 21	206
6.22	problem 22	207
6.23	problem 23	208
6.24	problem 24	209
6.25	problem 25	210
6.26	problem 26	212
6.27	problem 27(a)	213
6.28	problem 28(a)	215
6.29	problem 38	216
6.30	problem 39	217
6.31	problem 40	218

6.1 problem 1

Internal problem ID [1030]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$6x^2y^2 + 4y'yx^3 = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(6*x^2*y(x)^2+4*x^3*y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = -\frac{\sqrt{2}\sqrt{-c_1x}}{2x^2}$$
$$y(x) = \frac{\sqrt{2}\sqrt{-c_1x}}{2x^2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 23

```
DSolve[6*x^2*y[x]^2+4*x^3*y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$
$$y(x) \rightarrow \frac{c_1}{x^{3/2}}$$
$$y(x) \rightarrow 0$$

6.2 problem 2

Internal problem ID [1031]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$3y \cos(x) + 2x^3y + (3 \sin(x) + 3) y' = -4x e^x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 395

```
dsolve((3*y(x)*cos(x)+4*x*exp(x)+2*x^3*y(x))+(3*sin(x)+3)*diff(y(x),x)=0,y(x), singsol=all)
```

$y(x)$

$$= \frac{e^{\frac{(-24x+24ix e^{ix}) \operatorname{polylog}(2, ie^{ix}) + (-24i-24 e^{ix}) \operatorname{polylog}(3, ie^{ix}) + 4ix(x^2 + \frac{3}{4})e^{ix} - 3x}{3 e^{ix} + 3i}}}{4 \int \frac{-2ie^{\frac{24x(e^{ix}+i) \operatorname{polylog}(2, ie^{ix}) + 4e^{ix}x^3 + 3ix e^{ix} + 2}{3ie^{ix}}}}{dx}}$$

✓ Solution by Mathematica

Time used: 27.47 (sec). Leaf size: 193

```
DSolve[(3*y[x]*Cos[x]+4*x*Exp[x]+2*x^3*y[x])+(3*SIN[x]+3)*y'[x]==0,y[x],x,IncludeSingularSol
```

$y(x)$

$$\rightarrow (1 + ie^{-ix})^{-4x^2} \exp(-8ix \operatorname{PolyLog}(2, -ie^{-ix}) - 8 \operatorname{PolyLog}(3, -ie^{-ix}) + \frac{2}{3}x^3(\cot(\frac{1}{4}(2x + \pi)) - i)) \left(\int_1^x \dots \right)$$

6.3 problem 3

Internal problem ID [1032]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$14y^3x^2 + 21x^2y^2y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve((14*x^2*y(x)^3)+(21*x^2*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = c_1 e^{-\frac{2x}{3}}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 25

```
DSolve[(14*x^2*y[x]^3)+(21*x^2*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$
$$y(x) \rightarrow c_1 e^{-2x/3}$$
$$y(x) \rightarrow 0$$

6.4 problem 4

Internal problem ID [1033]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$-2y^2 + (12y^2 - 4yx) y' = -2x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 429

```
dsolve((2*x-2*y(x)^2)+(12*y(x)^2-4*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-6x^5 - 6c_1x^3 + 81x^4 + 162c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}{6} + \frac{x}{6 \left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-6x^5 - 6c_1x^3 + 81x^4 + 162c_1x^2 + 81c_1^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{\left(-\left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}} + x\right) \left(i \left(\left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}}\right)\right)}{12 \left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{\left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}} (i\sqrt{3} - 1)}{12} - \frac{x \left(i x \sqrt{3} + x - 2 \left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}}\right)}{12 \left(-27x^2 - 27c_1 + x^3 + 3\sqrt{-3(x^2 + c_1)(2x^3 - 27x^2 - 27c_1)}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 4.8 (sec). Leaf size: 414

`DSolve[(2*x-2*y[x]^2)+(12*y[x]^2-4*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -\frac{x^2}{3^{2/3} \sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}$$

$$-\frac{\sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}{6\sqrt[3]{2}} + \frac{x}{6}$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3})x^2}{6^{2/3} \sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}$$

$$+ \frac{(1 - i\sqrt{3}) \sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}{12\sqrt[3]{2}} + \frac{x}{6}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3})x^2}{6^{2/3} \sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}$$

$$+ \frac{(1 + i\sqrt{3}) \sqrt[3]{-2x^3 + 54x^2 + \sqrt{-4x^6 + 4(x^3 - 27x^2 - 54c_1)^2 + 108c_1}}}{12\sqrt[3]{2}} + \frac{x}{6}$$

6.5 problem 5

Internal problem ID [1034]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$(x + y)^2 + (x + y)^2 y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
dsolve((x+y(x))^2+(x+y(x))^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x$$

$$y(x) = c_1 - x$$

$$y(x) = -\frac{c_1}{2} - \frac{i\sqrt{3}c_1}{2} - x$$

$$y(x) = -\frac{c_1}{2} + \frac{i\sqrt{3}c_1}{2} - x$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 18

```
DSolve[(x+y[x])^2+(x+y[x])^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x$$

$$y(x) \rightarrow -x + c_1$$

6.6 problem 6

Internal problem ID [1035]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$7y + (3x + 4y)y' = -4x$$

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 54

```
dsolve((4*x+7*y(x))+(3*x+4*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x \left(-2 \operatorname{RootOf} \left(_Z^{36} + 3 _Z^6 c_1 x^6 - 2 c_1 x^6 \right)^6 + 1 \right)}{\operatorname{RootOf} \left(_Z^{36} + 3 _Z^6 c_1 x^6 - 2 c_1 x^6 \right)^6}$$

✓ Solution by Mathematica

Time used: 3.222 (sec). Leaf size: 409

```
DSolve[(4*x+7*y[x])+(3*x+4*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 1\right]$$

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 2\right]$$

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 3\right]$$

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 4\right]$$

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 5\right]$$

$$y(x) \rightarrow \text{Root}\left[2\#1^6 + 21\#1^5x + 90\#1^4x^2 + 200\#1^3x^3 + 240\#1^2x^4 + 144\#1x^5 + 32x^6 - e^{3c_1}\&, 6\right]$$

6.7 problem 7

Internal problem ID [1036]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact]

$$-2 \sin(x) y^2 + 3y^3 + (4 \cos(x) y + 9xy^2) y' = 2x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 638

`dsolve((-2*y(x)^2*sin(x)+3*y(x)^3-2*x)+(4*y(x)*cos(x)+9*x*y(x)^2)*diff(y(x),x)=0,y(x), sings`

$$y(x) = \frac{\left(972x^4 + 36\sqrt{3} \sqrt{(-x^2 + c_1)(-243x^4 + 32 \cos(x)^3 + 243c_1x^2)} x - 64 \cos(x)^3 - 972c_1x^2\right)^{\frac{1}{3}}}{2} + \frac{8 \cos(x)^2}{9x}$$

$$y(x) = \frac{\left(\left(972x^4 + 36\sqrt{3} \sqrt{32 \cos(x)^3 (-x^2 + c_1) + 243(x^2 - c_1)^2 x^2 x - 64 \cos(x)^3 - 972c_1x^2}\right)^{\frac{1}{3}} + 4 \cos(x)\right)}{9x}$$

$$y(x) = \frac{\left(i \left(\left(972x^4 + 36\sqrt{3} \sqrt{32 \cos(x)^3 (-x^2 + c_1) + 243(x^2 - c_1)^2 x^2 x - 64 \cos(x)^3 - 972c_1x^2}\right)^{\frac{1}{3}} - 4 \cos(x)\right)\right)}{9x}$$

✓ Solution by Mathematica

Time used: 34.362 (sec). Leaf size: 520

`DSolve[(-2*y[x]^2*Sin[x]+3*y[x]^3-2*x)+(4*y[x]*Cos[x]+9*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSin`

$$y(x) \rightarrow \frac{2^{2/3} \sqrt[3]{-16 \cos^3(x) + 9 \left(27x^4 + 27c_1x^2 + \sqrt{3} \sqrt{x^2(x^2 + c_1)} (-32 \cos^3(x) + 243x^2(x^2 + c_1)) \right)} + \sqrt[3]{-8c_1}}{18x}$$

$$y(x) \rightarrow \frac{i 2^{2/3} (\sqrt{3} + i) \sqrt[3]{-16 \cos^3(x) + 9 \left(27x^4 + 27c_1x^2 + \sqrt{3} \sqrt{x^2(x^2 + c_1)} (-32 \cos^3(x) + 243x^2(x^2 + c_1)) \right)}}{36}$$

$$y(x) \rightarrow \frac{2^{2/3} (1 + i\sqrt{3}) \sqrt[3]{-16 \cos^3(x) + 9 \left(27x^4 + 27c_1x^2 + \sqrt{3} \sqrt{x^2(x^2 + c_1)} (-32 \cos^3(x) + 243x^2(x^2 + c_1)) \right)}}{36}$$

6.8 problem 8

Internal problem ID [1037]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$y + (2y + 2x)y' = -2x$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 51

```
dsolve((2*x+y(x))+(2*y(x)+2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x(\sqrt{7} \tan(\text{RootOf}(\sqrt{7} \ln(\sec(_Z)^2 x^2) + \sqrt{7} \ln(7) - 3\sqrt{7} \ln(2) + 2\sqrt{7} c_1 + 2_Z)) - 3)}{4}$$

✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 62

```
DSolve[(2*x+y[x])+(2*y[x]+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{\arctan\left(\frac{4y(x)+3}{\sqrt{7}}\right)}{2\sqrt{7}} + \frac{1}{4} \log\left(\frac{2y(x)^2}{x^2} + \frac{3y(x)}{x} + 2\right) = -\frac{\log(x)}{2} + c_1, y(x) \right]$$

6.9 problem 9

Internal problem ID [1038]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_Abel, '2nd type', 'class B']]

$$2yx + 4y^2 + (x^2 + 8yx + 18y) y' = -3x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 77

```
dsolve((3*x^2+2*x*y(x)+4*y(x)^2)+(x^2+8*x*y(x)+18*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-x^2 + \sqrt{-15x^4 - 36x^3 - 16c_1x - 36c_1}}{8x + 18}$$

$$y(x) = \frac{-x^2 - \sqrt{-15x^4 - 36x^3 - 16c_1x - 36c_1}}{8x + 18}$$

✓ Solution by Mathematica

Time used: 0.616 (sec). Leaf size: 84

```
DSolve[(3*x^2+2*x*y[x]+4*y[x]^2)+(x^2+8*x*y[x]+18*y[x])*y'[x]==0,y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\frac{x^2 + \sqrt{-15x^4 - 36x^3 + 16c_1x + 36c_1}}{8x + 18}$$

$$y(x) \rightarrow -\frac{x^2 - \sqrt{-15x^4 - 36x^3 + 16c_1x + 36c_1}}{8x + 18}$$

6.10 problem 10

Internal problem ID [1039]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational]

$$8yx + y^2 + \left(2x^2 + \frac{xy^3}{3}\right)y' = -2x^2$$

X Solution by Maple

```
dsolve((2*x^2+8*x*y(x)+y(x)^2)+(2*x^2+x*y(x)^3/3)*diff(y(x),x)=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(2*x^2+8*x*y[x]+y[x]^2)+(2*x^2+x*y[x]^3/3)*y'[x]==0,y[x],x,IncludeSingularSolutions -
```

Not solved

6.11 problem 11

Internal problem ID [1040]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$\left(\frac{1}{y} + 2y\right) y' = -\frac{1}{x} - 2x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve((1/x+2*x)+(1/y(x)+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-x^2-c_1} \sqrt{2}}{2 \sqrt{\frac{e^{-2x^2-2c_1}}{x^2 \text{LambertW}\left(\frac{2e^{-2x^2-2c_1}}{x^2}\right)} x}}$$

✓ Solution by Mathematica

Time used: 9.517 (sec). Leaf size: 71

```
DSolve[(1/x+2*x)+(1/y[x]+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{W\left(\frac{2e^{-2x^2+2c_1}}{x^2}\right)}}{\sqrt{2}}$$
$$y(x) \rightarrow \frac{\sqrt{W\left(\frac{2e^{-2x^2+2c_1}}{x^2}\right)}}{\sqrt{2}}$$
$$y(x) \rightarrow 0$$

6.12 problem 12

Internal problem ID [1041]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [NONE]

$$\sin(x)y^2 + xy^3 \cos(x) + (\sin(x)xy + xy^3 \cos(x))y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 5

```
dsolve((y(x)^2*sin(x)+x*y(x)^3*cos(x))+(x*sin(x)*y(x)+x*y(x)^3*cos(x))*diff(y(x),x)=0,y(x),
```

$$y(x) = 0$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y[x]^2*Sin[x]+x*y[x]^3*Cos[x])+(x*Sin[x]*y[x]+x*y[x]^3*Cos[x])*y'[x]==0,y[x],x,Inclu
```

Not solved

6.13 problem 13

Internal problem ID [1042]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$\frac{x}{(x^2 + y^2)^{\frac{3}{2}}} + \frac{yy'}{(x^2 + y^2)^{\frac{3}{2}}} = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve((x/(x^2+y(x)^2)^(3/2))+y(x)/(x^2+y(x)^2)^(3/2))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x^2 + c_1}$$
$$y(x) = -\sqrt{-x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.094 (sec). Leaf size: 39

```
DSolve[(x/(x^2+y[x]^2)^(3/2))+y[x]/(x^2+y[x]^2)^(3/2))*y'[x]==0,y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\sqrt{-x^2 + 2c_1}$$
$$y(x) \rightarrow \sqrt{-x^2 + 2c_1}$$

6.14 problem 14

Internal problem ID [1043]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, [_Abel, '2nd type', 'class B']]

$$e^x(x^2y^2 + 2xy^2) + (2x^2y e^x + 2)y' = -6x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 67

```
dsolve((exp(x)*(x^2*y(x)^2+2*x*y(x)^2)+6*x)+(2*x^2*y(x)*exp(x)+2)*diff(y(x),x)=0,y(x), singularities)
```

$$y(x) = \frac{(-1 + \sqrt{-3e^x x^4 - e^x c_1 x^2 + 1}) e^{-x}}{x^2}$$
$$y(x) = \frac{(-1 - \sqrt{-3e^x x^4 - e^x c_1 x^2 + 1}) e^{-x}}{x^2}$$

✓ Solution by Mathematica

Time used: 36.472 (sec). Leaf size: 76

```
DSolve[(Exp[x]*(x^2*y[x]^2+2*x*y[x]^2)+6*x)+(2*x^2*y[x]*Exp[x]+2)*y'[x]==0,y[x],x,IncludeSingularities]
```

$$y(x) \rightarrow -\frac{e^{-x} \left(1 + \sqrt{1 + e^x (-3x^4 + c_1 x^2)} \right)}{x^2}$$
$$y(x) \rightarrow \frac{e^{-x} \left(-1 + \sqrt{1 + e^x (-3x^4 + c_1 x^2)} \right)}{x^2}$$

6.15 problem 15

Internal problem ID [1044]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [exact]

$$x^2 e^{x^2+y} (2x^2 + 3) + (x^3 e^{x^2+y} - 12y^2) y' = -4x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve((x^2*exp(x^2+y(x))*(2*x^2+3)+4*x)+(x^3*exp(x^2+y(x))-12*y(x)^2)*diff(y(x),x)=0,y(x),
```

$$x^3 e^{x^2+y(x)} - 4y(x)^3 + 2x^2 + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.428 (sec). Leaf size: 30

```
DSolve[(x^2*Exp[x^2+y[x]]*(2*x^2+3)+4*x)+(x^3*Exp[x^2+y[x]]-12*y[x]^2)*y'[x]==0,y[x],x,Inclu
```

$$\text{Solve}\left[2x^2 + x^3 e^{x^2+y(x)} - 4y(x)^3 = c_1, y(x)\right]$$

6.16 problem 16

Internal problem ID [1045]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, [_1st_order, ‘_with_symmetry_[F(x),G(x)*y+H(x)]’]]

$$e^{yx}(yx^4 + 4x^3) + 3y + (x^5e^{yx} + 3x)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve((exp(x*y(x))*(x^4*y(x)+4*x^3)+3*y(x))+( x^5*exp(x*y(x))+3*x)*diff(y(x),x)=0,y(x), sin
```

$$y(x) = \frac{-3 \operatorname{LambertW}\left(\frac{x^4 e^{-\frac{c_1}{3}}}{3}\right) - c_1}{3x}$$

✓ Solution by Mathematica

Time used: 4.486 (sec). Leaf size: 33

```
DSolve[(Exp[x*y[x]]*(x^4*y[x]+4*x^3)+3*y[x])+(x^5*Exp[x*y[x]]+3*x)*y'[x]==0,y[x],x,IncludeSi
```

$$y(x) \rightarrow \frac{c_1 - 3W\left(\frac{1}{3}e^{\frac{c_1}{3}}x^4\right)}{3x}$$

6.17 problem 17

Internal problem ID [1046]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Abel, '2nd type', 'class B']]`

$$3 \cos(x) y x^2 - x^3 y^2 \sin(x) + (8y - x^4 \sin(x) y) y' = -4x$$

X Solution by Maple

```
dsolve((3*x^2*cos(x)*y(x)-x^3*y(x)*sin(x)*y(x)+4*x)+(8*y(x)-x^4*sin(x)*y(x))*diff(y(x),x)=0,
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(3*x^2*Cos[x]*y[x]-x^3*y[x]*Sin[x]*y[x]+4*x)+(8*y[x]-x^4*SIn[x]*y[x])*y'[x]==0,y[x],x
```

Not solved

6.18 problem 18

Internal problem ID [1047]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_exact, _rational, [_Abel, '2nd type', 'class B']]

$$4x^3y^2 - 6x^2y + (2yx^4 - 2x^3)y' = 2x + 3$$

With initial conditions

$$[y(1) = 3]$$

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 22

```
dsolve([(4*x^3*y(x)^2-6*x^2*y(x)-2*x-3)+(2*x^4*y(x)-2*x^3)*diff(y(x),x)=0,y(1) = 3],y(x), si
```

$$y(x) = \frac{x + \sqrt{2x^2 + 3x - 1}}{x^2}$$

✓ Solution by Mathematica

Time used: 0.743 (sec). Leaf size: 31

```
DSolve[{(4*x^3*y[x]^2-6*x^2*y[x]-2*x-3)+(2*x^4*y[x]-2*x^3)*y'[x]==0,y[1]==3},y[x],x,IncludeS
```

$$y(x) \rightarrow \frac{x^3 + \sqrt{x^4(2x^2 + 3x - 1)}}{x^4}$$

6.19 problem 19

Internal problem ID [1048]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, [_1st_order, ' _with_symmetry_[F(x),G(x)] '], [_Abel, ']`

$$-4 \cos(x) y + (4y - 4 \sin(x)) y' = -4 \cos(x) \sin(x) - \sec(x)^2$$

With initial conditions

$$\left[y\left(\frac{\pi}{4}\right) = 0 \right]$$

✓ Solution by Maple

Time used: 1.719 (sec). Leaf size: 32

```
dsolve([(-4*y(x)*cos(x)+4*sin(x)*cos(x)+sec(x)^2)+(4*y(x)-4*sin(x))*diff(y(x),x)=0,y(1/4*Pi)
```

$$y(x) = \sin(x) - \frac{\sec(x)^2 \sqrt{2} \sqrt{\cos(x)^3 (2 \cos(x) - \sin(x))}}{2}$$

✓ Solution by Mathematica

Time used: 11.693 (sec). Leaf size: 38

```
DSolve[{-4*y[x]*Cos[x]+4*Sin[x]*Cos[x]+Sec[x]^2)+(4*y[x]-4*Sin[x])*y'[x]==0,y[Pi/4]==0},y[x]
```

$$y(x) \rightarrow \sin(x) + \frac{1}{2} \sqrt{-\sec^2(x)} \sqrt{\sin(2x) - 2 \cos(2x) - 2}$$

6.20 problem 20

Internal problem ID [1049]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$(y^3 - 1)e^x + 3y^2(1 + e^x)y' = 0$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 91

```
dsolve([(y(x)^3-1)*exp(x))+(3*y(x)^2*(exp(x)+1))*diff(y(x),x)=0,y(0) = 0],y(x), singsol=all
```

$$y(x) = \frac{((e^x - 1)(1 + e^x)^2)^{\frac{1}{3}}}{1 + e^x}$$

$$y(x) = \frac{(i\sqrt{3} - 1)((e^x - 1)(1 + e^x)^2)^{\frac{1}{3}}}{2 + 2e^x}$$

$$y(x) = -\frac{(1 + i\sqrt{3})((e^x - 1)(1 + e^x)^2)^{\frac{1}{3}}}{2 + 2e^x}$$

✓ Solution by Mathematica

Time used: 0.984 (sec). Leaf size: 77

```
DSolve[{((y[x]^3-1)*Exp[x])+(3*y[x]^2*(Exp[x]+1))*y'[x]==0,y[0]==0},y[x],x,IncludeSingularSo
```

$$y(x) \rightarrow \sqrt[3]{\frac{e^x - 1}{e^x + 1}}$$

$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{\frac{e^x - 1}{e^x + 1}}$$

$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{\frac{e^x - 1}{e^x + 1}}$$

6.21 problem 21

Internal problem ID [1050]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$-\sin(x)y + y' \cos(x) = -\sin(x) + 2 \cos(x)$$

With initial conditions

$$[y(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve([(sin(x)-y(x)*sin(x)-2*cos(x))+(cos(x))*diff(y(x),x)=0,y(0) = 1],y(x), singsol=all)
```

$$y(x) = 2 \tan(x) + 1$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 11

```
DSolve[{(Sin[x]-y[x]*Sin[x]-2*Cos[x])+(Cos[x])*y'[x]==0,y[0]==1},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow 2 \tan(x) + 1$$

6.22 problem 22

Internal problem ID [1051]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(2x - 1)(y - 1) + (2 + x)(x - 3)y' = 0$$

With initial conditions

$$[y(1) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve([(2*x-1)*(y(x)-1))+((x+2)*(x-3))*diff(y(x),x)=0,y(1) = -1],y(x), singsol=all)
```

$$y(x) = \frac{x^2 - x + 6}{(2 + x)(x - 3)}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 24

```
DSolve[{((2*x-1)*(y[x]-1))+((x+2)*(x-3))*y'[x]==0,y[1]==-1},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{x^2 - x + 6}{x^2 - x - 6}$$

6.23 problem 23

Internal problem ID [1052]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, [_Abel, '2nd ty`

$$4y + (4x + 3y)y' = -7x$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 53

```
dsolve((7*x+4*y(x))+(4*x+3*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-4c_1x - \sqrt{-5c_1^2x^2 + 3}}{3c_1}$$
$$y(x) = \frac{-4c_1x + \sqrt{-5c_1^2x^2 + 3}}{3c_1}$$

✓ Solution by Mathematica

Time used: 0.483 (sec). Leaf size: 118

```
DSolve[(7*x+4*y[x])+(4*x+3*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \left(-4x - \sqrt{-5x^2 + 3e^{2c_1}} \right)$$
$$y(x) \rightarrow \frac{1}{3} \left(-4x + \sqrt{-5x^2 + 3e^{2c_1}} \right)$$
$$y(x) \rightarrow \frac{1}{3} \left(-\sqrt{5}\sqrt{-x^2} - 4x \right)$$
$$y(x) \rightarrow \frac{1}{3} \left(\sqrt{5}\sqrt{-x^2} - 4x \right)$$

6.24 problem 24

Internal problem ID [1053]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_exact`, `_Bernoulli`]

$$e^x(x^4y^2 + 4x^3y^2 + 1) + (2x^4ye^x + 2y)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 64

```
dsolve((exp(x)*(x^4*y(x)^2+4*x^3*y(x)^2+1))+(2*x^4*y(x)*exp(x)+2*y(x))*diff(y(x),x)=0,y(x),
```

$$y(x) = \frac{\sqrt{(e^xx^4 + 1)(-e^x + c_1)}}{e^xx^4 + 1}$$
$$y(x) = -\frac{\sqrt{(e^xx^4 + 1)(-e^x + c_1)}}{e^xx^4 + 1}$$

✓ Solution by Mathematica

Time used: 1.05 (sec). Leaf size: 64

```
DSolve[(Exp[x]*(x^4*y[x]^2+4*x^3*y[x]^2+1))+(2*x^4*y[x]*Exp[x]+2*y[x])*y'[x]==0,y[x],x,Inclu
```

$$y(x) \rightarrow -\frac{\sqrt{-2e^x + c_1}}{\sqrt{2e^xx^4 + 2}}$$
$$y(x) \rightarrow \frac{\sqrt{-2e^x + c_1}}{\sqrt{2e^xx^4 + 2}}$$

6.25 problem 25

Internal problem ID [1054]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$x^3y^4 + (y^3x^4 + y)y' = -x$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 111

```
dsolve((x^3*y(x)^4+x)+(x^4*y(x)^3+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-1 - \sqrt{-2x^6 - 4c_1x^4 + 1}}}{x^2}$$

$$y(x) = \frac{\sqrt{-1 + \sqrt{-2x^6 - 4c_1x^4 + 1}}}{x^2}$$

$$y(x) = -\frac{\sqrt{-1 - \sqrt{-2x^6 - 4c_1x^4 + 1}}}{x^2}$$

$$y(x) = -\frac{\sqrt{-1 + \sqrt{-2x^6 - 4c_1x^4 + 1}}}{x^2}$$

✓ Solution by Mathematica

Time used: 11.648 (sec). Leaf size: 135

```
DSolve[(x^3*y[x]^4+x)+(x^4*y[x]^3+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{-\frac{1 + \sqrt{-2x^6 + 4c_1x^4 + 1}}{x^4}}$$

$$y(x) \rightarrow \sqrt{-\frac{1 + \sqrt{-2x^6 + 4c_1x^4 + 1}}{x^4}}$$

$$y(x) \rightarrow -\sqrt{\frac{-1 + \sqrt{-2x^6 + 4c_1x^4 + 1}}{x^4}}$$

$$y(x) \rightarrow \sqrt{\frac{-1 + \sqrt{-2x^6 + 4c_1x^4 + 1}}{x^4}}$$

6.26 problem 26

Internal problem ID [1055]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_exact, _rational, [_1st_order, ‘_with_symmetry_[F(x),G(x)]’]`

$$2y + (2y + 2x)y' = -3x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve((3*x^2+2*y(x))+(2*y(x)+2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x - \sqrt{-x^3 + x^2 - c_1}$$
$$y(x) = -x + \sqrt{-x^3 + x^2 - c_1}$$

✓ Solution by Mathematica

Time used: 0.131 (sec). Leaf size: 49

```
DSolve[(3*x^2+2*y[x])+(2*y[x]+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{-x^3 + x^2 + c_1}$$
$$y(x) \rightarrow -x + \sqrt{-x^3 + x^2 + c_1}$$

6.27 problem 27(a)

Internal problem ID [1056]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 27(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_exact`, `_rational`]

$$x^3y^4 + (y^3x^4 + 3y)y' = -2x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 111

```
dsolve((x^3*y(x)^4+2*x)+(x^4*y(x)^3+3*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{-3 - \sqrt{-4x^6 - 4c_1x^4 + 9}}}{x^2}$$

$$y(x) = \frac{\sqrt{-3 + \sqrt{-4x^6 - 4c_1x^4 + 9}}}{x^2}$$

$$y(x) = -\frac{\sqrt{-3 - \sqrt{-4x^6 - 4c_1x^4 + 9}}}{x^2}$$

$$y(x) = -\frac{\sqrt{-3 + \sqrt{-4x^6 - 4c_1x^4 + 9}}}{x^2}$$

✓ Solution by Mathematica

Time used: 11.546 (sec). Leaf size: 135

```
DSolve[(x^3*y[x]^4+2*x)+(x^4*y[x]^3+3*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -\sqrt{-\frac{3 + \sqrt{-4x^6 + 4c_1x^4 + 9}}{x^4}}$$

$$y(x) \rightarrow \sqrt{-\frac{3 + \sqrt{-4x^6 + 4c_1x^4 + 9}}{x^4}}$$

$$y(x) \rightarrow -\sqrt{\frac{-3 + \sqrt{-4x^6 + 4c_1x^4 + 9}}{x^4}}$$

$$y(x) \rightarrow \sqrt{\frac{-3 + \sqrt{-4x^6 + 4c_1x^4 + 9}}{x^4}}$$

6.28 problem 28(a)

Internal problem ID [1057]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 28(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _exact, _rational, _Bernoulli]`

$$y^2 + 2y'xy = -x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve((x^2+y(x)^2)+(2*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{3} \sqrt{-x(x^3 - 3c_1)}}{3x}$$
$$y(x) = \frac{\sqrt{3} \sqrt{-x(x^3 - 3c_1)}}{3x}$$

✓ Solution by Mathematica

Time used: 0.22 (sec). Leaf size: 60

```
DSolve[(x^2+y[x]^2)+(2*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$
$$y(x) \rightarrow \frac{\sqrt{-x^3 + 3c_1}}{\sqrt{3}\sqrt{x}}$$

6.29 problem 38

Internal problem ID [1058]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 38.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$y' + \frac{2y}{x} + \frac{2xy}{x^2 + 2x^2y + 1} = 0$$

With initial conditions

$$[y(1) = -2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)+2/x*y(x)= -(2*x*y(x))/(x^2+2*x^2*y(x)+1),y(1) = -2],y(x), singsol=all)
```

$$y(x) = \frac{-x^2 - 1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.694 (sec). Leaf size: 38

```
DSolve[{y'[x]+2/x*y[x]== -(2*x*y[x])/(x^2+2*x^2*y[x]+1),y[1]==-2},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \frac{1}{2} \left(-\frac{1}{x^2} - \frac{\sqrt{x^3(x^2+1)^2}}{x^{7/2}} - 1 \right)$$

6.30 problem 39

Internal problem ID [1059]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 39.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$y' - \frac{3y}{x} - \frac{2x^4(4x^3 - 3y)}{3x^5 + 3x^3 + 2y} = 0$$

With initial conditions

$$[y(1) = 1]$$

✓ Solution by Maple

Time used: 0.109 (sec). Leaf size: 30

```
dsolve([diff(y(x),x)-3/x*y(x)= (2*x^4*(4*x^3-3*y(x)))/(3*x^5+3*x^3+2*y(x)),y(1) = 1],y(x), s
```

$$y(x) = \frac{(-3x^2 + \sqrt{9x^4 + 34x^2 + 21} - 3)x^3}{2}$$

✓ Solution by Mathematica

Time used: 0.734 (sec). Leaf size: 49

```
DSolve[{y'[x]-3/x*y[x]== (2*x^4*(4*x^3-3*y[x]))/(3*x^5+3*x^3+2*y[x]),y[1]==1},y[x],x,Include
```

$$y(x) \rightarrow \frac{1}{2} \left(-3x^5 - 3x^3 + \sqrt{\frac{1}{x^7} \sqrt{x(9x^4 + 34x^2 + 21)} x^6} \right)$$

6.31 problem 40

Internal problem ID [1060]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Section 2.5 Page 79

Problem number: 40.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_Abel, '2nd type', 'class B']]`

$$y' + 2yx + \frac{e^{-x^2}(3x + 2y e^{x^2})}{2x + 3y e^{x^2}} = 0$$

With initial conditions

$$[y(0) = -1]$$

✓ Solution by Maple

Time used: 0.312 (sec). Leaf size: 36

```
dsolve([diff(y(x),x)+2*x*y(x)= -exp(-x^2)*(3*x+2*y(x)*exp(x^2))/(2*x+3*y(x)*exp(x^2)),y(0) =
```

$$y(x) = -\frac{\left(2x e^{x^2} + \sqrt{e^{2x^2}(-5x^2 + 9)}\right) e^{-2x^2}}{3}$$

✓ Solution by Mathematica

Time used: 33.105 (sec). Leaf size: 44

```
DSolve[{y'[x]+2*x*y[x]== -Exp[-x^2]*(3*x+2*y[x]*Exp[x^2])/(2*x+3*y[x]*Exp[x^2]),y[0]==-1},y[
```

$$y(x) \rightarrow -\frac{1}{3}e^{-2x^2} \left(2e^{x^2} x + \sqrt{e^{2x^2} (9 - 5x^2)}\right)$$

7 Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6 Page 91

7.1	problem 1(a)	220
7.2	problem 2(a)	221
7.3	problem 3	222
7.4	problem 4	223
7.5	problem 5	224
7.6	problem 6	225
7.7	problem 7	226
7.8	problem 8	227
7.9	problem 9	230
7.10	problem 10	231
7.11	problem 11	232
7.12	problem 12	234
7.13	problem 13	235
7.14	problem 14	236
7.15	problem 15	237
7.16	problem 16	238
7.17	problem 18	239
7.18	problem 19	240
7.19	problem 20	241
7.20	problem 21	242
7.21	problem 22	243
7.22	problem 23	244
7.23	problem 24	245
7.24	problem 25	247
7.25	problem 26	249
7.26	problem 27	250

7.1 problem 1(a)

Internal problem ID [1061]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, [_Abel, '2nd type', 'cl`

$$y + \left(2x + \frac{1}{y}\right) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
dsolve(y(x)+(2*x+1/y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-1 + \sqrt{4c_1x + 1}}{2x}$$
$$y(x) = \frac{-1 - \sqrt{4c_1x + 1}}{2x}$$

✓ Solution by Mathematica

Time used: 0.321 (sec). Leaf size: 54

```
DSolve[y[x]+(2*x+1/y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1 + \sqrt{1 + 4c_1x}}{2x}$$
$$y(x) \rightarrow \frac{-1 + \sqrt{1 + 4c_1x}}{2x}$$
$$y(x) \rightarrow 0$$

7.2 problem 2(a)

Internal problem ID [1062]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$-y^2 + y'x^2 = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(-y(x)^2+x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{c_1x + 1}$$

✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 21

```
DSolve[-y[x]^2+x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{1 - c_1x}$$
$$y(x) \rightarrow 0$$

7.3 problem 3

Internal problem ID [1063]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$-y'x + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(y(x)-x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1x$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 14

```
DSolve[y[x]-x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x$$
$$y(x) \rightarrow 0$$

7.4 problem 4

Internal problem ID [1064]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$3x^2y + 2x^3y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(3*x^2*y(x)+2*x^3*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 18

```
DSolve[3*x^2*y[x]+2*x^3*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^{3/2}}$$
$$y(x) \rightarrow 0$$

7.5 problem 5

Internal problem ID [1065]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$2y^3 + 3y^2y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(2*y(x)^3+3*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = c_1 e^{-\frac{2x}{3}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 25

```
DSolve[2*y[x]^3+3*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$
$$y(x) \rightarrow c_1 e^{-2x/3}$$
$$y(x) \rightarrow 0$$

7.6 problem 6

Internal problem ID [1066]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$5yx + 2y + 2y'x = -5$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve((5*x*y(x)+2*y(x)+5)+(2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{5x}{2}} c_1 - 1}{x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 21

```
DSolve[(5*x*y[x]+2*y[x]+5)+(2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-1 + c_1 e^{-5x/2}}{x}$$

7.7 problem 7

Internal problem ID [1067]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [linear]

$$yx + 2y + (x + 1)y' = -x - 1$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve((x*y(x)+x+2*y(x)+1)+(x+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-x}c_1 - x}{x + 1}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 23

```
DSolve[(x*y[x]+x+2*y[x]+1)+(x+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x - c_1 e^{-x}}{x + 1}$$

7.8 problem 8

Internal problem ID [1068]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class A', _rational, [_Abel, '2nd type', 'cl`

$$27xy^2 + 8y^3 + (18x^2y + 12xy^2) y' = 0$$

✓ Solution by Maple

Time used: 0.344 (sec). Leaf size: 1355

`dsolve((27*x*y(x)^2+8*y(x)^3)+(18*x^2*y(x)+12*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = 0$$

$$y(x) = \frac{x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = \frac{x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = \frac{x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = \frac{x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = \frac{x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) =$$

$$\frac{2x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{i \left(-3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}} \right) \sqrt{3} + 3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = -\frac{2x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{(1 - i\sqrt{3}) \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}} + 3c_1 (1 + i\sqrt{3}) x^5}$$

$$y(x) =$$

$$\frac{2x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{i \left(-3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}} \right) \sqrt{3} + 3c_1 x^5 + \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}}}$$

$$y(x) = -\frac{2x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}{(1 - i\sqrt{3}) \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{2}{3}} + 3c_1 (1 + i\sqrt{3}) x^5}$$

$$y(x) =$$

$$\frac{228}{2x \left(2c_1 x^5 + \sqrt{-27c_1^3 x^{15} + 4c_1^2 x^{10}} \right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 59.109 (sec). Leaf size: 534

`DSolve[(27*x*y[x]^2+8*y[x]^3)+(18*x^2*y[x]+12*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolut`

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{1}{4} \left(\frac{9x^2}{\sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}}} + \sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}} - 3x \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(-\frac{9(1+i\sqrt{3})x^2}{\sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}}} + i(\sqrt{3}+i) \sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}} - 6x \right)$$

$$y(x) \rightarrow \frac{1}{8} \left(\frac{9i(\sqrt{3}+i)x^2}{\sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}}} - (1+i\sqrt{3}) \sqrt[3]{\frac{-27x^5 + 4\sqrt{e^{6c_1}(-27x^5 + 4e^{6c_1})} + 8e^{6c_1}}{x^2}} - 6x \right)$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{3\left(\sqrt[3]{-x^3} + x\right)\left(-2x + (1 - i\sqrt{3})\sqrt[3]{-x^3}\right)}{8x}$$

$$y(x) \rightarrow \frac{3\left(\sqrt[3]{-x^3} + x\right)\left(-2x + (1 + i\sqrt{3})\sqrt[3]{-x^3}\right)}{8x}$$

$$y(x) \rightarrow -\frac{3\left(-\sqrt[3]{-x^3}x + (-x^3)^{2/3}\right)}{4x}$$

7.9 problem 9

Internal problem ID [1069]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class B']]`

$$6xy^2 + 2y + (12x^2y + 12xy^2) y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 5

```
dsolve((6*x*y(x)^2+2*y(x))+(12*x^2*y(x)+12*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(6*x*y[x]^2+2*y[x])+(12*x^2*y[x]+12*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolution
```

Not solved

7.10 problem 10

Internal problem ID [1070]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6

Page 91

Problem number: 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ' _with_symmetry_[F(x)*G(y),0]']]`

$$y^2 + \left(xy^2 + 6yx + \frac{1}{y}\right)y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 38

```
dsolve((y(x)^2)+(x*y(x)^2+3*x*y(x)+3*x*y(x)+1/y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{xy(x)^6 + y(x)^3 - 3y(x)^2 - e^{-y(x)}c_1 + 6y(x) - 6}{y(x)^6} = 0$$

✓ Solution by Mathematica

Time used: 0.2 (sec). Leaf size: 41

```
DSolve[(y[x]^2)+(x*y[x]^2+3*x*y[x]+3*x*y[x]+1/y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions
```

$$\text{Solve} \left[x = -\frac{y(x)^3 - 3y(x)^2 + 6y(x) - 6}{y(x)^6} + \frac{c_1 e^{-y(x)}}{y(x)^6}, y(x) \right]$$

7.11 problem 11

Internal problem ID [1071]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 11.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_rational, [_Abel, '2nd type', 'class B']]

$$12yx^3 + 24x^2y^2 + (9x^4 + 32yx^3 + 4y)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve((12*x^3*y(x)+24*x^2*y(x)^2)+(9*x^4+32*x^3*y(x)+4*y(x))*diff(y(x),x)=0,y(x), singsol=a
```

$$3x^4y(x)^3 + 8x^3y(x)^4 + y(x)^4 + c_1 = 0$$

✓ Solution by Mathematica

Time used: 61.708 (sec). Leaf size: 1733

`DSolve[(12*x^3*y[x]+24*x^2*y[x]^2)+(9*x^4+32*x^3*y[x]+4*y[x])*y'[x]==0,y[x],x,IncludeSingularities->True]`

$$y(x) \rightarrow -\frac{3x^4}{32x^3 + 4}$$

$$+ \frac{1}{2} \sqrt{\frac{9x^8}{4(8x^3 + 1)^2} + \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} - \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

$$- \frac{1}{2} \sqrt{\frac{9x^8}{2(8x^3 + 1)^2} - \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} + \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

$$y(x) \rightarrow -\frac{3x^4}{32x^3 + 4}$$

$$+ \frac{1}{2} \sqrt{\frac{9x^8}{4(8x^3 + 1)^2} + \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} - \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

$$+ \frac{1}{2} \sqrt{\frac{9x^8}{2(8x^3 + 1)^2} - \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} + \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

$$y(x) \rightarrow -\frac{3x^4}{32x^3 + 4}$$

$$- \frac{1}{2} \sqrt{\frac{9x^8}{4(8x^3 + 1)^2} + \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} - \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

$$- \frac{1}{2} \sqrt{\frac{9x^8}{2(8x^3 + 1)^2} - \frac{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3} - 243c_1x^8}}{3\sqrt[3]{2}(8x^3 + 1)}} + \frac{4\sqrt[3]{2}c_1}{\sqrt[3]{\sqrt{59049c_1^2x^{16} + 6912c_1^3(8x^3 + 1)^3}}}$$

7.12 problem 12

Internal problem ID [1072]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 12.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x^2y + 4yx + 2y + (x^2 + x)y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((x^2*y(x)+4*x*y(x)+2*y(x))+(x^2+x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 e^{-x}}{(x+1)x^2}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 26

```
DSolve[(x^2*y[x]+4*x*y[x]+2*y[x])+(x^2+x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 e^{-x}}{x^2(x+1)}$$
$$y(x) \rightarrow 0$$

7.13 problem 13

Internal problem ID [1073]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 13.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$-y + (x^4 - x) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(-y(x)+(x^4-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1(x-1)^{\frac{1}{3}}(x^2+x+1)^{\frac{1}{3}}}{x}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 27

```
DSolve[-y[x]+(x^4-x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 \sqrt[3]{1-x^3}}{x}$$
$$y(x) \rightarrow 0$$

7.14 problem 14

Internal problem ID [1074]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 14.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_1st_order, '_with_symmetry_[F(x)*G(y),0]']]`

$$\cos(x) \cos(y) + (\sin(x) \cos(y) - \sin(x) \sin(y) + y) y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve((cos(x)*cos(y(x)))+(sin(x)*cos(y(x))-sin(x)*sin(y(x))+y(x))*diff(y(x),x)=0,y(x),sing
```

$$(\sin(x) \cos(y(x)) + y(x) - 1) e^{y(x)} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.218 (sec). Leaf size: 28

```
DSolve[(Cos[x]*Cos[y[x]])+(Sin[x]*Cos[y[x]]-Sin[x]*Sin[y[x]]+y[x])*y'[x]==0,y[x],x,IncludeSi
```

$$\text{Solve}[-2e^{y(x)}(y(x) - 1) - 2e^{y(x)} \sin(x) \cos(y(x)) = c_1, y(x)]$$

7.15 problem 15

Internal problem ID [1075]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [rational]

$$2yx + y^2 + (2yx + x^2 - 2xy^2 - 2xy^3) y' = 0$$

X Solution by Maple

```
dsolve((2*x*y(x)+y(x)^2)+(2*x*y(x)+x^2-2*x*y(x)^2-2*x*y(x)^3)*diff(y(x),x)=0,y(x), singsol=a
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(2*x*y[x]+y[x]^2)+(2*x*y[x]+x^2-2*x*y[x]^2-2*x*y[x]^3)*y'[x]==0,y[x],x,IncludeSingular
```

Not solved

7.16 problem 16

Internal problem ID [1076]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y \sin(y) + x(\sin(y) - y \cos(y)) y' = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 16

```
dsolve((y(x)*sin(y(x)))+(x*(sin(y(x))-y(x)*cos(y(x))))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\ln(x) + \ln(y(x)) - \ln(\sin(y(x))) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.569 (sec). Leaf size: 27

```
DSolve[(y[x]*Sin[y[x]])+(x*(Sin[y[x]]-y[x]*Cos[y[x]]))*y'[x]==0,y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow \text{InverseFunction}[\log(\sin(\#1)) - \log(\#1)\&][\log(x) + c_1]$$

$$y(x) \rightarrow 0$$

7.17 problem 18

Internal problem ID [1077]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$ay + bxy + (cx + dxy)y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 56

```
dsolve((a*y(x)+b*x*y(x))+(c*x+d*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = x^{-\frac{a}{c}} e^{\frac{-bx - c \operatorname{LambertW}\left(\frac{dx^{-\frac{a}{c}} e^{-bx+c_1}}{c}\right) - c_1}{c}}$$

✓ Solution by Mathematica

Time used: 1.133 (sec). Leaf size: 42

```
DSolve[(a*y[x]+b*x*y[x])+(c*x+d*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{cW\left(\frac{dx^{-\frac{a}{c}} e^{-bx+c_1}}{c}\right)}{d}$$
$$y(x) \rightarrow 0$$

7.18 problem 19

Internal problem ID [1078]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 19.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_Abel, '2nd type', 'class C']]`

$$3y^3x^2 - y^2 + y + (-yx + 2x)y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 59

```
dsolve((3*x^2*y(x)^3-y(x)^2+y(x))+(-x*y(x)+2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{4}{\sqrt{x} \sqrt{\frac{c_1x+48x^2+4}{x}} + 2}$$
$$y(x) = -\frac{4}{\sqrt{x} \sqrt{\frac{c_1x+48x^2+4}{x}} - 2}$$

✓ Solution by Mathematica

Time used: 0.776 (sec). Leaf size: 80

```
DSolve[(3*x^2*y[x]^3-y[x]^2+y[x])+(-x*y[x]+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{2}{1 + \sqrt{-\frac{1}{x^2}x\sqrt{-12x^2 - 4c_1x - 1}}}$$
$$y(x) \rightarrow \frac{2x}{x + \frac{\sqrt{-12x^2 - 4c_1x - 1}}{\sqrt{-\frac{1}{x^2}}}}$$
$$y(x) \rightarrow 0$$

7.19 problem 20

Internal problem ID [1079]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 20.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$2y + 3(x^2 + y^3x^2)y' = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 30

```
dsolve((2*y(x))+3*(x^2+x^2*y(x)^3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x \operatorname{LambertW}\left(e^{-\frac{-2c_1x+2}{x}}\right) + 2c_1x - 2}{3x}}$$

✓ Solution by Mathematica

Time used: 4.79 (sec). Leaf size: 82

```
DSolve[(2*y[x])+3*(x^2+x^2*y[x]^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{W\left(e^{\frac{2}{x}+3c_1}\right)}$$

$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{W\left(e^{\frac{2}{x}+3c_1}\right)}$$

$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{W\left(e^{\frac{2}{x}+3c_1}\right)}$$

$$y(x) \rightarrow 0$$

7.20 problem 21

Internal problem ID [1080]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 21.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_linear]`

$$a \cos(x) y - \sin(x) y^2 + (b \cos(x) y - \sin(x) x y) y' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 68

```
dsolve((a*cos(x)*y(x)-y(x)*sin(x)*y(x))+(b*cos(x)*y(x)-x*sin(x)*y(x))*diff(y(x),x)=0,y(x), s
```

$$y(x) = 0$$

$$y(x) = \left(a \left(\int \frac{\cos(x) e^{\int \frac{\sin(x)}{x \sin(x) - \cos(x) b} dx}}{x \sin(x) - \cos(x) b} dx \right) + c_1 \right) e^{-\left(\int \frac{\sin(x)}{x \sin(x) - \cos(x) b} dx \right)}$$

✓ Solution by Mathematica

Time used: 3.564 (sec). Leaf size: 106

```
DSolve[(a*Cos[x]*y[x]-y[x]*Sin[x]*y[x])+(b*Cos[x]*y[x]-x*SIn[x]*y[x])*y'[x]==0,y[x],x,Includ
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \exp \left(\int_1^x \frac{\sin(K[1])}{b \cos(K[1]) - K[1] \sin(K[1])} dK[1] \right) \left(\int_1^x \frac{a \exp \left(- \int_1^{K[2]} \frac{\sin(K[1])}{b \cos(K[1]) - K[1] \sin(K[1])} dK[1] \right) \cos(K[2])}{b \cos(K[2]) - K[2] \sin(K[2])} dK[2] + c_1 \right)$$

$$y(x) \rightarrow 0$$

7.21 problem 22

Internal problem ID [1081]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 22.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_separable]

$$x^4 y^4 + x^5 y^3 y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((x^4*y(x)^4)+(x^5*y(x)^3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 21

```
DSolve[(x^4*y[x]^4)+(x^5*y[x]^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{c_1}{x}$$

$$y(x) \rightarrow 0$$

7.22 problem 23

Internal problem ID [1082]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 23.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y(x \cos(x) + 2 \sin(x)) + x(1 + y) y' = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 18

```
dsolve((y(x)*(x*cos(x)+2*sin(x)))+(x*(y(x)+1))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{LambertW}(e^{-\sin(x)-2 \text{Si}(x)-c_1})$$

✓ Solution by Mathematica

Time used: 2.869 (sec). Leaf size: 24

```
DSolve[(y[x]*(x*Cos[x]+2*Sin[x]))+(x*(y[x]+1))*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow W(e^{-2\text{Si}(x)-\sin(x)+c_1})$$
$$y(x) \rightarrow 0$$

7.23 problem 24

Internal problem ID [1083]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 24.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$y^3 x^4 + y + (y^2 x^5 - x) y' = 0$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 242

```
dsolve((x^4*y(x)^3+y(x))+(x^5*y(x)^2-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{2}{3}} + 4c_1}{2x^2 \left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{1}{3}} \sqrt{c_1}}$$
$$y(x) = \frac{-i\left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{2}{3}} \sqrt{3} + 4i\sqrt{3} c_1 - \left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{2}{3}} - 4c_1}{4x^2 \left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{1}{3}} \sqrt{c_1}}$$
$$y(x) = -\frac{-i\left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{2}{3}} \sqrt{3} + 4i\sqrt{3} c_1 + \left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{2}{3}} + 4c_1}{4x^2 \left(4x^3 + 4\sqrt{x^6 - 4c_1^3}\right)^{\frac{1}{3}} \sqrt{c_1}}$$

✓ Solution by Mathematica

Time used: 39.244 (sec). Leaf size: 300

`DSolve[(x^4*y[x]^3+y[x])+(x^5*y[x]^2-x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{2 + \frac{\sqrt[3]{2}(3c_1x^9 + \sqrt{x^{12}(-4 + 9c_1^2x^6)})^{2/3}}{x^4}}{2^{2/3}\sqrt[3]{3c_1x^9 + \sqrt{x^{12}(-4 + 9c_1^2x^6)}}$$
$$y(x) \rightarrow \frac{i\left((\sqrt{3} + i)\left(6c_1x^9 + 2\sqrt{x^{12}(-4 + 9c_1^2x^6)}\right)^{2/3} - 2\sqrt[3]{2}(\sqrt{3} - i)x^4\right)}{4x^4\sqrt[3]{3c_1x^9 + \sqrt{x^{12}(-4 + 9c_1^2x^6)}}$$
$$y(x) \rightarrow \frac{2i\sqrt[3]{2}(\sqrt{3} + i)x^4 + (-1 - i\sqrt{3})\left(6c_1x^9 + 2\sqrt{x^{12}(-4 + 9c_1^2x^6)}\right)^{2/3}}{4x^4\sqrt[3]{3c_1x^9 + \sqrt{x^{12}(-4 + 9c_1^2x^6)}}$$

7.24 problem 25

Internal problem ID [1084]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6

Page 91

Problem number: 25.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class D', _rational, [_Abel, '2nd type', 'cl`

$$3yx + 2y^2 + y + (x^2 + 2yx + x + 2y) y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 85

```
dsolve((3*x*y(x)+2*y(x)^2+y(x))+(x^2+2*x*y(x)+x+2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-\sqrt{4 + x^2 (x + 1)^2} c_1^2 + (-x^2 - x) c_1}{2c_1 (x + 1)}$$

$$y(x) = \frac{\sqrt{4 + x^2 (x + 1)^2} c_1^2 + (-x^2 - x) c_1}{2c_1 (x + 1)}$$

✓ Solution by Mathematica

Time used: 14.424 (sec). Leaf size: 105

```
DSolve[(3*x*y[x]+2*y[x]^2+y[x])+(x^2+2*x*y[x]+x+2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{1}{2} \left(-x - \sqrt{x^2 + \frac{4e^{c_1}}{(x+1)^2}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-x + \sqrt{x^2 + \frac{4e^{c_1}}{(x+1)^2}} \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(-\sqrt{x^2} - x \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{x^2} - x \right)$$

7.25 problem 26

Internal problem ID [1085]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 26.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class G'], _rational]`

$$12yx + 6y^3 + (9x^2 + 10xy^2)y' = 0$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 35

```
dsolve((12*x*y(x)+6*y(x)^3)+(9*x^2+10*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$\ln(x) - c_1 + \frac{6 \ln\left(\frac{y(x)}{\sqrt{x}}\right)}{11} + \frac{2 \ln\left(\frac{2y(x)^2+3x}{x}\right)}{11} = 0$$

✓ Solution by Mathematica

Time used: 8.107 (sec). Leaf size: 151

```
DSolve[(12*x*y[x]+6*y[x]^3)+(9*x^2+10*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$\begin{aligned}y(x) &\rightarrow \text{Root}\left[2\#1^5x^3 + 3\#1^3x^4 - c_1\&, 1\right] \\y(x) &\rightarrow \text{Root}\left[2\#1^5x^3 + 3\#1^3x^4 - c_1\&, 2\right] \\y(x) &\rightarrow \text{Root}\left[2\#1^5x^3 + 3\#1^3x^4 - c_1\&, 3\right] \\y(x) &\rightarrow \text{Root}\left[2\#1^5x^3 + 3\#1^3x^4 - c_1\&, 4\right] \\y(x) &\rightarrow \text{Root}\left[2\#1^5x^3 + 3\#1^3x^4 - c_1\&, 5\right]\end{aligned}$$

7.26 problem 27

Internal problem ID [1086]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 2, First order equations. Exact equations. Integrating factors. Section 2.6
Page 91

Problem number: 27.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Bernoulli]`

$$3x^2y^2 + 2y + 2y'x = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve((3*x^2*y(x)^2+2*y(x))+(2*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2}{(3x + 2c_1)x}$$

✓ Solution by Mathematica

Time used: 0.146 (sec). Leaf size: 25

```
DSolve[(3*x^2*y[x]^2+2*y[x])+(2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3x^2 + 2c_1x}$$
$$y(x) \rightarrow 0$$

8 Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

8.1	problem 1	252
8.2	problem 2c	253
8.3	problem 2d	254
8.4	problem 3c	255
8.5	problem 3d	256
8.6	problem 4	257
8.7	problem 10	258
8.8	problem 11	259
8.9	problem 12	260
8.10	problem 13	261
8.11	problem 14	262
8.12	problem 15	263
8.13	problem 16	264
8.14	problem 17	265
8.15	problem 18	266
8.16	problem 19	267
8.17	problem 20	268
8.18	problem 21	269
8.19	problem 22	270
8.20	problem 23	271

8.1 problem 1

Internal problem ID [1087]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 7y' + 10y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([diff(y(x),x$2)-7*diff(y(x),x)+10*y(x)=0,y(0) = -1, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = e^{5x} - 2e^{2x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[{y''[x]-7*y'[x]+10*y[x]==0,{y[0]==-1,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(e^{3x} - 2)$$

8.2 problem 2c

Internal problem ID [1088]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 2c.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 2y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = -2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+2*y(x)=0,y(0) = 3, D(y)(0) = -2],y(x), singsol=all)
```

$$y(x) = e^x(-5 \sin(x) + 3 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[{y'[x]-2*y'[x]+2*y[x]==0,{y[0]==3,y'[0]==-2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(3 \cos(x) - 5 \sin(x))$$

8.3 problem 2d

Internal problem ID [1089]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 2d.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + 2y = 0$$

With initial conditions

$$[y(0) = k_0, y'(0) = k_1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+2*y(x)=0,y(0) = k__0, D(y)(0) = k__1],y(x), singsol=all)
```

$$y(x) = e^x((k_1 - k_0) \sin(x) + k_0 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[{y''[x]-2*y'[x]+2*y[x]==0,{y[0]==k0,y'[0]==k1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x((k1 - k0) \sin(x) + k0 \cos(x))$$

8.4 problem 3c

Internal problem ID [1090]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 3c.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + y = 0$$

With initial conditions

$$[y(0) = 7, y'(0) = 4]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+y(x)=0,y(0) = 7, D(y)(0) = 4],y(x), singsol=all)
```

$$y(x) = e^x(7 - 3x)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 14

```
DSolve[{y''[x]-2*y'[x]+y[x]==0,{y[0]==7,y'[0]==4}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(7 - 3x)$$

8.5 problem 3d

Internal problem ID [1091]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 3d.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + y = 0$$

With initial conditions

$$[y(0) = k_0, y'(0) = k_1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+y(x)=0,y(0) = k__0, D(y)(0) = k__1],y(x), singsol=all)
```

$$y(x) = -e^x((x - 1)k_0 - xk_1)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[{y''[x]-2*y'[x]+y[x]==0,{y[0]==k0,y'[0]==k1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^x(k_0(-x) + k_0 + k_1x)$$

8.6 problem 4

Internal problem ID [1092]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 1)y'' + 4y'x + 2y = 0$$

With initial conditions

$$[y(0) = -5, y'(0) = 1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([(x^2-1)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(0) = -5, D(y)(0) = 1],y(x), sings
```

$$y(x) = \frac{-x + 5}{x^2 - 1}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 18

```
DSolve[{(x^2-1)*y'[x]+4*x*y'[x]+2*y[x]==0,{y[0]==-5,y'[0]==1}},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow \frac{5 - x}{x^2 - 1}$$

8.7 problem 10

Internal problem ID [1093]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' - 3y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-3*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + c_2e^{3x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[y''[x]-2*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2e^{4x} + c_1)$$

8.8 problem 11

Internal problem ID [1094]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 6y' + 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{3x}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]-6*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2x + c_1)$$

8.9 problem 12

Internal problem ID [1095]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2ay' + a^2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)-2*a*diff(y(x),x)+a^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{ax}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[y''[x]-2*a*y'[x]+a^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ax}(c_2x + c_1)$$

8.10 problem 13

Internal problem ID [1096]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2y'' + y'x - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1x^2 + c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[x^2*y'[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x} + c_2x$$

8.11 problem 14

Internal problem ID [1097]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$x^2 y'' - y'x + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = x(c_2 \ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 15

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \log(x) + c_1)$$

8.12 problem 15

Internal problem ID [1098]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$x^2 y'' - (2a - 1) x y' + a^2 y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x^2*diff(y(x),x$2)-(2*a-1)*x*diff(y(x),x)+a^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_2 \ln(x) + c_1) x^a$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]-(2*a-1)*x*y'[x]+a^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^a (a c_2 \log(x) + c_1)$$

8.13 problem 16

Internal problem ID [1099]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' - 4y'x + (-16x^2 + 3)y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+(3-16*x^2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{x} (c_1 \sinh(2x) + c_2 \cosh(2x))$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 32

```
DSolve[4*x^2*y'[x]-4*x*y'[x]+(3-16*x^2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-2x}\sqrt{x}(c_2e^{4x} + 4c_1)$$

8.14 problem 17

Internal problem ID [1100]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 1)y'' - y'x + y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve((x-1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x + c_2e^x$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 17

```
DSolve[(x-1)*y'[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x - c_2x$$

8.15 problem 18

Internal problem ID [1101]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x + (x^2 + 2)y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(x^2+2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = x(c_1 \sin(x) + c_2 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 33

```
DSolve[x^2*y'[x]-2*x*y'[x]+(x^2+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-ix} x - \frac{1}{2} i c_2 e^{ix} x$$

8.16 problem 19

Internal problem ID [1102]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2 \sin(x) y'' - 4x(x \cos(x) + \sin(x)) y' + (2x \cos(x) + 3 \sin(x)) y = 0$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 14

```
dsolve(4*x^2*sin(x)*diff(y(x),x$2)-4*x*(x*cos(x)+sin(x))*diff(y(x),x)+(2*x*cos(x)+3*sin(x))*
```

$$y(x) = \sqrt{x} (c_1 + c_2 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.356 (sec). Leaf size: 21

```
DSolve[4*x^2*Sin[x]*y''[x]-4*x*(x*Cos[x]+Sin[x])*y'[x]+(2*x*Cos[x]+3*Sin[x])*y[x]==0,y[x],x,
```

$$y(x) \rightarrow \sqrt{\arccos(\cos(x))} (c_2 \cos(x) + c_1)$$

8.17 problem 20

Internal problem ID [1103]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x - 1)y'' - (3x + 2)y' + (6x - 8)y = 0$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 177

```
dsolve((3*x-1)*diff(y(x),x$2)-(3*x+2)*diff(y(x),x)+(6*x-8)*y(x)=0,y(x), singsol=all)
```

$y(x) =$

$$6615 e^{-\frac{x(i\sqrt{7}-1)}{2}} \left(x - \frac{1}{3}\right)^2 \left(\left(\left(-\frac{23x}{35} + \frac{296}{735}\right) \sqrt{7} + ix - \frac{248i}{105} \right) c_1 \text{KummerM} \left(\frac{1}{2} - \frac{5i\sqrt{7}}{14}, 3, \frac{i\sqrt{7}(3x-1)}{3} \right) - \left(\left(\left(-\frac{23x}{35} + \frac{296}{735}\right) \sqrt{7} + ix - \frac{248i}{105} \right) c_2 \text{KummerM} \left(\frac{1}{2} + \frac{5i\sqrt{7}}{14}, 3, \frac{i\sqrt{7}(3x-1)}{3} \right) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.13 (sec). Leaf size: 109

```
DSolve[(3*x-1)*y''[x]-(3*x+2)*y'[x]+(6*x-8)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 4e^{\frac{1}{6}(1-i\sqrt{7})(3x-1)}(1-3x)^2 \left(c_1 \text{HypergeometricU} \left(\frac{3}{2} - \frac{5i}{2\sqrt{7}}, 3, \frac{1}{3}i\sqrt{7}(3x-1) \right) + c_2 L^2_{-\frac{3}{2} + \frac{5i}{2\sqrt{7}}} \left(\frac{1}{3}i\sqrt{7}(3x-1) \right) \right)$$

8.18 problem 21

Internal problem ID [1104]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 4)y'' + 4y'x + 2y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve((x^2-4)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1x + c_2}{x^2 - 4}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 20

```
DSolve[(x^2-4)*y'[x]+4*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x + c_1}{x^2 - 4}$$

8.19 problem 22

Internal problem ID [1105]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1 + 2x)y'' - 2(2x^2 - 1)y' - 4(x + 1)y = 0$$

✓ Solution by Maple

Time used: 0.187 (sec). Leaf size: 32

```
dsolve((2*x+1)*diff(y(x),x$2)-2*(2*x^2-1)*diff(y(x),x)-4*(x+1)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \operatorname{HeunB}\left(-\frac{1}{2}, -2, -\frac{1}{2}, 3, x + \frac{1}{2}\right) + c_2 \operatorname{HeunB}\left(\frac{1}{2}, -2, -\frac{1}{2}, 3, x + \frac{1}{2}\right) \sqrt{4x + 2}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(2*x+1)*y'[x]-2*(2*x^2-1)*y'[x]-4*(x+1)*y[x]==0,y[x],x,IncludeSingularSolutions -> T
```

Not solved

8.20 problem 23

Internal problem ID [1106]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.1 Homogeneous linear equations. Page 203

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$(x^2 - 2x)y'' + (-x^2 + 2)y' + (2x - 2)y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve((x^2-2*x)*diff(y(x),x$2)+(2-x^2)*diff(y(x),x)+(2*x-2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^2 + c_2e^x$$

✓ Solution by Mathematica

Time used: 0.091 (sec). Leaf size: 18

```
DSolve[(x^2-2*x)*y''[x]+(2-x^2)*y'[x]+(2*x-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^2 + c_1e^x$$

9 Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

9.1	problem 1	274
9.2	problem 2	275
9.3	problem 3	276
9.4	problem 4	277
9.5	problem 5	278
9.6	problem 6	279
9.7	problem 7	280
9.8	problem 8	281
9.9	problem 9	282
9.10	problem 10	283
9.11	problem 11	284
9.12	problem 12	285
9.13	problem 13	286
9.14	problem 14	287
9.15	problem 15	288
9.16	problem 16	289
9.17	problem 17	290
9.18	problem 18	291
9.19	problem 19	292
9.20	problem 20	293
9.21	problem 21	294
9.22	problem 22	295
9.23	problem 23	296
9.24	problem 24	297
9.25	problem 25	298
9.26	problem 26	299
9.27	problem 27	300
9.28	problem 28	301
9.29	problem 29	302
9.30	problem 30	303
9.31	problem 31	304
9.32	problem 32	305
9.33	problem 33	306
9.34	problem 34	307
9.35	problem 35	308
9.36	problem 38 part (a)	309

9.37	problem 38 part (b)	310
9.38	problem 38 part (c)	311
9.39	problem 38 part (d)	312
9.40	problem 38 part (e)	313
9.41	problem 38 part (f)	314
9.42	problem 38 part (g)	315
9.43	problem 39 part(a)	316
9.44	problem 39 part(b)	317
9.45	problem 39 part(c)	318
9.46	problem 39 part(d)	319
9.47	problem 39 part(e)	320
9.48	problem 39 part(f)	321

9.1 problem 1

Internal problem ID [1107]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1 + 2x)y'' - 2y' - (2x + 3)y = (1 + 2x)^2$$

Given that one solution of the ode is

$$y_1 = e^{-x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve([(2*x+1)*diff(y(x),x$2)-2*diff(y(x),x)-(2*x+3)*y(x)=(2*x+1)^2,exp(-x)],singsol=all)
```

$$y(x) = e^{-x}c_2 + x e^x c_1 + 1 - 2x$$

✓ Solution by Mathematica

Time used: 0.126 (sec). Leaf size: 33

```
DSolve[(2*x+1)*y'[x]-2*y'[x]-(2*x+3)*y[x]==(2*x+1)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x-\frac{1}{2}} + x \left(-2 + c_2 e^{x+\frac{1}{2}} \right) + 1$$

9.2 problem 2

Internal problem ID [1108]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + y'x - y = \frac{4}{x^2}$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=4/x^2,x],singsol=all)
```

$$y(x) = c_2 x + \frac{4}{3x^2} + \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 23

```
DSolve[x^2*y''[x]+x*y'[x]-y[x]==4/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{4}{3x^2} + \frac{c_1}{x} + c_2 x$$

9.3 problem 3

Internal problem ID [1109]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - y' x + y = x$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=x,x],singsol=all)
```

$$y(x) = x \left(c_2 + \ln(x) c_1 + \frac{\ln(x)^2}{2} \right)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} x (\log^2(x) + 2c_2 \log(x) + 2c_1)$$

9.4 problem 4

Internal problem ID [1110]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' + 2y = \frac{1}{1 + e^{-x}}$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve([diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=1/(1+exp(-x)),exp(2*x)],singsol=all)
```

$$y(x) = e^x(\ln(1 + e^x)(1 + e^x) + (-1 - e^x)\ln(e^x) + e^x c_1 + c_2 - 1)$$

✓ Solution by Mathematica

Time used: 0.078 (sec). Leaf size: 34

```
DSolve[y''[x]-3*y'[x]+2*y[x]==1/(1+Exp[-x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(2(e^x + 1) \operatorname{arctanh}(2e^x + 1) + c_2 e^x - 1 + c_1)$$

9.5 problem 5

Internal problem ID [1111]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = 7x^{\frac{3}{2}}e^x$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+y(x)=7*x^(3/2)*exp(x),exp(x)],singsol=all)
```

$$y(x) = e^x \left(c_2 + c_1 x + \frac{4x^{\frac{7}{2}}}{5} \right)$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 29

```
DSolve[y''[x]-2*y'[x]+y[x]==7*x^(3/2)*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^x(4x^{7/2} + 5c_2x + 5c_1)$$

9.6 problem 6

Internal problem ID [1112]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' + (-8x^2 + 4x)y' + (4x^2 - 4x - 1)y = 4\sqrt{x}e^x(4x + 1)$$

Given that one solution of the ode is

$$y_1 = \sqrt{x}e^x$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 26

```
dsolve([4*x^2*dif(y(x),x$2)+(4*x-8*x^2)*dif(y(x),x)+(4*x^2-4*x-1)*y(x)=4*x^(1/2)*exp(x)*(1
```

$$y(x) = \frac{(x \ln(x) + 2x^2 + (c_1 - 1)x + c_2)e^x}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 32

```
DSolve[4*x^2*y'[x]+(4*x-8*x^2)*y'[x]+(4*x^2-4*x-1)*y[x]==4*x^(1/2)*Exp[x]*(1+4*x),y[x],x,Int
```

$$y(x) \rightarrow \frac{e^x(2x^2 + x \log(x) + (-1 + c_2)x + c_1)}{\sqrt{x}}$$

9.7 problem 7

Internal problem ID [1113]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + 2y = e^x \sec(x)$$

Given that one solution of the ode is

$$y_1 = \cos(x) e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+2*y(x)=exp(x)*sec(x),exp(x)*cos(x)],singsol=all)
```

$$y(x) = (-\cos(x) \ln(\sec(x)) + \cos(x) c_1 + \sin(x) (c_2 + x)) e^x$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 26

```
DSolve[y''[x]-2*y'[x]+2*y[x]==Exp[x]*Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x((x + c_1) \sin(x) + \cos(x)(\log(\cos(x)) + c_2))$$

9.8 problem 8

Internal problem ID [1114]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y'x + (4x^2 + 2)y = 8e^{-x(2+x)}$$

Given that one solution of the ode is

$$y_1 = e^{-x^2}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve([diff(y(x),x$2)+4*x*diff(y(x),x)+(4*x^2+2)*y(x)=8*exp(-x*(x+2)),exp(-x^2)],singsol=all)
```

$$y(x) = (c_1x + c_2)e^{-x^2} + 2e^{-x(2+x)}$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 29

```
DSolve[y''[x]+4*x*y'[x]+(4*x^2+2)*y[x]==8*Exp[-x*(x+2)],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-x(x+2)}(2 + e^{2x}(c_2x + c_1))$$

9.9 problem 9

Internal problem ID [1115]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x - 4y = -6x - 4$$

Given that one solution of the ode is

$$y_1 = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=-6*x-4,x^2],singsol=all)
```

$$y(x) = \frac{c_2}{x^2} + c_1 x^2 + 2x + 1$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x]==-6*x-4,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 x^2 + \frac{c_1}{x^2} + 2x + 1$$

9.10 problem 10

Internal problem ID [1116]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' + 2x(x-1)y' + (x^2 - 2x + 2)y = e^{2x} x^3$$

Given that one solution of the ode is

$$y_1 = x e^{-x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve([x^2*diff(y(x),x$2)+2*x*(x-1)*diff(y(x),x)+(x^2-2*x+2)*y(x)=x^3*exp(2*x),x*exp(-x)],s
```

$$y(x) = \frac{x(e^{2x} + 9c_1 x e^{-x} + 9e^{-x}c_2)}{9}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 30

```
DSolve[x^2*y''[x]+2*x*(x-1)*y'[x]+(x^2-2*x+2)*y[x]==x^3*Exp[2*x],y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{1}{9} e^{-x} x (e^{3x} + 9(c_2 x + c_1))$$

9.11 problem 11

Internal problem ID [1117]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - x(2x - 1)y' + (x^2 - x - 1)y = x^2 e^x$$

Given that one solution of the ode is

$$y_1 = x e^x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve([x^2*diff(y(x),x$2)-x*(2*x-1)*diff(y(x),x)+(x^2-x-1)*y(x)=x^2*exp(x),x*exp(x)],singular
```

$$y(x) = \frac{e^x(3c_1 x^2 + x^3 + 3c_2)}{3x}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 32

```
DSolve[x^2*y'[x]-x*(2*x-1)*y'[x]+(x^2-x-1)*y[x]==x^2*Exp[x],y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{e^x(2x^3 + 3c_2 x^2 + 6c_1)}{6x}$$

9.12 problem 12

Internal problem ID [1118]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(1 - 2x)y'' + 2y' + (-3 + 2x)y = (4x^2 - 4x + 1)e^x$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve([(1-2*x)*diff(y(x),x$2)+2*diff(y(x),x)+(2*x-3)*y(x)=(1-4*x+4*x^2)*exp(x),exp(x)],sing
```

$$y(x) = c_1 x e^{-x} - \frac{e^x(x^2 - 2c_2 - x)}{2}$$

✓ Solution by Mathematica

Time used: 0.254 (sec). Leaf size: 77

```
DSolve[(1-2*x)*y'[x]+2*y'[x]+(2*x-3)*y[x]==(1-4*x+4*x^2)*Exp[x],y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\frac{1}{2}e^x(x-1)x - \frac{c_2 e^{\frac{1}{2}-x}\sqrt{1-2xx}}{\sqrt{2x-1}} + \frac{c_1 e^{x-\frac{1}{2}}\sqrt{1-2x}}{\sqrt{2x-1}}$$

9.13 problem 13

Internal problem ID [1119]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 3y'x + 4y = 4x^4$$

Given that one solution of the ode is

$$y_1 = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+4*y(x)=4*x^4,x^2],singsol=all)
```

$$y(x) = x^2(\ln(x) c_1 + x^2 + c_2)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 21

```
DSolve[x^2*y'[x]-3*x*y'[x]+4*y[x]==4*x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(x^2 + 2c_2 \log(x) + c_1)$$

9.14 problem 14

Internal problem ID [1120]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$2xy'' + (4x + 1)y' + (1 + 2x)y = 3\sqrt{x}e^{-x}$$

Given that one solution of the ode is

$$y_1 = e^{-x}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([2*x*diff(y(x),x$2)+(4*x+1)*diff(y(x),x)+(2*x+1)*y(x)=3*x^(1/2)*exp(-x),exp(-x)],sing
```

$$y(x) = e^{-x} \left(c_2 + c_1 \sqrt{x} + x^{\frac{3}{2}} \right)$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 28

```
DSolve[2*x*y'[x]+(4*x+1)*y'[x]+(2*x+1)*y[x]==3*x^(1/2)*Exp[-x],y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow e^{-x} \left(x^{3/2} + 2c_2 \sqrt{x} + c_1 \right)$$

9.15 problem 15

Internal problem ID [1121]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' - (1 + 2x)y' + (x + 1)y = -e^{-x}$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 38

```
dsolve([x*diff(y(x),x$2)-(2*x+1)*diff(y(x),x)+(x+1)*y(x)=-exp(-x),exp(x)],singsol=all)
```

$$y(x) = -\exp\text{Integral}_1(2x)x^2e^x + \frac{e^{-x}(2x-1)}{4} + e^x(c_1x^2 + c_2)$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 52

```
DSolve[x*y''[x]-(2*x+1)*y'[x]+(x+1)*y[x]==-Exp[-x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^x x^2 \text{ExpIntegralEi}(-2x) + \frac{1}{4} e^{-x} (2c_2 e^{2x} x^2 + 2x + 4c_1 e^{2x} - 1)$$

9.16 problem 16

Internal problem ID [1122]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' - 4x(x+1)y' + (2x+3)y = 4x^{5/2}e^{2x}$$

Given that one solution of the ode is

$$y_1 = \sqrt{x}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve([4*x^2*diff(y(x),x$2)-4*x*(x+1)*diff(y(x),x)+(2*x+3)*y(x)=4*x^(5/2)*exp(2*x),x^(1/2)]
```

$$y(x) = \left(c_2 + e^x c_1 + \frac{e^{2x}}{2} \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 31

```
DSolve[4*x^2*y''[x]-4*x*(x+1)*y'[x]+(2*x+3)*y[x]==4*x^(5/2)*Exp[2*x],y[x],x,IncludeSingularS
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{x}(e^{2x} + 2c_2e^x + 2c_1)$$

9.17 problem 17

Internal problem ID [1123]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' - 5y'x + 8y = 4x^2$$

Given that one solution of the ode is

$$y_1 = x^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+8*y(x)=4*x^2,x^2],singsol=all)
```

$$y(x) = x^2(c_2x^2 - 2\ln(x) + c_1 - 1)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 23

```
DSolve[x^2*y'[x]-5*x*y'[x]+8*y[x]==4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_2x^2 - 2\log(x) - 1 + c_1)$$

9.18 problem 18

Internal problem ID [1124]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + (-2x + 2)y' + (-2 + x)y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([x*diff(y(x),x$2)+(2-2*x)*diff(y(x),x)+(x-2)*y(x)=0,exp(x)],singsol=all)
```

$$y(x) = \frac{e^x(c_1x + c_2)}{x}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 19

```
DSolve[x*y''[x]+(2-2*x)*y'[x]+(x-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x(c_2x + c_1)}{x}$$

9.19 problem 19

Internal problem ID [1125]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' - 4y'x + 6y = 0$$

Given that one solution of the ode is

$$y_1 = x^2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=0,x^2],singsol=all)
```

$$y(x) = x^2(c_1x + c_2)$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 16

```
DSolve[x^2*y''[x]-4*x*y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_2x + c_1)$$

9.20 problem 20

Internal problem ID [1126]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 \ln(x)^2 y'' - 2x \ln(x) y' + (2 + \ln(x)) y = 0$$

Given that one solution of the ode is

$$y_1 = \ln(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([x^2*(ln(x))^2*diff(y(x),x$2)-2*x*ln(x)*diff(y(x),x)+(2+ln(x))*y(x)=0,ln(x)],singsol=
```

$$y(x) = \ln(x) (c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 15

```
DSolve[x^2*Log[x]^2*y''[x]-2*x*Log[x]*y'[x]+(2+Log[x])*y[x]==0,y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow (c_2 x + c_1) \log(x)$$

9.21 problem 21

Internal problem ID [1127]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$4xy'' + 2y' + y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(\sqrt{x})$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([4*x*diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,sin(sqrt(x))],singsol=all)
```

$$y(x) = c_1 \sin(\sqrt{x}) + c_2 \cos(\sqrt{x})$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 24

```
DSolve[4*x*y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(\sqrt{x}) + c_2 \sin(\sqrt{x})$$

9.22 problem 22

Internal problem ID [1128]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' - (2 + 2x)y' + (2 + x)y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x$2)-(2*x+2)*diff(y(x),x)+(x+2)*y(x)=0,exp(x)],singsol=all)
```

$$y(x) = e^x(c_2x^3 + c_1)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 23

```
DSolve[x*y''[x]-(2*x+2)*y'[x]+(x+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^x(c_2x^3 + 3c_1)$$

9.23 problem 23

Internal problem ID [1129]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' - (2a - 1)xy' + a^2 y = 0$$

Given that one solution of the ode is

$$y_1 = x^a$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([x^2*diff(y(x),x$2)-(2*a-1)*x*diff(y(x),x)+a^2*y(x)=0,x^a],singsol=all)
```

$$y(x) = (c_2 \ln(x) + c_1) x^a$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]-(2*a-1)*x*y'[x]+a^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^a (ac_2 \log(x) + c_1)$$

9.24 problem 24

Internal problem ID [1130]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 2y'x + (x^2 + 2)y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x) x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(x^2+2)*y(x)=0,x*sin(x)],singsol=all)
```

$$y(x) = x(c_1 \sin(x) + c_2 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 33

```
DSolve[x^2*y'[x]-2*x*y'[x]+(x^2+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-ix} x - \frac{1}{2} i c_2 e^{ix} x$$

9.25 problem 25

Internal problem ID [1131]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' - (4x + 1)y' + (4x + 2)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)-(4*x+1)*diff(y(x),x)+(4*x+2)*y(x)=0,exp(2*x)],singsol=all)
```

$$y(x) = e^{2x}(c_2x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 25

```
DSolve[x*y''[x]-(4*x+1)*y'[x]+(4*x+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{2x}(c_2x^2 + 2c_1)$$

9.26 problem 26

Internal problem ID [1132]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2 \sin(x) y'' - 4x(x \cos(x) + \sin(x)) y' + (2x \cos(x) + 3 \sin(x)) y = 0$$

Given that one solution of the ode is

$$y_1 = \sqrt{x}$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 14

```
dsolve([4*x^2*sin(x)*diff(y(x),x$2)-4*x*(x*cos(x)+sin(x))*diff(y(x),x)+(2*x*cos(x)+3*sin(x))*y(x)=0,y(x),x)
```

$$y(x) = \sqrt{x} (c_1 + c_2 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.322 (sec). Leaf size: 21

```
DSolve[4*x^2*Sin[x]*y''[x]-4*x*(x*Cos[x]+Sin[x])*y'[x]+(2*x*Cos[x]+3*Sin[x])*y[x]==0,y[x],x,
```

$$y(x) \rightarrow \sqrt{\arccos(\cos(x))} (c_2 \cos(x) + c_1)$$

9.27 problem 27

Internal problem ID [1133]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' - 4y'x + (-16x^2 + 3)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}\sqrt{x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([4*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+(3-16*x^2)*y(x)=0,sqrt(x)*exp(2*x)],singsol=all)
```

$$y(x) = \sqrt{x}(c_1 \sinh(2x) + c_2 \cosh(2x))$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 32

```
DSolve[4*x^2*y''[x]-4*x*y'[x]+(3-16*x^2)*y[x]==0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-2x}\sqrt{x}(c_2e^{4x} + 4c_1)$$

9.28 problem 28

Internal problem ID [1134]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1 + 2x)xy'' - 2(2x^2 - 1)y' - 4(x + 1)y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{1}{x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([(2*x+1)*x*diff(y(x),x$2)-2*(2*x^2-1)*diff(y(x),x)-4*(x+1)*y(x)=0,1/x],singsol=all)
```

$$y(x) = \frac{c_2 e^{2x} x + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 28

```
DSolve[(2*x+1)*x*y''[x]-2*(2*x^2-1)*y'[x]-4*(x+1)*y[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{c_2 e^{2x+1} x + c_1}{\sqrt{e} x}$$

9.29 problem 29

Internal problem ID [1135]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 2x)y'' + (-x^2 + 2)y' + (2x - 2)y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([(x^2-2*x)*diff(y(x),x$2)+(2-x^2)*diff(y(x),x)+(2*x-2)*y(x)=0,exp(x)],singsol=all)
```

$$y(x) = c_1x^2 + c_2e^x$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 18

```
DSolve[(x^2-2*x)*y''[x]+(2-x^2)*y'[x]+(2*x-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^2 + c_1e^x$$

9.30 problem 30

Internal problem ID [1136]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' - (4x + 1)y' + (4x + 2)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([x*diff(y(x),x$2)-(4*x+1)*diff(y(x),x)+(4*x+2)*y(x)=0,exp(2*x)],singsol=all)
```

$$y(x) = e^{2x}(c_2x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 25

```
DSolve[x*y''[x]-(4*x+1)*y'[x]+(4*x+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{2x}(c_2x^2 + 2c_1)$$

9.31 problem 31

Internal problem ID [1137]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 3y'x + 4y = 4x^4$$

Given that one solution of the ode is

$$y_1 = x^2$$

With initial conditions

$$[y(-1) = 7, y'(-1) = -8]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 21

```
dsolve([x^2*diff(diff(y(x),x),x)-3*x*diff(y(x),x)+4*y(x) = 4*x^4, x^2, y(-1) = 7, D(y)(-1) =
```

$$y(x) = (8i\pi + x^2 - 8 \ln(x) + 6) x^2$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 37

```
DSolve[x^2*y''[x]-3*x*y'[x]+4*y[x]==4*x^2,{y[-1]==7,y'[-1]==8},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow x^2(2 \log^2(x) + (-22 - 4i\pi) \log(x) - 2\pi^2 + 22i\pi + 7)$$

9.32 problem 32

Internal problem ID [1138]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x - 1)y'' - (3x + 2)y' - (6x - 8)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

With initial conditions

$$[y(0) = 2, y'(0) = 3]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 18

```
dsolve([(3*x-1)*diff(diff(y(x),x),x)-(3*x+2)*diff(y(x),x)-(6*x-8)*y(x) = 0, exp(2*x), y(0) =
```

$$y(x) = 2e^{2x} - xe^{-x}$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 21

```
DSolve[(3*x-1)*y'[x]-(3*x+2)*y'[x]-(6*x-8)*y[x]==0,{y[0]==2,y'[0]==3},y[x],x,IncludeSingular
```

$$y(x) \rightarrow 2e^{2x} - e^{-x}x$$

9.33 problem 33

Internal problem ID [1139]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction of order. Page 253

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(x + 1)^2 y'' - 2(x + 1) y' - (x^2 + 2x - 1) y = (x + 1)^3 e^x$$

Given that one solution of the ode is

$$y_1 = (x + 1) e^x$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 22

```
dsolve([(1+x)^2*diff(diff(y(x),x),x)-2*(1+x)*diff(y(x),x)-(x^2+2*x-1)*y(x) = (1+x)^3*exp(x),
```

$$y(x) = \frac{(x + 1)(x e^x - 5 \sinh(x) + 2 \cosh(x))}{2}$$

✓ Solution by Mathematica

Time used: 21.262 (sec). Leaf size: 5749

```
DSolve[(x+1)^2*y'[x]-2*(x+1)*x*y'[x]-(x^2+2*x-1)*y[x]==(x+1)^3*Exp[x],{y[0]==1,y'[0]==-1},y
```

Too large to display

9.34 problem 34

Internal problem ID [1140]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 2y'x - 2y = x^2$$

Given that one solution of the ode is

$$y_1 = x$$

With initial conditions

$$\left[y(1) = \frac{5}{4}, y'(1) = \frac{3}{2} \right]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 11

```
dsolve([x^2*diff(diff(y(x),x),x)+2*x*diff(y(x),x)-2*y(x) = x^2, x, y(1) = 5/4, D(y)(1) = 3/2]
```

$$y(x) = x + \frac{1}{4}x^2$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 13

```
DSolve[x^2*y''[x]+2*x*y'[x]-2*y[x]==x^2,{y[1]==5/4,y'[1]==3/2},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{1}{4}x(x + 4)$$

9.35 problem 35

Internal problem ID [1141]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$(x^2 - 4)y'' + 4y'x + 2y = 2 + x$$

Given that one solution of the ode is

$$y_1 = \frac{1}{-2 + x}$$

With initial conditions

$$\left[y(0) = -\frac{1}{3}, y'(0) = -1 \right]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve([(x^2-4)*diff(diff(y(x),x),x)+4*x*diff(y(x),x)+2*y(x) = 2+x, 1/(-2+x), y(0) = -1/3, D
```

$$y(x) = \frac{x^3 + 6x^2 + 24x + 8}{6x^2 - 24}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 30

```
DSolve[(x^2-4)*y''[x]+4*x*y'[x]+2*y[x]==x+2,{y[1]==5/4,y'[1]==3/2},y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow \frac{-2x^3 - 12x^2 + 54x + 5}{48 - 12x^2}$$

9.36 problem 38 part (a)

Internal problem ID [1142]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + y^2 = -k^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)+y(x)^2+k^2=0,y(x), singsol=all)
```

$$y(x) = -\tan(k(c_1 + x))k$$

✓ Solution by Mathematica

Time used: 4.112 (sec). Leaf size: 35

```
DSolve[y'[x]+y[x]^2+k^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -k \tan(k(x - c_1))$$

$$y(x) \rightarrow -ik$$

$$y(x) \rightarrow ik$$

9.37 problem 38 part (b)

Internal problem ID [1143]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + y^2 - 3y = -2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)+y(x)^2-3*y(x)+2=0,y(x), singsol=all)
```

$$y(x) = \frac{2e^x c_1 - 1}{e^x c_1 - 1}$$

✓ Solution by Mathematica

Time used: 0.93 (sec). Leaf size: 40

```
DSolve[y' [x]+y [x]^2-3*y [x]+2==0,y [x] ,x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow \frac{2e^x - e^{c_1}}{e^x - e^{c_1}} \\y(x) &\rightarrow 1 \\y(x) &\rightarrow 2\end{aligned}$$

9.38 problem 38 part (c)

Internal problem ID [1144]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + y^2 + 5y = 6$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)+y(x)^2+5*y(x)-6=0,y(x), singsol=all)
```

$$y(x) = \frac{6 + c_1 e^{7x}}{c_1 e^{7x} - 1}$$

✓ Solution by Mathematica

Time used: 0.71 (sec). Leaf size: 46

```
DSolve[y'[x]+y[x]^2+5*y[x]-6==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow \frac{e^{7x} + 6e^{7c_1}}{e^{7x} - e^{7c_1}} \\y(x) &\rightarrow -6 \\y(x) &\rightarrow 1\end{aligned}$$

9.39 problem 38 part (d)

Internal problem ID [1145]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$y' + y^2 + 8y = -7$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)+y(x)^2+8*y(x)+7=0,y(x), singsol=all)
```

$$y(x) = \frac{7 - c_1 e^{6x}}{c_1 e^{6x} - 1}$$

✓ Solution by Mathematica

Time used: 0.655 (sec). Leaf size: 47

```
DSolve[y'[x]+y[x]^2+8*y[x]+7==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow -\frac{e^{6x} - 7e^{6c_1}}{e^{6x} - e^{6c_1}} \\y(x) &\rightarrow -7 \\y(x) &\rightarrow -1\end{aligned}$$

9.40 problem 38 part (e)

Internal problem ID [1146]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' + y^2 + 14y = -50$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+y(x)^2+14*y(x)+50=0,y(x), singsol=all)
```

$$y(x) = -7 - \tan(c_1 + x)$$

✓ Solution by Mathematica

Time used: 0.553 (sec). Leaf size: 30

```
DSolve[y'[x]+y[x]^2+14*y[x]+50==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -7 - \tan(x - c_1)$$

$$y(x) \rightarrow -7 - i$$

$$y(x) \rightarrow -7 + i$$

9.41 problem 38 part (f)

Internal problem ID [1147]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (f).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$6y' + 6y^2 - y = 1$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 24

```
dsolve(6*diff(y(x),x)+6*y(x)^2-y(x)-1=0,y(x), singsol=all)
```

$$y(x) = \frac{1 + c_1 e^{\frac{5x}{6}}}{2c_1 e^{\frac{5x}{6}} - 3}$$

✓ Solution by Mathematica

Time used: 0.234 (sec). Leaf size: 56

```
DSolve[6*y'[x]+6*y[x]^2-y[x]-1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{5x/6} - e^{5c_1}}{2e^{5x/6} + 3e^{5c_1}}$$

$$y(x) \rightarrow -\frac{1}{3}$$

$$y(x) \rightarrow \frac{1}{2}$$

9.42 problem 38 part (g)

Internal problem ID [1148]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 38 part (g).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [quadrature]

$$36y' + 36y^2 - 12y = -1$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(36*diff(y(x),x)+36*y(x)^2-12*y(x)+1=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 + x + 6}{6x + 6c_1}$$

✓ Solution by Mathematica

Time used: 0.119 (sec). Leaf size: 30

```
DSolve[36*y'[x]+36*y[x]^2-12*y[x]+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x + 6 - 36c_1}{6x - 216c_1}$$

$$y(x) \rightarrow \frac{1}{6}$$

9.43 problem 39 part(a)

Internal problem ID [1149]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x),G(x)]’], _Riccati]`

$$x^2(y' + y^2) - x(2 + x)y = -x - 2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(x^2*(diff(y(x),x)+y(x)^2)-x*(x+2)*y(x)+x+2=0,y(x), singsol=all)
```

$$y(x) = \frac{x e^x + e^x - c_1}{(-c_1 + e^x) x}$$

✓ Solution by Mathematica

Time used: 0.179 (sec). Leaf size: 49

```
DSolve[x^2*(y'[x]+y[x]) - x*(x+2)*y[x] + x + 2 == 0, y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{\frac{1}{x}-x} \left(\int_1^x \frac{e^{K[1]-\frac{1}{K[1]}} (K[1]^2 + K[1] - 2)}{K[1]^2} dK[1] + c_1 \right)$$

9.44 problem 39 part(b)

Internal problem ID [1150]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(b).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[[_homogeneous, 'class C'], _Riccati]`

$$y' + y^2 + 4yx = -4x^2 - 2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)+y(x)^2+4*x*y(x)+4*x^2+2=0,y(x), singsol=all)
```

$$y(x) = \frac{-2c_1x + 2x^2 - 1}{c_1 - x}$$

✓ Solution by Mathematica

Time used: 0.129 (sec). Leaf size: 22

```
DSolve[y'[x]+y[x]^2+4*x*y[x]+4*x^2+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x + \frac{1}{x + c_1}$$
$$y(x) \rightarrow -2x$$

9.45 problem 39 part(c)

Internal problem ID [1151]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(c).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, _Riccati]`

$$(1 + 2x)(y' + y^2) - 2y = 2x + 3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve((2*x+1)*(diff(y(x),x)+y(x)^2)-2*y(x)-(2*x+3)=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{2x}x + e^{2x} - c_1}{e^{2x}x + c_1}$$

✓ Solution by Mathematica

Time used: 0.403 (sec). Leaf size: 41

```
DSolve[(2*x+1)*(y'[x]+y[x]^2)-2*y[x]-(2*x+3)==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{2x+1}(x+1) - c_1}{e^{2x+1}x + c_1}$$
$$y(x) \rightarrow -1$$

9.46 problem 39 part(d)

Internal problem ID [1152]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(d).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, _Riccati]`

$$(3x - 1)(y' + y^2) - y(3x + 2) = 6x - 8$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 31

```
dsolve((3*x-1)*(diff(y(x),x)+y(x)^2)-(3*x+2)*y(x)-6*x+8=0,y(x), singsol=all)
```

$$y(x) = \frac{-c_1x + 2e^{3x-1} + c_1}{c_1x + e^{3x-1}}$$

✓ Solution by Mathematica

Time used: 0.557 (sec). Leaf size: 41

```
DSolve[(3*x-1)*(y'[x]+y[x]^2)-(3*x+2)*y[x]-6*x+8==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2(-ex + c_1e^{3x} + e)}{2ex + c_1e^{3x}}$$
$$y(x) \rightarrow 2$$

9.47 problem 39 part(e)

Internal problem ID [1153]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(e).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_rational, [_1st_order, ‘_with_symmetry_[F(x),G(x)]’], _Riccati]`

$$x^2(y' + y^2) + yx = -x^2 + \frac{1}{4}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 46

```
dsolve(x^2*(diff(y(x),x)+y(x)^2)+x*y(x)+x^2-1/4=0,y(x), singsol=all)
```

$$y(x) = \frac{-4c_1x - e^{-2ix} - 2ie^{-2ix}x - 2ic_1}{2x(e^{-2ix} + 2ic_1)}$$

✓ Solution by Mathematica

Time used: 0.377 (sec). Leaf size: 22

```
DSolve[x^2*(y'[x]+y[x]^2)+x*y[x]+x^2-1/4==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2x} - \tan(x - c_1)$$

9.48 problem 39 part(f)

Internal problem ID [1154]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.6 Reduction or order. Page 253

Problem number: 39 part(f).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_homogeneous, 'class G', _rational, _Riccati]`

$$x^2(y' + y^2) - 7yx = -7$$

✓ Solution by Maple

Time used: 0.375 (sec). Leaf size: 24

```
dsolve(x^2*(diff(y(x),x)+y(x)^2)-7*x*y(x)+7=0,y(x), singsol=all)
```

$$y(x) = \frac{-7x^6 + c_1}{x(-x^6 + c_1)}$$

✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 34

```
DSolve[x^2*(y'[x]+y[x]^2)-7*x*y[x]+7==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{7x^6 - 6c_1}{x^7 - 6c_1x}$$
$$y(x) \rightarrow \frac{1}{x}$$

10 Chapter 5 linear second order equations.

Section 5.7 Variation of Parameters. Page 262

10.1 problem 1	323
10.2 problem 2	324
10.3 problem 3	325
10.4 problem 4	326
10.5 problem 5	327
10.6 problem 6	328
10.7 problem 7	329
10.8 problem 8	330
10.9 problem 9	331
10.10problem 10	332
10.11problem 11	333
10.12problem 12	334
10.13problem 13	335
10.14problem 14	336
10.15problem 15	337
10.16problem 16	338
10.17problem 17	339
10.18problem 18	340
10.19problem 19	341
10.20problem 20	343
10.21problem 21	344
10.22problem 22	345
10.23problem 23	346
10.24problem 24	347
10.25problem 25	348
10.26problem 26	349
10.27problem 27	350
10.28problem 28	351
10.29problem 29	352
10.30problem 30	353
10.31problem 31	354
10.32problem 32	355
10.33problem 33	356
10.34problem 34	357
10.35problem 35	358

10.1 problem 1

Internal problem ID [1155]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 9y = \tan(3x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+9*y(x)=tan(3*x),y(x), singsol=all)
```

$$y(x) = \sin(3x) c_2 + \cos(3x) c_1 - \frac{\cos(3x) \ln(\sec(3x) + \tan(3x))}{9}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==Tan[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{9} \cos(3x) \operatorname{arctanh}(\sin(3x)) + c_1 \cos(3x) + c_2 \sin(3x)$$

10.2 problem 2

Internal problem ID [1156]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y = \sin(2x) \sec(2x)^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$2)+4*y(x)=sin(2*x)*sec(2*x)^2,y(x), singsol=all)
```

$$y(x) = \frac{\ln(\sec(2x)) \sin(2x)}{4} + \frac{(4c_2 - 1) \sin(2x)}{4} + \frac{\cos(2x)(2c_1 + x)}{2}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 33

```
DSolve[y''[x]+4*y[x]==Sin[2*x]*Sec[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (-x + c_1) \cos(2x) + \sin(x) \cos(x)(2 \log(\cos(x)) - 1 + 2c_2)$$

10.3 problem 3

Internal problem ID [1157]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 3y' + 2y = \frac{4}{1 + e^{-x}}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=4/(1+exp(-x)),y(x), singsol=all)
```

$$y(x) = e^x((4e^x + 4) \ln(1 + e^x) + (-4e^x - 4) \ln(e^x) + e^x c_1 + c_2 - 4)$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 34

```
DSolve[y''[x]-3*y'[x]+2*y[x]==4/(1+Exp[-x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(8(e^x + 1) \operatorname{arctanh}(2e^x + 1) + c_2 e^x - 4 + c_1)$$

10.4 problem 4

Internal problem ID [1158]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + 2y = 3e^x \sec(x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+2*y(x)=3*exp(x)*sec(x),y(x), singsol=all)
```

$$y(x) = e^x(\sin(x)c_2 + \cos(x)c_1 + 3x \sin(x) - 3 \cos(x) \ln(\sec(x)))$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 30

```
DSolve[y''[x]-2*y'[x]+2*y[x]==3*Exp[x]*Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x((3x + c_1) \sin(x) + \cos(x)(3 \log(\cos(x)) + c_2))$$

10.5 problem 5

Internal problem ID [1159]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - 2y' + y = 14x^{\frac{3}{2}}e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=14*x^(3/2)*exp(x),y(x), singsol=all)
```

$$y(x) = e^x \left(c_2 + c_1 x + \frac{8x^{\frac{7}{2}}}{5} \right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 29

```
DSolve[y''[x]-2*y'[x]+y[x]==14*x^(3/2)*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^x(8x^{7/2} + 5c_2x + 5c_1)$$

10.6 problem 6

Internal problem ID [1160]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' - y = \frac{4e^{-x}}{1 - e^{-2x}}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve(diff(y(x),x$2)-y(x)=4*exp(-x)/(1-exp(-2*x)),y(x), singsol=all)
```

$$y(x) = \ln(1 - e^{-2x})e^x + e^{-x} \ln(e^{-2x}) - e^{-x} \ln(-1 + e^{-2x}) + c_2 e^x + e^{-x} c_1$$

✓ Solution by Mathematica

Time used: 0.121 (sec). Leaf size: 67

```
DSolve[y''[x]-y[x]==4*Exp[-x]/(1-Exp[-2*x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(-2e^{2x} \log(e^x) + (e^{2x} - 1) \log(e^x - 1) + e^{2x} \log(e^x + 1) - \log(e^x + 1) + c_1 e^{2x} + c_2)$$

10.7 problem 7

Internal problem ID [1161]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2 y'' + y'x - y = 2x^2 + 2$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=2*x^2+2,y(x), singsol=all)
```

$$y(x) = c_2 x + \frac{2x^2}{3} + \frac{c_1}{x} - 2$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 24

```
DSolve[x^2*y'[x]+x*y'[x]-y[x]==2*x^2+2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x^2}{3} + c_2 x + \frac{c_1}{x} - 2$$

10.8 problem 8

Internal problem ID [1162]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _linear, _nonhomogeneous]`

$$x^2 y'' + (-2x + 2) y' + (-2 + x) y = e^{2x}$$

X Solution by Maple

```
dsolve(x^2*diff(y(x),x$2)+(2-2*x)*diff(y(x),x)+(x-2)*y(x)=exp(2*x),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[x^2*y'[x]+(2-2*x)*y'[x]+(x-2)*y[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True
```

Not solved

10.9 problem 9

Internal problem ID [1163]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' + (-8x^2 + 4x)y' + (4x^2 - 4x - 1)y = 4\sqrt{x}e^x$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(y(x),x$2)+(4*x-8*x^2)*diff(y(x),x)+(4*x^2-4*x-1)*y(x)=4*x^(1/2)*exp(x),y(x)
```

$$y(x) = \frac{e^x(x \ln(x) + (c_1 - 1)x + c_2)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 27

```
DSolve[4*x^2*y''[x]+(4*x-8*x^2)*y'[x]+(4*x^2-4*x-1)*y[x]==4*x^(1/2)*Exp[x],y[x],x,IncludeSin
```

$$y(x) \rightarrow \frac{e^x(x \log(x) + (-1 + c_2)x + c_1)}{\sqrt{x}}$$

10.10 problem 10

Internal problem ID [1164]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$y'' + 4y'x + (4x^2 + 2)y = 4e^{-x(2+x)}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+4*x*diff(y(x),x)+(4*x^2+2)*y(x)=4*exp(-x*(x+2)),y(x), singsol=all)
```

$$y(x) = (c_1x + c_2)e^{-x^2} + e^{-x(2+x)}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 27

```
DSolve[4*x^2*y''[x]+(4*x-8*x^2)*y'[x]+(4*x^2-4*x-1)*y[x]==4*x^(1/2)*Exp[x],y[x],x,IncludeSin
```

$$y(x) \rightarrow \frac{e^x(x \log(x) + (-1 + c_2)x + c_1)}{\sqrt{x}}$$

10.11 problem 11

Internal problem ID [1165]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 4y'x + 6y = x^{\frac{5}{2}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=x^(5/2),y(x), singsol=all)
```

$$y(x) = c_2 x^3 + c_1 x^2 - 4x^{\frac{5}{2}}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 23

```
DSolve[x^2*y'[x]-4*x*y'[x]+6*y[x]==x^(5/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(-4\sqrt{x} + c_2 x + c_1)$$

10.12 problem 12

Internal problem ID [1166]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$x^2 y'' - 3y'x + 3y = 2 \sin(x) x^4$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+3*y(x)=2*x^4*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{(c_1 x^2 - 4x \sin(x) - 4 \cos(x) + 2c_2) x}{2}$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]-3*x*y'[x]+3*y[x]==2*x^4*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 x^2 - 2x \sin(x) - 2 \cos(x) + c_1)$$

10.13 problem 13

Internal problem ID [1167]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(1 + 2x)y'' - 2y' - (2x + 3)y = (1 + 2x)^2 e^{-x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve((2*x+1)*diff(y(x),x$2)-2*diff(y(x),x)-(2*x+3)*y(x)=(2*x+1)^2*exp(-x),y(x), singsol=all)
```

$$y(x) = \frac{(-x^2 + 2c_2 - x)e^{-x}}{2} + x e^x c_1$$

✓ Solution by Mathematica

Time used: 0.084 (sec). Leaf size: 42

```
DSolve[(2*x+1)*y'[x]-2*y'[x]-(2*x+3)*y[x]==(2*x+1)^2*Exp[-x],y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow -\frac{1}{2}e^{-x}x(x+1) + c_1e^{-x-\frac{1}{2}} + c_2e^{x+\frac{1}{2}x}$$

10.14 problem 14

Internal problem ID [1168]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$2xy'' + 2y' + 2y = \sin(\sqrt{x})$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 71

```
dsolve(2*x*diff(y(x),x$2)+2*diff(y(x),x)+2*y(x)=sin(sqrt(x)),y(x), singsol=all)
```

$$y(x) = \text{BesselJ}(0, 2\sqrt{x}) c_2 + \text{BesselY}(0, 2\sqrt{x}) c_1 + \frac{\pi \left(\int \text{BesselJ}(0, 2\sqrt{x}) \sin(\sqrt{x}) dx \right) \text{BesselY}(0, 2\sqrt{x})}{2} - \frac{\pi \left(\int \text{BesselY}(0, 2\sqrt{x}) \sin(\sqrt{x}) dx \right) \text{BesselJ}(0, 2\sqrt{x})}{2}$$

✓ Solution by Mathematica

Time used: 11.396 (sec). Leaf size: 110

```
DSolve[2*x*y'[x]+2*y'[x]+2*y[x]==Sin[Sqrt[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{BesselJ}(0, 2\sqrt{x}) \int_1^x -\frac{1}{2} \pi \text{BesselY}(0, 2\sqrt{K[1]}) \sin(\sqrt{K[1]}) dK[1] + 2 \text{BesselY}(0, 2\sqrt{x}) \int_1^x \frac{1}{4} \pi \text{BesselJ}(0, 2\sqrt{K[2]}) \sin(\sqrt{K[2]}) dK[2] + c_1 \text{BesselJ}(0, 2\sqrt{x}) + 2c_2 \text{BesselY}(0, 2\sqrt{x})$$

10.15 problem 15

Internal problem ID [1169]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' - (2 + 2x)y' + (2 + x)y = 6e^x x^3$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(x*diff(y(x),x$2)-(2*x+2)*diff(y(x),x)+(x+2)*y(x)=6*x^3*exp(x),y(x), singsol=all)
```

$$y(x) = e^x \left(c_2 + c_1 x^3 + \frac{3}{2} x^4 \right)$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 29

```
DSolve[x*y''[x]-(2*x+2)*y'[x]+(x+2)*y[x]==6*x^3*Exp[x],y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow \frac{1}{6} e^x (9x^4 + 2c_2 x^3 + 6c_1)$$

10.16 problem 16

Internal problem ID [1170]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - (2a - 1) x y' + a^2 y = x^{1+a}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)-(2*a-1)*x*diff(y(x),x)+a^2*y(x)=x^(a+1),y(x), singsol=all)
```

$$y(x) = x^a(c_2 + \ln(x) c_1 + x)$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 19

```
DSolve[x^2*y'[x]-(2*a-1)*x*y'[x]+a^2*y[x]==x^(a+1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^a(ac_2 \log(x) + x + c_1)$$

10.17 problem 17

Internal problem ID [1171]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2y'x + (x^2 + 2)y = x^3 \cos(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+(x^2+2)*y(x)=x^3*cos(x),y(x), singsol=all)
```

$$y(x) = \frac{((x + 2c_2) \sin(x) + 2 \cos(x) c_1) x}{2}$$

✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 49

```
DSolve[x^2*y''[x]-2*x*y'[x]+(x^2+2)*y[x]==x^3*Cos[x],y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{8} e^{-ix} x (2ix + e^{2ix} (-2ix + 1 - 4ic_2) + 1 + 8c_1)$$

10.18 problem 18

Internal problem ID [1172]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$xy'' - y' - 4yx^3 = 8x^5$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(x*diff(y(x),x$2)-diff(y(x),x)-4*x^3*y(x)=8*x^5,y(x), singsol=all)
```

$$y(x) = \sinh(x^2) c_2 + \cosh(x^2) c_1 - 2x^2$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 28

```
DSolve[x*y''[x]-y'[x]-4*x^3*y[x]==8*x^5,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x^2 + c_1 \cosh(x^2) + ic_2 \sinh(x^2)$$

10.19 problem 19

Internal problem ID [1173]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$\sin(x) y'' + (2 \sin(x) - \cos(x)) y' + (-\cos(x) + \sin(x)) y = e^{-x}$$

✓ Solution by Maple

Time used: 0.093 (sec). Leaf size: 58

```
dsolve(sin(x)*diff(y(x),x$2)+(2*sin(x)-cos(x))*diff(y(x),x)+(sin(x)-cos(x))*y(x)=exp(-x),y(x)
```

$$y(x) = -e^{\arcsin(\cos(x))} \left(\left(\int \csc(x)^2 e^{-\arcsin(\cos(x))-x} dx \right) \cos(x) - \cos(x) c_1 \right. \\ \left. - \left(\int \cot(x) \csc(x) e^{-\arcsin(\cos(x))-x} dx \right) - c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.809 (sec). Leaf size: 121

```
DSolve[Sin[x]*y''[x]+(2*Sin[x]-Cos[x])*y'[x]+(Sin[x]-Cos[x])*y[x]==Exp[-x],y[x],x,IncludeSins]
```

$$\begin{aligned} y(x) \rightarrow & -\sqrt{\sin^2(x)} \exp\left(-\arccos(\cos(x)) - 4 \arctan\left(\frac{\sqrt{\sin^2(x)}}{\cos(x)+1}\right)\right. \\ & \left. + 4 \cot^{-1}\left(\frac{\cos(x)+1}{\sqrt{\sin^2(x)}}\right)\right) \\ & + c_2 \cos(x) \exp\left(2\left(\cot^{-1}\left(\frac{\cos(x)+1}{\sqrt{\sin^2(x)}}\right) - 2 \arctan\left(\frac{\sqrt{\sin^2(x)}}{\cos(x)+1}\right)\right)\right) \\ & + c_1 e^{-2 \cot^{-1}\left(\frac{\cos(x)+1}{\sqrt{\sin^2(x)}}\right)} \end{aligned}$$

10.20 problem 20

Internal problem ID [1174]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' - 4y'x + (-16x^2 + 3)y = 8x^{\frac{5}{2}}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(4*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+(3-16*x^2)*y(x)=8*x^(5/2),y(x), singsol=all)
```

$$y(x) = \sqrt{x} \left(-\frac{1}{2} + \sinh(2x)c_2 + \cosh(2x)c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 39

```
DSolve[4*x^2*y''[x]-4*x*y'[x]+(3-16*x^2)*y[x]==8*x^(5/2),y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{4}e^{-2x}\sqrt{x}(-2e^{2x} + c_2e^{4x} + 4c_1)$$

10.21 problem 21

Internal problem ID [1175]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' - 4y'x + (4x^2 + 3)y = x^{\frac{7}{2}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+(4*x^2+3)*y(x)=x^(7/2),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x}(x + 4 \sin(x) c_2 + 4 \cos(x) c_1)}{4}$$

✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 48

```
DSolve[4*x^2*y''[x]-4*x*y'[x]+(4*x^2+3)*y[x]==x^(7/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-ix}\sqrt{x}(e^{ix}x - 2ic_2e^{2ix} + 4c_1)$$

10.22 problem 22

Internal problem ID [1176]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2y'x - (x^2 - 2)y = 3x^4$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)-(x^2-2)*y(x)=3*x^4,y(x), singsol=all)
```

$$y(x) = \sinh(x) x c_2 + \cosh(x) x c_1 - 3x^2$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 29

```
DSolve[x^2*y'[x]-2*x*y'[x]-(x^2-2)*y[x]==3*x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x(-6x + 2c_1e^{-x} + c_2e^x)$$

10.23 problem 23

Internal problem ID [1177]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2x(x+1)y' + (x^2 + 2x + 2)y = e^x x^3$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)-2*x*(x+1)*diff(y(x),x)+(x^2+2*x+2)*y(x)=x^3*exp(x),y(x), singsol=a
```

$$y(x) = \frac{e^x x(2c_1 x + x^2 + 2c_2)}{2}$$

✓ Solution by Mathematica

Time used: 0.867 (sec). Leaf size: 227

```
DSolve[x^2*y''[x]-2*x*y'[x]+(x^2+2*x+2)*y[x]==x^3*Exp[x],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow e^{ix} x \left(\text{HypergeometricU}(-i, 0, -2ix) \int_1^x \frac{ie^{(1-i)K[1]} L_i^{-1}(-2iK[1])}{4 \text{Hypergeometric1F1}(1-i, 2, -2iK[1]) \text{HypergeometricU}(1-i, 1, -2iK[1]) K[1]} - 2 \text{Hypergeomet}$$

$$+ L_i^{-1}(-2ix) \int_1^x \frac{ie^{(1-i)K[2]} \text{HypergeometricU}(-i, 1, -2ix)}{4 \text{Hypergeometric1F1}(1-i, 2, -2iK[2]) \text{HypergeometricU}(1-i, 1, -2iK[2]) K[2]} - 2 \text{Hypergeomet}$$

$$+ c_1 \text{HypergeometricU}(-i, 0, -2ix) + c_2 L_i^{-1}(-2ix) \right)$$

10.24 problem 24

Internal problem ID [1178]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$x^2y'' - y'x - 3y = x^{\frac{3}{2}}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)-3*y(x)=x^(3/2),y(x), singsol=all)
```

$$y(x) = \frac{15c_2x^4 - 4x^{\frac{5}{2}} + 15c_1}{15x}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 27

```
DSolve[x^2*y'[x]-x*y'[x]-3*y[x]==x^(3/2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4x^{3/2}}{15} + c_2x^3 + \frac{c_1}{x}$$

10.25 problem 25

Internal problem ID [1179]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - x(x+4)y' + 2y(x+3) = e^x x^4$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-x*(x+4)*diff(y(x),x)+2*(x+3)*y(x)=x^4*exp(x),y(x), singsol=all)
```

$$y(x) = ((c_2 + x) e^x + c_1) x^2$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 21

```
DSolve[x^2*y''[x]-x*(x+4)*y'[x]+2*(x+3)*y[x]==x^4*Exp[x],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow x^2(e^x(x-1+c_2) + c_1)$$

10.26 problem 26

Internal problem ID [1180]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 2x(2+x)y' + (x^2 + 4x + 6)y = 2x e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-2*x*(x+2)*diff(y(x),x)+(x^2+4*x+6)*y(x)=2*x*exp(x),y(x), singsol=a
```

$$y(x) = e^x x (c_1 x^2 + c_2 x + 1)$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 22

```
DSolve[x^2*y'[x]-2*x*(x+2)*y'[x]+(x^2+4*x+6)*y[x]==2*x*Exp[x],y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow e^x x (c_2 x^2 + c_1 x + 1)$$

10.27 problem 27

Internal problem ID [1181]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$x^2 y'' - 4y'x + (x^2 + 6)y = x^4$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+(x^2+6)*y(x)=x^4,y(x), singsol=all)
```

$$y(x) = x^2(1 + \sin(x) c_2 + \cos(x) c_1)$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 38

```
DSolve[x^2*y''[x]-4*x*y'[x]+(x^2+6)*y[x]==x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x^2(2c_1e^{-ix} - ic_2e^{ix} + 2)$$

10.28 problem 28

Internal problem ID [1182]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x - 1)y'' - y'x + y = 2(x - 1)^2 e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve((x-1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=2*(x-1)^2*exp(x),y(x), singsol=all)
```

$$y(x) = (x^2 + c_1 - 2x) e^x + c_2 x$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 24

```
DSolve[(x-1)*y'[x]-x*y'[x]+y[x]==2*(x-1)^2*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (x^2 - 2x + c_1) - c_2 x$$

10.29 problem 29

Internal problem ID [1183]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$4x^2y'' - 4x(x+1)y' + (2x+3)y = x^{\frac{5}{2}}e^x$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(y(x),x$2)-4*x*(x+1)*diff(y(x),x)+(2*x+3)*y(x)=x^(5/2)*exp(x),y(x), singular
```

$$y(x) = \frac{\sqrt{x}((x+4c_1)e^x + 4c_2)}{4}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 30

```
DSolve[4*x^2*y''[x]-4*x*(x+1)*y'[x]+(2*x+3)*y[x]==x^(5/2)*Exp[x],y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{1}{4}\sqrt{x}(e^x(x-1+4c_2) + 4c_1)$$

10.30 problem 30

Internal problem ID [1184]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(3x - 1)y'' - (3x + 2)y' - (6x - 8)y = (3x - 1)^2 e^{2x}$$

With initial conditions

$$[y(0) = 1, y'(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 28

```
dsolve([(3*x-1)*diff(y(x),x$2)-(3*x+2)*diff(y(x),x)-(6*x-8)*y(x)=(3*x-1)^2*exp(2*x),y(0) = 1
```

$$y(x) = \frac{(3x^2 - 2x + 6)e^{2x}}{6} + \frac{x e^{-x}}{3}$$

✓ Solution by Mathematica

Time used: 0.13 (sec). Leaf size: 34

```
DSolve[{(3*x-1)*y'[x]-(3*x+2)*y[x]-(6*x-8)*y[x]==(3*x-1)^2*Exp[2*x],{y[0]==1,y'[0]==2}},y[x]
```

$$y(x) \rightarrow \frac{1}{6}e^{-x}(e^{3x}(3x^2 - 2x + 6) + 2x)$$

10.31 problem 31

Internal problem ID [1185]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$(x - 1)^2 y'' - 2(x - 1) y' + 2y = (x - 1)^2$$

With initial conditions

$$[y(0) = 3, y'(0) = -6]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 33

```
dsolve([(x-1)^2*diff(y(x),x$2)-2*(x-1)*diff(y(x),x)+2*y(x)=(x-1)^2,y(0) = 3, D(y)(0) = -6],y
```

$$y(x) = (-i\pi x + i\pi + \ln(x - 1)x - \ln(x - 1) + 2x - 3)(x - 1)$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 30

```
DSolve[{(x-1)^2*y'[x]-2*(x-1)*y'[x]+2*y[x]==(x-1)^2,{y[0]==3,y'[0]==-6}},y[x],x,IncludeSing
```

$$y(x) \rightarrow (x - 1)(-i\pi(x - 1) + 2x + (x - 1) \log(x - 1) - 3)$$

10.32 problem 32

Internal problem ID [1186]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _linear, _nonhomogeneous]]`

$$(x - 1)^2 y'' - (x^2 - 1) y' + (x - 1)^3 y = (x - 1)^3 e^x$$

With initial conditions

$$[y(0) = 4, y'(0) = -6]$$

X Solution by Maple

```
dsolve([(x-1)^2*diff(y(x),x$2)-(x^2-1)*diff(y(x),x)+(x-1)^3*y(x)=(x-1)^3*exp(x),y(0) = 4, D
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(x-1)^2*y''[x]-(x^2-1)*y'[x]+(x-1)^3*y[x]==(x-1)^3*Exp[x],{y[0]==4,y'[0]==-6}},y[x],
```

Not solved

10.33 problem 33

Internal problem ID [1187]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$(x - 1)^2 y'' + 4y'x + 2y = 2x$$

With initial conditions

$$[y(0) = 0, y'(0) = -2]$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 67

```
dsolve([(x-1)^2*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=2*x,y(0) = 0, D(y)(0) = -2],y(x), sin
```

$$y(x) = \frac{x^3 - 28 e^{\frac{4}{x-1}} \operatorname{expIntegral}_1(-4) - 7 e^{\frac{4x}{x-1}} + 28 e^{\frac{4}{x-1}} \operatorname{expIntegral}_1\left(\frac{4}{x-1}\right) - 2x^2 - 6x + 7}{3(x-1)^2}$$

✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 71

```
DSolve[{(x-1)^2*y''[x]+4*x*y'[x]+2*y[x]==2*x,{y[0]==0,y'[0]==-2}},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \frac{-28 e^{\frac{4}{x-1}} \operatorname{ExpIntegralEi}\left(-\frac{4}{x-1}\right) + 28 \operatorname{ExpIntegralEi}(4) e^{\frac{4}{x-1}} + x^3 - 2x^2 - 6x - 7 e^{\frac{4x}{x-1}} + 7}{3(x-1)^2}$$

10.34 problem 34

Internal problem ID [1188]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 2y'x - 2y = -2x^2$$

With initial conditions

$$[y(1) = 1, y'(1) = -1]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=-2*x^2,y(1) = 1, D(y)(1) = -1],y(x), sing
```

$$y(x) = \frac{1}{2x^2} + x - \frac{x^2}{2}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 21

```
DSolve[{x^2*y'[x]+2*x*y'[x]-2*y[x]==-2*x^2,{y[1]==1,y'[1]==-1}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\frac{x^2}{2} + \frac{1}{2x^2} + x$$

10.35 problem 35

Internal problem ID [1189]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 5 linear second order equations. Section 5.7 Variation of Parameters. Page 262

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _nonhomogeneous]]`

$$(x + 1)(2x + 3)y'' + 2(2 + x)y' - 2y = (2x + 3)^2$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve([(x+1)*(2*x+3)*diff(y(x),x$2)+2*(x+2)*diff(y(x),x)-2*y(x)=(2*x+3)^2,y(0) = 0, D(y)(0)
```

$$y(x) = \frac{x^2(4x + 9)}{6x + 6}$$

✓ Solution by Mathematica

Time used: 0.694 (sec). Leaf size: 22

```
DSolve[{(x+1)*(2*x+3)*y''[x]+2*(x+2)*y'[x]-2*y[x]==(2*x+3)^2,{y[0]==0,y'[0]==0}},y[x],x,Incl
```

$$y(x) \rightarrow \frac{x^2(4x + 9)}{6(x + 1)}$$

11 Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

11.1 problem 11	360
11.2 problem 12	361
11.3 problem 13	362
11.4 problem 14	363
11.5 problem 15	364
11.6 problem 16	365
11.7 problem 17	366
11.8 problem 18	367
11.9 problem 19	368
11.10 problem 21	369
11.11 problem 22	370
11.12 problem 23	371
11.13 problem 24	372
11.14 problem 25	373
11.15 problem 26	374

11.1 problem 11

Internal problem ID [1190]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2 + x)y'' + y'x + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;  
dsolve((2+x)*diff(y(x),x$2)+x*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{4}x^2 + \frac{1}{8}x^3 + \frac{1}{8}x^4 - \frac{9}{160}x^5\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{40}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(2+x)*y'[x]+x*y'[x]+3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{40} + \frac{x^4}{12} - \frac{x^3}{3} + x \right) + c_1 \left(-\frac{9x^5}{160} + \frac{x^4}{8} + \frac{x^3}{8} - \frac{3x^2}{4} + 1 \right)$$

11.2 problem 12

Internal problem ID [1191]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x^2 + 1)y'' + 3y'x^2 - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
Order:=6;
```

```
dsolve((1+3*x^2)*diff(y(x),x$2)+3*x^2*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x^2 - \frac{1}{3}x^4 - \frac{3}{10}x^5\right)y(0) + \left(x + \frac{1}{3}x^3 - \frac{1}{4}x^4 - \frac{4}{15}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 52

```
AsymptoticDSolveValue[(1+3*x^2)*y'[x]+3*x^2*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{4x^5}{15} - \frac{x^4}{4} + \frac{x^3}{3} + x \right) + c_1 \left(-\frac{3x^5}{10} - \frac{x^4}{3} + x^2 + 1 \right)$$

11.3 problem 13

Internal problem ID [1192]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + 1)y'' + (-3x + 2)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve((1+2*x^2)*diff(y(x),x$2)+(2-3*x)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - 2x^2 + \frac{4}{3}x^3 - \frac{1}{3}x^4 - \frac{1}{3}x^5\right) y(0) + \left(x - x^2 + \frac{1}{2}x^3 - \frac{1}{12}x^4 - \frac{17}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 66

```
AsymptoticDSolveValue[(1+2*x^2)*y'[x]+(2-3*x)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{3} - \frac{x^4}{3} + \frac{4x^3}{3} - 2x^2 + 1 \right) + c_2 \left(-\frac{17x^5}{120} - \frac{x^4}{12} + \frac{x^3}{2} - x^2 + x \right)$$

11.4 problem 14

Internal problem ID [1193]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + (2 - x)y' + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 52

```
Order:=6;
```

```
dsolve((1+x^2)*diff(y(x),x$2)+(2-x)*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{2}x^2 + x^3 - \frac{1}{8}x^4 - \frac{1}{4}x^5\right) y(0) + \left(x - x^2 + \frac{1}{3}x^3 + \frac{1}{12}x^4 - \frac{2}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 64

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+(2-x)*y'[x]+3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{4} - \frac{x^4}{8} + x^3 - \frac{3x^2}{2} + 1 \right) + c_2 \left(-\frac{2x^5}{15} + \frac{x^4}{12} + \frac{x^3}{3} - x^2 + x \right)$$

11.5 problem 15

Internal problem ID [1194]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x^2 + 1)y'' - 2y'x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
Order:=6;
```

```
dsolve((1+3*x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (x^4 - 2x^2 + 1)y(0) + \left(x - \frac{1}{3}x^3 + \frac{4}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[(1+3*x^2)*y'[x]-2*x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{4x^5}{15} - \frac{x^3}{3} + x \right) + c_1 (x^4 - 2x^2 + 1)$$

11.6 problem 16

Internal problem ID [1195]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$xy'' + (2x + 4)y' + (2 + x)y = 0$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 52

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+(4+2*x)*diff(y(x),x)+(2+x)*y(x)=0,y(x),type='series',x=-1);
```

$$y(x) = \left(1 + \frac{(x+1)^2}{2} + \frac{2(x+1)^3}{3} + \frac{7(x+1)^4}{8} + \frac{17(x+1)^5}{15}\right) y(-1) \\ + \left(x+1 + (x+1)^2 + \frac{3(x+1)^3}{2} + 2(x+1)^4 + \frac{103(x+1)^5}{40}\right) D(y)(-1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 81

```
AsymptoticDSolveValue[(x)*y''[x]+(4+2*x)*y'[x]+(2+x)*y[x]==0,y[x],{x,-1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{17}{15}(x+1)^5 + \frac{7}{8}(x+1)^4 + \frac{2}{3}(x+1)^3 + \frac{1}{2}(x+1)^2 + 1 \right) \\ + c_2 \left(\frac{103}{40}(x+1)^5 + 2(x+1)^4 + \frac{3}{2}(x+1)^3 + (x+1)^2 + x + 1 \right)$$

11.7 problem 17

Internal problem ID [1196]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$x^2 y'' + 2y'x - 3yx = 0$$

With the expansion point for the power series method at $x = 2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-3*x*y(x)=0,y(x),type='series',x=2);
```

$$y(x) = \left(1 + \frac{3(-2+x)^2}{4} - \frac{3(-2+x)^3}{8} + \frac{9(-2+x)^4}{32} - \frac{27(-2+x)^5}{160}\right) y(2) \\ + \left(-2+x - \frac{(-2+x)^2}{2} + \frac{(-2+x)^3}{2} - \frac{5(-2+x)^4}{16} + \frac{31(-2+x)^5}{160}\right) D(y)(2) \\ + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^2*y''[x]+2*x*y'[x]-3*x*y[x]==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{27}{160}(x-2)^5 + \frac{9}{32}(x-2)^4 - \frac{3}{8}(x-2)^3 + \frac{3}{4}(x-2)^2 + 1 \right) \\ + c_2 \left(\frac{31}{160}(x-2)^5 - \frac{5}{16}(x-2)^4 + \frac{1}{2}(x-2)^3 - \frac{1}{2}(x-2)^2 + x - 2 \right)$$

11.8 problem 18

Internal problem ID [1197]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$(2 - x)y'' + 2y = 0$$

With initial conditions

$$[y(0) = a_0, y'(0) = a_1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 40

```
Order:=6;  
dsolve([(2-x)*diff(y(x),x$2)+2*y(x)=0,y(0) = a__0, D(y)(0) = a__1],y(x),type='series',x=0);
```

$$y(x) = a_0 + a_1x - \frac{1}{2}a_0x^2 + \left(-\frac{a_1}{6} - \frac{a_0}{12}\right)x^3 + \left(\frac{a_0}{48} - \frac{a_1}{24}\right)x^4 + \left(-\frac{a_1}{240} + \frac{a_0}{96}\right)x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 79

```
AsymptoticDSolveValue[{(2-x)*y''[x]+2*y[x]==0,{y[0]==a0,y'[0]==a1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{1}{20}x^5 \left(\frac{1}{6} \left(\frac{a_0}{2} + a_1 \right) + \frac{a_0}{8} - \frac{a_1}{4} \right) + \frac{1}{12}x^4 \left(\frac{a_0}{4} - \frac{a_1}{2} \right) + \frac{1}{6}x^3 \left(-\frac{a_0}{2} - a_1 \right) - \frac{a_0x^2}{2} + a_0 + a_1x$$

11.9 problem 19

Internal problem ID [1198]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + 2(x - 1)^2 y' + 3y = 0$$

With initial conditions

$$[y(1) = a_0, y'(1) = a_1]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

Order:=6;

```
dsolve([(1+x)*diff(y(x),x$2)+2*(x-1)^2*diff(y(x),x)+3*y(x)=0,y(1) = a__0, D(y)(1) = a__1],y(x),{x,
```

$$y(x) = a_0 + a_1(x - 1) - \frac{3}{4}a_0(x - 1)^2 + \left(\frac{a_0}{8} - \frac{a_1}{4}\right)(x - 1)^3 \\ + \left(\frac{a_0}{16} - \frac{a_1}{48}\right)(x - 1)^4 + \left(\frac{3a_0}{64} + \frac{a_1}{40}\right)(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 95

```
AsymptoticDSolveValue[{(1+x)*y''[x]+2*(x-1)^2*y'[x]+3*y[x]==0,{y[1]==a0,y'[1]==a1}},y[x],{x,
```

$$y(x) \rightarrow \frac{1}{20}(x - 1)^5 \left(\frac{1}{4} \left(\frac{3a_1}{2} - \frac{3a_0}{4} \right) + \frac{9a_0}{8} + \frac{a_1}{8} \right) + \frac{1}{12}(x - 1)^4 \left(\frac{3a_0}{4} - \frac{a_1}{4} \right) \\ + \frac{1}{6}(x - 1)^3 \left(\frac{3a_0}{4} - \frac{3a_1}{2} \right) - \frac{3}{4}a_0(x - 1)^2 + a_0 + a_1(x - 1)$$

11.10 problem 21

Internal problem ID [1199]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1-x)y'' + x(x+4)y' + (2-x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 55

```
Order:=6;
```

```
dsolve(x^2*(1-x)*diff(y(x),x$2)+x*(4+x)*diff(y(x),x)+(2-x)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{\ln(x)(9x + 18x^2 + 3x^3 + O(x^6))c_2 + c_1(1 + 2x + \frac{1}{3}x^2 + O(x^6))x + (1 - 5x - 55x^2 - \frac{53}{3}x^3 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 56

```
AsymptoticDSolveValue[x^2*(1-x)*y'[x]+x*(4+x)*y'[x]+(2-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{3(x^2 + 6x + 3) \log(x)}{x} - \frac{21x^3 + 75x^2 + 15x - 1}{x^2} \right) + c_2 \left(\frac{x}{3} + \frac{1}{x} + 2 \right)$$

11.11 problem 22

Internal problem ID [1200]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' + x(1+2x)y' - (6x+4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

```
Order:=6;
```

```
dsolve(x^2*(1+x)*diff(y(x),x$2)+x*(1+2*x)*diff(y(x),x)-(4+6*x)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = c_1 x^2 (1 + O(x^6)) + \frac{c_2 (\ln(x) (576x^4 + O(x^6)) + (-144 + 192x - 288x^2 + 576x^3 - 576x^4 - 576x^5 + O(x^6)))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 48

```
AsymptoticDSolveValue[x^2*(1+x)*y'[x]+x*(1+2*x)*y'[x]-(4+6*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x^2 + c_1 \left(\frac{3x^4 - 12x^3 + 6x^2 - 4x + 3}{3x^2} - 4x^2 \log(x) \right)$$

11.12 problem 23

Internal problem ID [1201]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' - x(-x^2 - 6x + 1)y' + (x^2 + 6x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 63

Order:=6;

`dsolve(x^2*(1+x)*diff(y(x),x$2)-x*(1-6*x-x^2)*diff(y(x),x)+(1+6*x+x^2)*y(x)=0,y(x),type='ser`

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - 12x + \frac{119}{2}x^2 - \frac{583}{3}x^3 + \frac{1981}{4}x^4 - \frac{80287}{75}x^5 + O(x^6) \right) + \left(17x - \frac{471}{4}x^2 + 445x^3 - \frac{118285}{96}x^4 + \frac{702451}{250}x^5 + O(x^6) \right) c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 114

`AsymptoticDSolveValue[x^2*(1+x)*y'[x]-x*(1-6*x-x^2)*y'[x]+(1+6*x+x^2)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_1 x \left(-\frac{80287x^5}{75} + \frac{1981x^4}{4} - \frac{583x^3}{3} + \frac{119x^2}{2} - 12x + 1 \right) + c_2 \left(x \left(\frac{702451x^5}{250} - \frac{118285x^4}{96} + 445x^3 - \frac{471x^2}{4} + 17x \right) + x \left(-\frac{80287x^5}{75} + \frac{1981x^4}{4} - \frac{583x^3}{3} + \frac{119x^2}{2} - 12x + 1 \right) \log(x) \right)$$

11.13 problem 24

Internal problem ID [1202]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear,`

$$x^2(3x + 1)y'' + x(x^2 + 12x + 2)y' + 2x(x + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
Order:=6;
```

```
dsolve(x^2*(1+3*x)*diff(y(x),x$2)+x*(2+12*x+x^2)*diff(y(x),x)+2*x*(3+x)*y(x)=0,y(x),type='se
```

$$y(x) = c_1 \left(1 - 3x + \frac{26}{3}x^2 - \frac{101}{4}x^3 + \frac{4441}{60}x^4 - \frac{26141}{120}x^5 + O(x^6) \right) \\ + \frac{c_2 \left(1 - 6x + \frac{35}{2}x^2 - \frac{101}{2}x^3 + \frac{1177}{8}x^4 - \frac{17251}{40}x^5 + O(x^6) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 60

```
AsymptoticDSolveValue[x^2*(1+3*x)*y'[x]+x*(2+12*x+x^2)*y'[x]+2*x*(3+x)*y[x]==0,y[x],{x,0,5}
```

$$y(x) \rightarrow c_1 \left(\frac{571x^3}{8} - \frac{49x^2}{2} + \frac{17x}{2} + \frac{1}{x} - 3 \right) + c_2 \left(\frac{4441x^4}{60} - \frac{101x^3}{4} + \frac{26x^2}{3} - 3x + 1 \right)$$

11.14 problem 25

Internal problem ID [1203]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(2x^2 + 1)y'' + x(2x^2 + 4)y' + 2(-x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
Order:=6;
```

```
dsolve(x^2*(1+2*x^2)*diff(y(x),x$2)+x*(4+2*x^2)*diff(y(x),x)+2*(1-x^2)*y(x)=0,y(x),type='ser
```

$$y(x) = \frac{c_1(1 + O(x^6))x + c_2(1 - 3x^2 - \frac{1}{2}x^4 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 25

```
AsymptoticDSolveValue[x^2*(1+2*x^2)*y'[x]+x*(4+2*x^2)*y'[x]+2*(1-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^2}{2} + \frac{1}{x^2} - 3 \right) + \frac{c_2}{x}$$

11.15 problem 26

Internal problem ID [1204]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.1 Exercises. Page 318

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 2)y'' + 2x(x^2 + 5)y' + 2(-x^2 + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(x^2*(2+x^2)*diff(y(x),x$2)+2*x*(x^2+5)*diff(y(x),x)+2*(3-x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1(1 + \frac{1}{8}x^2 + O(x^6))}{x} + \frac{c_2(\ln(x)(2x^2 + \frac{1}{4}x^4 + O(x^6)) + (-2 - \frac{3}{2}x^2 - \frac{1}{4}x^4 + O(x^6)))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 51

```
AsymptoticDSolveValue[x^2*(2+x^2)*y'[x]+2*x*(x^2+5)*y'[x]+2*(3-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4 + 7x^2 + 4}{4x^3} - \frac{(x^2 + 8) \log(x)}{8x} \right) + c_2 \left(\frac{x}{8} + \frac{1}{x} \right)$$

12 Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

12.1 problem 1	377
12.2 problem 2	378
12.3 problem 3	379
12.4 problem 4	380
12.5 problem 5	381
12.6 problem 6	382
12.7 problem 7	383
12.8 problem 8	384
12.9 problem 9(a)	385
12.10problem 10	386
12.11problem 11	387
12.12problem 12	388
12.13problem 13	389
12.14problem 16	390
12.15problem 17	391
12.16problem 18	392
12.17problem 19	393
12.18problem 20	394
12.19problem 21	395
12.20problem 22	396
12.21problem 23	397
12.22problem 24	398
12.23problem 25	399
12.24problem 26	400
12.25problem 27	401
12.26problem 31	402
12.27problem 33	403
12.28problem 34	404
12.29problem 35	405
12.30problem 36	406
12.31problem 37	407
12.32problem 39	408
12.33problem 40	409

12.34problem 41	410
12.35problem 42	411
12.36problem 43	412
12.37problem 44	413

12.1 problem 1

Internal problem ID [1205]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + 6y'x + 6y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)+6*x*diff(y(x),x)+6*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (5x^4 - 3x^2 + 1)y(0) + (3x^5 - 2x^3 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+6*x*y'[x]+6*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(3x^5 - 2x^3 + x) + c_1(5x^4 - 3x^2 + 1)$$

12.2 problem 2

Internal problem ID [1206]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + 2y'x - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x^2 - \frac{1}{3}x^4\right)y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 23

```
AsymptoticDSolveValue[(1+x^2)*y'[x]+2*x*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^4}{3} + x^2 + 1\right) + c_2 x$$

12.3 problem 3

Internal problem ID [1207]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' - 8y'x + 20y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((1+x^2)*diff(y(x),x$2)-8*x*diff(y(x),x)+20*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (5x^4 - 10x^2 + 1)y(0) + \left(x - 2x^3 + \frac{1}{5}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[(1+x^2)*y'[x]-8*x*y'[x]+20*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{5} - 2x^3 + x \right) + c_1(5x^4 - 10x^2 + 1)$$

12.4 problem 4

Internal problem ID [1208]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 8y'x - 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((1-x^2)*diff(y(x),x$2)-8*x*diff(y(x),x)-12*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (15x^4 + 6x^2 + 1)y(0) + \left(x + \frac{10}{3}x^3 + 7x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-8*x*y'[x]-12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(7x^5 + \frac{10x^3}{3} + x\right) + c_1(15x^4 + 6x^2 + 1)$$

12.5 problem 5

Internal problem ID [1209]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + 1)y'' + 7y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((1+2*x^2)*diff(y(x),x$2)+7*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{5}{3}x^4\right) y(0) + \left(x - \frac{3}{2}x^3 + \frac{21}{8}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[(1+2*x^2)*y'[x]+7*x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{21x^5}{8} - \frac{3x^3}{2} + x \right) + c_1 \left(\frac{5x^4}{3} - x^2 + 1 \right)$$

12.6 problem 6

Internal problem ID [1210]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + 2y'x + \frac{y}{4} = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((1+x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)+1/4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{8}x^2 + \frac{25}{384}x^4\right) y(0) + \left(x - \frac{3}{8}x^3 + \frac{147}{640}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+2*x*y'[x]+1/4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{147x^5}{640} - \frac{3x^3}{8} + x \right) + c_1 \left(\frac{25x^4}{384} - \frac{x^2}{8} + 1 \right)$$

12.7 problem 7

Internal problem ID [1211]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [_Gegenbauer]

$$(-x^2 + 1)y'' - 5y'x - 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((1-x^2)*diff(y(x),x$2)-5*x*diff(y(x),x)-4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + 2x^2 + \frac{8}{3}x^4\right)y(0) + \left(x + \frac{3}{2}x^3 + \frac{15}{8}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-5*x*y'[x]-4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{15x^5}{8} + \frac{3x^3}{2} + x \right) + c_1 \left(\frac{8x^4}{3} + 2x^2 + 1 \right)$$

12.8 problem 8

Internal problem ID [1212]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' - 10y'x + 28y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)-10*x*diff(y(x),x)+28*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{35}{3}x^4 - 14x^2\right)y(0) + \left(x - 3x^3 + \frac{3}{5}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 38

```
AsymptoticDSolveValue[(1+x^2)*y'[x]-10*x*y'[x]+28*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{3x^5}{5} - 3x^3 + x\right) + c_1 \left(\frac{35x^4}{3} - 14x^2 + 1\right)$$

12.9 problem 9(a)

Internal problem ID [1213]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 9(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{1}{3}x^4\right) y(0) + \left(x - \frac{1}{2}x^3 + \frac{1}{8}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[y''[x]+x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{8} - \frac{x^3}{2} + x \right) + c_1 \left(\frac{x^4}{3} - x^2 + 1 \right)$$

12.10 problem 10

Internal problem ID [1214]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y'x + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+2*x*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{2}x^2 + \frac{7}{8}x^4\right) y(0) + \left(x - \frac{5}{6}x^3 + \frac{3}{8}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]+2*x*y'[x]+3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{3x^5}{8} - \frac{5x^3}{6} + x \right) + c_1 \left(\frac{7x^4}{8} - \frac{3x^2}{2} + 1 \right)$$

12.11 problem 11

Internal problem ID [1215]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$(x^2 + 1)y'' + y'x + y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+x^2)*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(0) = 2, D(y)(0) = -1],y(x),type='series')
```

$$y(x) = 2 - x - x^2 + \frac{1}{3}x^3 + \frac{5}{12}x^4 - \frac{1}{6}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{(1+x^2)*y''[x]+x*y'[x]+y[x]==0,{y[0]==2,y'[0]==-1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{6} + \frac{5x^4}{12} + \frac{x^3}{3} - x^2 - x + 2$$

12.12 problem 12

Internal problem ID [1216]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + 1)y'' - 9y'x - 6y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+2*x^2)*diff(y(x),x$2)-9*x*diff(y(x),x)-6*y(x)=0,y(0) = 1, D(y)(0) = -1],y(x),type
```

$$y(x) = 1 - x + 3x^2 - \frac{5}{2}x^3 + 5x^4 - \frac{21}{8}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{{(1+2*x^2)*y''[x]-9*x*y'[x]-6*y[x]==0,{y[0]==1,y'[0]==-1}},y[x],{x,0,5
```

$$y(x) \rightarrow -\frac{21x^5}{8} + 5x^4 - \frac{5x^3}{2} + 3x^2 - x + 1$$

12.13 problem 13

Internal problem ID [1217]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$(8x^2 + 1)y'' + 2y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+8*x^2)*diff(y(x),x$2)+2*y(x)=0,y(0) = 2, D(y)(0) = -1],y(x),type='series',x=0);
```

$$y(x) = 2 - x - 2x^2 + \frac{1}{3}x^3 + 3x^4 - \frac{5}{6}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{(1+8*x^2)*y'[x]+2*y[x]==0,{y[0]==2,y'[0]==-1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{5x^5}{6} + 3x^4 + \frac{x^3}{3} - 2x^2 - x + 2$$

12.14 problem 16

Internal problem ID [1218]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)-y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \left(1 + \frac{(x-3)^2}{2} + \frac{(x-3)^4}{24}\right) y(3) + \left(x-3 + \frac{(x-3)^3}{6} + \frac{(x-3)^5}{120}\right) D(y)(3) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 51

```
AsymptoticDSolveValue[y''[x]-y[x]==0,y[x],{x,3,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{24}(x-3)^4 + \frac{1}{2}(x-3)^2 + 1 \right) + c_2 \left(\frac{1}{120}(x-3)^5 + \frac{1}{6}(x-3)^3 + x-3 \right)$$

12.15 problem 17

Internal problem ID [1219]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' - (x - 3)y' - y = 0$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)-(x-3)*diff(y(x),x)-y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \left(1 + \frac{(x-3)^2}{2} + \frac{(x-3)^4}{8}\right) y(3) + \left(x-3 + \frac{(x-3)^3}{3} + \frac{(x-3)^5}{15}\right) D(y)(3) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 51

```
AsymptoticDSolveValue[y''[x]-(x-3)*y'[x]-y[x]==0,y[x],{x,3,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{8}(x-3)^4 + \frac{1}{2}(x-3)^2 + 1 \right) + c_2 \left(\frac{1}{15}(x-3)^5 + \frac{1}{3}(x-3)^3 + x-3 \right)$$

12.16 problem 18

Internal problem ID [1220]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(2x^2 - 4x + 1)y'' + 10(x - 1)y' + 6y = 0$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((1-4*x+2*x^2)*diff(y(x),x$2)+10*(x-1)*diff(y(x),x)+6*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(1 + 3(x-1)^2 + \frac{15(x-1)^4}{2}\right) y(1) + \left(x-1 + \frac{8(x-1)^3}{3} + \frac{32(x-1)^5}{5}\right) D(y)(1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[(1-4*x+2*x^2)*y'[x]+10*(x-1)*y'[x]+6*y[x]==0,y[x],{x,1,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{15}{2}(x-1)^4 + 3(x-1)^2 + 1 \right) + c_2 \left(\frac{32}{5}(x-1)^5 + \frac{8}{3}(x-1)^3 + x-1 \right)$$

12.17 problem 19

Internal problem ID [1221]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 - 8x + 11)y'' - 16(-2 + x)y' + 36y = 0$$

With the expansion point for the power series method at $x = 2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

Order:=6;

```
dsolve((11-8*x+2*x^2)*diff(y(x),x$2)-16*(x-2)*diff(y(x),x)+36*y(x)=0,y(x),type='series',x=2)
```

$$y(x) = \left(1 - 6(-2+x)^2 + \frac{4(-2+x)^4}{3}\right) y(2) + \left(-2+x - \frac{10(-2+x)^3}{9}\right) D(y)(2) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[(11-8*x+2*x^2)*y'[x]-16*(x-2)*y'[x]+36*y[x]==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{4}{3}(x-2)^4 - 6(x-2)^2 + 1 \right) + c_2 \left(-\frac{10}{9}(x-2)^3 + x - 2 \right)$$

12.18 problem 20

Internal problem ID [1222]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(3x^2 + 6x + 5)y'' + 9(x + 1)y' + 3y = 0$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

Order:=6;

```
dsolve((5+6*x+3*x^2)*diff(y(x),x$2)+9*(x+1)*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=-1);
```

$$y(x) = \left(1 - \frac{3(x+1)^2}{4} + \frac{27(x+1)^4}{32}\right) y(-1) + \left(x+1 - (x+1)^3 + \frac{6(x+1)^5}{5}\right) D(y)(-1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 70

```
AsymptoticDSolveValue[(5+6*x+2*x^2)*y'[x]+9*(x+1)*y'[x]+3*y[x]==0,y[x],{x,-1,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{93}{20}(x+1)^5 + \frac{17}{8}(x+1)^4 + (x+1)^3 - \frac{3}{2}(x+1)^2 + 1 \right) + c_2 \left(\frac{9}{5}(x+1)^5 + 2(x+1)^4 - 2(x+1)^3 + x + 1 \right)$$

12.19 problem 21

Internal problem ID [1223]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 4)y'' - y'x - 3y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([(x^2-4)*diff(y(x),x$2)-x*diff(y(x),x)-3*y(x)=0,y(0) = -1, D(y)(0) = 2],y(x),type='se
```

$$y(x) = -1 + 2x + \frac{3}{8}x^2 - \frac{1}{3}x^3 - \frac{3}{128}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 29

```
AsymptoticDSolveValue[{{(x^2-4)*y''[x]-x*y'[x]-3*y[x]==0,{y[0]==-1,y'[0]==2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{3x^4}{128} - \frac{x^3}{3} + \frac{3x^2}{8} + 2x - 1$$

12.20 problem 22

Internal problem ID [1224]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (x - 3)y' + 3y = 0$$

With initial conditions

$$[y(3) = -2, y'(3) = 3]$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
dsolve([diff(y(x),x$2)+(x-3)*diff(y(x),x)+3*y(x)=0,y(3) = -2, D(y)(3) = 3],y(x),type='series
```

$$y(x) = -2 + 3(x - 3) + 3(x - 3)^2 - 2(x - 3)^3 - \frac{5}{4}(x - 3)^4 + \frac{3}{5}(x - 3)^5 + O((x - 3)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[{y''[x]+(x-3)*y'[x]+3*y[x]==0,{y[3]==-2,y'[3]==3}},y[x],{x,3,5}]
```

$$y(x) \rightarrow \frac{3}{5}(x - 3)^5 - \frac{5}{4}(x - 3)^4 - 2(x - 3)^3 + 3(x - 3)^2 + 3(x - 3) - 2$$

12.21 problem 23

Internal problem ID [1225]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x^2 - 6x + 5)y'' + (x - 1)y' + 12y = 0$$

With initial conditions

$$[y(1) = -1, y'(1) = 1]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

```
dsolve([(5-6*x+3*x^2)*diff(y(x),x$2)+(x-1)*diff(y(x),x)+12*y(x)=0,y(1) = -1, D(y)(1) = 1],y(x))
```

$$y(x) = -1 + (x - 1) + 3(x - 1)^2 - \frac{13}{12}(x - 1)^3 - \frac{5}{2}(x - 1)^4 + \frac{143}{160}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[{(5-6*x+3*x^2)*y''[x]+(x-1)*y'[x]+12*y[x]==0,{y[1]==-1,y'[1]==1}},y[x]]
```

$$y(x) \rightarrow \frac{143}{160}(x - 1)^5 - \frac{5}{2}(x - 1)^4 - \frac{13}{12}(x - 1)^3 + 3(x - 1)^2 + x - 2$$

12.22 problem 24

Internal problem ID [1226]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$(4x^2 - 24x + 37)y'' + y = 0$$

With initial conditions

$$[y(3) = 4, y'(3) = -6]$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(4*x^2-24*x+37)*diff(y(x),x$2)+y(x)=0,y(3) = 4, D(y)(3) = -6],y(x),type='series',x=3)
```

$$y(x) = 4 - 6(x - 3) - 2(x - 3)^2 + (x - 3)^3 + \frac{3}{2}(x - 3)^4 - \frac{5}{4}(x - 3)^5 + O((x - 3)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[{(4*x^2-24*x+37)*y'[x]+y[x]==0,{y[3]==4,y'[3]==-6}},y[x],{x,3,5}]
```

$$y(x) \rightarrow -\frac{5}{4}(x - 3)^5 + \frac{3}{2}(x - 3)^4 + (x - 3)^3 - 2(x - 3)^2 - 6(x - 3) + 4$$

12.23 problem 25

Internal problem ID [1227]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 8x + 14)y'' - 8(x - 4)y' + 20y = 0$$

With initial conditions

$$[y(4) = 3, y'(4) = -4]$$

With the expansion point for the power series method at $x = 4$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(x^2-8*x+14)*diff(y(x),x$2)-8*(x-4)*diff(y(x),x)+20*y(x)=0,y(4) = 3, D(y)(4) = -4],y`

$$y(x) = 3 - 4(x - 4) + 15(x - 4)^2 - 4(x - 4)^3 + \frac{15}{4}(x - 4)^4 - \frac{1}{5}(x - 4)^5 + O((x - 4)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 44

`AsymptoticDSolveValue[{(x^2-8*x+14)*y''[x]+8*(x-4)*y'[x]+20*y[x]==0,{y[4]==3,y'[4]==-4}},y[x]`

$$y(x) \rightarrow -\frac{35}{3}(x - 4)^5 + \frac{95}{4}(x - 4)^4 - \frac{28}{3}(x - 4)^3 + 15(x - 4)^2 - 4(x - 4) + 3$$

12.24 problem 26

Internal problem ID [1228]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + 4x + 5)y'' - 20(x + 1)y' + 60y = 0$$

With initial conditions

$$[y(-1) = 3, y'(-1) = -3]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

`dsolve([(2*x^2+4*x+5)*diff(y(x),x$2)-20*(x+1)*diff(y(x),x)+60*y(x)=0,y(-1) = 3, D(y)(-1) = -3])`

$$y(x) = 3 - 3(x + 1) - 30(x + 1)^2 + \frac{20}{3}(x + 1)^3 + 20(x + 1)^4 - \frac{4}{3}(x + 1)^5 + O((x + 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

`AsymptoticDSolveValue[{(2*x^2+4*x+5)*y'[x]-20*(x+1)*y'[x]+60*y[x]==0,{y[-1]==3,y'[-1]==-3}]`

$$y(x) \rightarrow -\frac{4}{3}(x + 1)^5 + 20(x + 1)^4 + \frac{20}{3}(x + 1)^3 - 30(x + 1)^2 - 3(x + 1) + 3$$

12.25 problem 27

Internal problem ID [1229]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 1)y'' + 4y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
Order:=6;  
dsolve((1+x^2)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (x^4 - x^2 + 1)y(0) + (x^5 - x^3 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 30

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+4*x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(x^5 - x^3 + x) + c_1(x^4 - x^2 + 1)$$

12.26 problem 31

Internal problem ID [1230]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + 2\alpha y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 63

```
Order:=6;
```

```
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+2*alpha*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \alpha x^2 + \frac{\alpha(\alpha - 2)x^4}{6}\right) y(0) + \left(x - \frac{(\alpha - 1)x^3}{3} + \frac{(\alpha^2 - 4\alpha + 3)x^5}{30}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+2*\[Alpha]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{\alpha^2 x^5}{30} - \frac{2\alpha x^5}{15} + \frac{x^5}{10} - \frac{\alpha x^3}{3} + \frac{x^3}{3} + x \right) + c_1 \left(\frac{\alpha^2 x^4}{6} - \frac{\alpha x^4}{3} - \alpha x^2 + 1 \right)$$

12.27 problem 33

Internal problem ID [1231]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$y'' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x + \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{12} + x \right) + c_1 \left(\frac{x^3}{6} + 1 \right)$$

12.28 problem 34

Internal problem ID [1232]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-2x^3 + 1)y'' - 10y'x^2 - 8yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

Order:=6;

```
dsolve((1-2*x^3)*diff(y(x),x$2)-10*x^2*diff(y(x),x)-8*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{4x^3}{3}\right)y(0) + \left(x + \frac{3}{2}x^4\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[(1-2*x^3)*y'[x]-10*x^2*y'[x]-8*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{3x^4}{2} + x\right) + c_1 \left(\frac{4x^3}{3} + 1\right)$$

12.29 problem 35

Internal problem ID [1233]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^3 + 1) y'' + 7y'x^2 + 9yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve((1+x^3)*diff(y(x),x$2)+7*x^2*diff(y(x),x)+9*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3x^3}{2}\right) y(0) + \left(x - \frac{4}{3}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[(1+x^3)*y'[x]+7*x^2*y'[x]+9*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{4x^4}{3}\right) + c_1 \left(1 - \frac{3x^3}{2}\right)$$

12.30 problem 36

Internal problem ID [1234]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-2x^3 + 1)y'' + 6y'x^2 + 24yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;
```

```
dsolve((1-2*x^3)*diff(y(x),x$2)+6*x^2*diff(y(x),x)+24*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-4x^3 + 1)y(0) + \left(x - \frac{5}{2}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 26

```
AsymptoticDSolveValue[(1-2*x^3)*y'[x]+6*x^2*y'[x]+24*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{5x^4}{2}\right) + c_1(1 - 4x^3)$$

12.31 problem 37

Internal problem ID [1235]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-x^3 + 1)y'' + 15y'x^2 - 36yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

Order:=6;

```
dsolve((1-x^3)*diff(y(x),x$2)+15*x^2*diff(y(x),x)-36*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (6x^3 + 1)y(0) + \left(x + \frac{7}{4}x^4\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[(1-2*x^3)*y'[x]-10*x^2*y'[x]-8*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{3x^4}{2} + x \right) + c_1 \left(\frac{4x^3}{3} + 1 \right)$$

12.32 problem 39

Internal problem ID [1236]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^5 + 1)y'' + 14y'x^4 + 10yx^3 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

Order:=6;

```
dsolve((1+2*x^5)*diff(y(x),x$2)+14*x^4*diff(y(x),x)+10*x^3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^5}{2}\right)y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 20

```
AsymptoticDSolveValue[(1+2*x^5)*y'[x]+14*x^4*y'[x]+10*x^3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^5}{2}\right) + c_2 x$$

12.33 problem 40

Internal problem ID [1237]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_Emden, _Fowler]`

$$y'' + x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{20}\right) + c_1 \left(1 - \frac{x^4}{12}\right)$$

12.34 problem 41

Internal problem ID [1238]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x^6 + 7yx^5 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^6*diff(y(x),x)+7*x^5*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 10

```
AsymptoticDSolveValue[y''[x]+x^6*y'[x]+7*x^5*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x + c_1$$

12.35 problem 42

Internal problem ID [1239]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^8 + 1)y'' - 16y'x^7 + 72yx^6 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
Order:=6;  
dsolve((1+x^8)*diff(y(x),x$2)-16*x^7*diff(y(x),x)+72*x^6*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 10

```
AsymptoticDSolveValue[(1+x^8)*y''[x]-16*x^7*y'[x]+72*x^6*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x + c_1$$

12.36 problem 43

Internal problem ID [1240]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^6 + 1)y'' - 12y'x^5 - 30yx^4 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve((1-x^6)*diff(y(x),x$2)-12*x^5*diff(y(x),x)-30*x^4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 10

```
AsymptoticDSolveValue[(1-x^6)*y''[x]-12*x^5*y'[x]-30*x^4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x + c_1$$

12.37 problem 44

Internal problem ID [1241]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.2 SERIES SOLUTIONS NEAR AN ORDINARY POINT I. Exercises 7.2. Page 329

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x^5 + 6yx^4 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^5*diff(y(x),x)+6*x^4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 10

```
AsymptoticDSolveValue[y''[x]+x^5*y'[x]+6*x^4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x + c_1$$

13 Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3.

Page 338

13.1 problem 1	416
13.2 problem 2	417
13.3 problem 3	418
13.4 problem 4	419
13.5 problem 5	420
13.6 problem 6	421
13.7 problem 7	422
13.8 problem 8	423
13.9 problem 9	424
13.10problem 10	425
13.11problem 11	426
13.12problem 12	427
13.13problem 13	428
13.14problem 14	429
13.15problem 15	430
13.16problem 19	431
13.17problem 20	432
13.18problem 21	433
13.19problem 22	434
13.20problem 23	435
13.21problem 24	436
13.22problem 25	437
13.23problem 26	438
13.24problem 27	439
13.25problem 28	440
13.26problem 29	441
13.27problem 31(a)	444
13.28problem 31(b)	445
13.29problem 31(c)	446
13.30problem 31(d)	447
13.31problem 31(e)	448
13.32problem 32	449
13.33problem 33	450

13.34problem 34	451
13.35problem 35	452
13.36problem 36	453
13.37problem 37	454
13.38problem 38	455
13.39problem 39 (a)	456
13.40problem 39 (b)	457
13.41problem 40	458
13.42problem 41	459
13.43problem 42	460
13.44problem 43	461
13.45problem 44	462
13.46problem 45	463
13.47problem 46	464
13.48problem 47	465
13.49problem 48	466
13.50problem 49	467

13.1 problem 1

Internal problem ID [1242]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x + 1)y'' + y'x + 2y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;  
dsolve([(1+3*x)*diff(y(x),x$2)+x*diff(y(x),x)+2*y(x)=0,y(0) = 2, D(y)(0) = -3],y(x),type='series')
```

$$y(x) = 2 - 3x - 2x^2 + \frac{7}{2}x^3 - \frac{55}{12}x^4 + \frac{59}{8}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{(1+3*x)*y'[x]+x*y''[x]+2*y[x]==0,{y[0]==2,y'[0]==-3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{59x^5}{8} - \frac{55x^4}{12} + \frac{7x^3}{2} - 2x^2 - 3x + 2$$

13.2 problem 2

Internal problem ID [1243]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(2x^2 + x + 1)y'' + (2 + 8x)y' + 4y = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([(1+x+2*x^2)*diff(y(x),x$2)+(2+8*x)*diff(y(x),x)+4*y(x)=0,y(0) = -1, D(y)(0) = 2],y(x)
```

$$y(x) = -1 + 2x - 4x^3 + 4x^4 + 4x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 23

```
AsymptoticDSolveValue[{{(1+x+2*x^2)*y'[x]+(2+8*x)*y'[x]+4*y[x]==0,{y[0]==-1,y'[0]==2}},y[x],
```

$$y(x) \rightarrow 4x^5 + 4x^4 - 4x^3 + 2x - 1$$

13.3 problem 3

Internal problem ID [1244]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-2x^2 + 1)y'' + (2 - 6x)y' - 2y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

Order:=6;

`dsolve([(1-2*x^2)*diff(y(x),x$2)+(2-6*x)*diff(y(x),x)-2*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),t`

$$y(x) = 1 + x^2 - \frac{2}{3}x^3 + \frac{11}{6}x^4 - \frac{9}{5}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 29

`AsymptoticDSolveValue[{(1-2*x^2)*y'[x]+(2-6*x)*y'[x]-2*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,`

$$y(x) \rightarrow -\frac{9x^5}{5} + \frac{11x^4}{6} - \frac{2x^3}{3} + x^2 + 1$$

13.4 problem 4

Internal problem ID [1245]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x^2 + x + 1)y'' + (2 + 15x)y' + 12y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

Order:=6;

`dsolve([(1+x+3*x^2)*diff(y(x),x$2)+(2+15*x)*diff(y(x),x)+12*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x))`

$$y(x) = x - x^2 - \frac{7}{2}x^3 + \frac{15}{2}x^4 + \frac{45}{8}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 31

`AsymptoticDSolveValue[{(1+x+3*x^2)*y'[x]+(2+15*x)*y'[x]+12*y[x]==0,{y[0]==0,y'[0]==1}},y[x]]`

$$y(x) \rightarrow \frac{45x^5}{8} + \frac{15x^4}{2} - \frac{7x^3}{2} - x^2 + x$$

13.5 problem 5

Internal problem ID [1246]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2+x)y'' + (x+1)y' + 3y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(2+x)*diff(y(x),x$2)+(1+x)*diff(y(x),x)+3*y(x)=0,y(0) = 4, D(y)(0) = 3],y(x),type='s
```

$$y(x) = 4 + 3x - \frac{15}{4}x^2 + \frac{1}{4}x^3 + \frac{11}{16}x^4 - \frac{5}{16}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{(2+x)*y'[x]+(1+x)*y'[x]+3*y[x]==0,{y[0]==4,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{5x^5}{16} + \frac{11x^4}{16} + \frac{x^3}{4} - \frac{15x^2}{4} + 3x + 4$$

13.6 problem 6

Internal problem ID [1247]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 3x + 3)y'' + (6 + 4x)y' + 2y = 0$$

With initial conditions

$$[y(0) = 7, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(3+3*x+x^2)*diff(y(x),x$2)+(6+4*x)*diff(y(x),x)+2*y(x)=0,y(0) = 7, D(y)(0) = 3],y(x),{
```

$$y(x) = 7 + 3x - \frac{16}{3}x^2 + \frac{13}{3}x^3 - \frac{23}{9}x^4 + \frac{10}{9}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{{(3+3*x+x^2)*y''[x]+(6+4*x)*y'[x]+2*y[x]==0,{y[0]==7,y'[0]==3}},y[x],{
```

$$y(x) \rightarrow \frac{10x^5}{9} - \frac{23x^4}{9} + \frac{13x^3}{3} - \frac{16x^2}{3} + 3x + 7$$

13.7 problem 7

Internal problem ID [1248]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 4)y'' + (2 + x)y' + 2y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 5]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(4+x)*diff(y(x),x$2)+(2+x)*diff(y(x),x)+2*y(x)=0,y(0) = 2, D(y)(0) = 5],y(x),type='s
```

$$y(x) = 2 + 5x - \frac{7}{4}x^2 - \frac{3}{16}x^3 + \frac{37}{192}x^4 - \frac{7}{192}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{{(4+x)*y''[x]+(2+x)*y'[x]+2*y[x]==0,{y[0]==4,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{7x^5}{192} + \frac{25x^4}{192} + \frac{x^3}{16} - \frac{7x^2}{4} + 3x + 4$$

13.8 problem 8

Internal problem ID [1249]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(2x^2 - 3x + 2)y'' - (4 - 6x)y' + 2y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = -1]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

Order:=6;

```
dsolve([(2-3*x+2*x^2)*diff(y(x),x$2)-(4-6*x)*diff(y(x),x)+2*y(x)=0,y(1) = 1, D(y)(1) = -1],y
```

$$y(x) = 1 - (x - 1) + \frac{4}{3}(x - 1)^3 - \frac{4}{3}(x - 1)^4 - \frac{4}{5}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[{(2-3*x+2*x^2)*y'[x]-(4-6*x)*y'[x]+2*y[x]==0,{y[1]==1,y'[1]==-1}},y[x]
```

$$y(x) \rightarrow -\frac{4}{5}(x - 1)^5 - \frac{4}{3}(x - 1)^4 + \frac{4}{3}(x - 1)^3 - x + 2$$

13.9 problem 9

Internal problem ID [1250]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + 3x)y'' + 10(x + 1)y' + 8y = 0$$

With initial conditions

$$[y(-1) = 1, y'(-1) = -1]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(3*x+2*x^2)*diff(y(x),x$2)+10*(1+x)*diff(y(x),x)+8*y(x)=0,y(-1) = 1, D(y)(-1) = -1], y(x))`

$$y(x) = 1 - (x + 1) + 4(x + 1)^2 - \frac{13}{3}(x + 1)^3 + \frac{77}{6}(x + 1)^4 - \frac{278}{15}(x + 1)^5 + O((x + 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

`AsymptoticDSolveValue[{(3*x+2*x^2)*y''[x]+10*(1+x)*y'[x]+8*y[x]==0,{y[-1]==1,y'[-1]==-1}},y[x],{x,-1,0}]`

$$y(x) \rightarrow -\frac{278}{15}(x + 1)^5 + \frac{77}{6}(x + 1)^4 - \frac{13}{3}(x + 1)^3 + 4(x + 1)^2 - x$$

13.10 problem 10

Internal problem ID [1251]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - x + 1)y'' - (-4x + 1)y' + 2y = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = -1]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

```
dsolve([(1-x+x^2)*diff(y(x),x$2)-(1-4*x)*diff(y(x),x)+2*y(x)=0,y(1) = 2, D(y)(1) = -1],y(x),
```

$$y(x) = 2 - (x - 1) - \frac{1}{2}(x - 1)^2 + \frac{5}{3}(x - 1)^3 - \frac{19}{12}(x - 1)^4 + \frac{7}{30}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 44

```
AsymptoticDSolveValue[{{(1-x+x^2)*y''[x]-(1-4*x)*y'[x]+2*y[x]==0,{y[1]==2,y'[1]==-1}},y[x],{x
```

$$y(x) \rightarrow \frac{7}{30}(x - 1)^5 - \frac{19}{12}(x - 1)^4 + \frac{5}{3}(x - 1)^3 - \frac{1}{2}(x - 1)^2 - x + 3$$

13.11 problem 11

Internal problem ID [1252]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(2+x)y'' + (2+x)y' + y = 0$$

With initial conditions

$$[y(-1) = -2, y'(-1) = 3]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(2+x)*diff(y(x),x$2)+(2+x)*diff(y(x),x)+y(x)=0,y(-1) = -2, D(y)(-1) = 3],y(x),type='
```

$$y(x) = -2 + 3(x+1) - \frac{1}{2}(x+1)^2 - \frac{2}{3}(x+1)^3 + \frac{5}{8}(x+1)^4 - \frac{11}{30}(x+1)^5 + O((x+1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{{(2+x)*y''[x]+(2+x)*y'[x]+y[x]==0},{y[-1]==-2,y'[-1]==3}},y[x],{x,-1,5}
```

$$y(x) \rightarrow -\frac{11}{30}(x+1)^5 + \frac{5}{8}(x+1)^4 - \frac{2}{3}(x+1)^3 - \frac{1}{2}(x+1)^2 + 3(x+1) - 2$$

13.12 problem 12

Internal problem ID [1253]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - (6 - 7x) y' + 8y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = -2]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([x^2*diff(y(x),x$2)-(6-7*x)*diff(y(x),x)+8*y(x)=0,y(1) = 1, D(y)(1) = -2],y(x),type='
```

$$y(x) = 1 - 2(x - 1) - 3(x - 1)^2 + 8(x - 1)^3 - 4(x - 1)^4 - \frac{42}{5}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[{x^2*y''[x]-(6-7*x)*y'[x]+8*y[x]==0,{y[1]==1,y'[1]==-2}},y[x],{x,1,5}]
```

$$y(x) \rightarrow -\frac{42}{5}(x - 1)^5 - 4(x - 1)^4 + 8(x - 1)^3 - 3(x - 1)^2 - 2(x - 1) + 1$$

13.13 problem 13

Internal problem ID [1254]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 + x + 1)y'' + (1 + 7x)y' + 2y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([(1+x+2*x^2)*diff(y(x),x$2)+(1+7*x)*diff(y(x),x)+2*y(x)=0,y(1) = 1, D(y)(1) = 0],y(x),{
```

$$y(x) = 1 - \frac{1}{4}(x-1)^2 + \frac{13}{48}(x-1)^3 - \frac{77}{384}(x-1)^4 + \frac{287}{2560}(x-1)^5 + O((x-1)^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 41

```
AsymptoticDSolveValue[{{(1+x+2*x^2)*y''[x]+(1+7*x)*y'[x]+2*y[x]==0},{y[1]==1,y'[1]==0}},y[x],{
```

$$y(x) \rightarrow \frac{287(x-1)^5}{2560} - \frac{77}{384}(x-1)^4 + \frac{13}{48}(x-1)^3 - \frac{1}{4}(x-1)^2 + 1$$

13.14 problem 14

Internal problem ID [1255]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 3)y'' + (1 + 2x)y' - (2 - x)y = 0$$

With initial conditions

$$[y(-1) = 1, y'(-1) = 0]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([(3+x)*diff(y(x),x$2)+(1+2*x)*diff(y(x),x)-(2-x)*y(x)=0,y(-1) = 1, D(y)(-1) = 0],y(x),{
```

$$y(x) = 1 + \frac{3}{4}(x + 1)^2 - \frac{1}{12}(x + 1)^3 - \frac{1}{48}(x + 1)^4 - \frac{1}{120}(x + 1)^5 + O((x + 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[{{(3+x)*y''[x]+(1+2*x)*y'[x]-(2-x)*y[x]==0,{y[-1]==1,y'[-1]==0}},y[x],{
```

$$y(x) \rightarrow -\frac{1}{120}(x + 1)^5 - \frac{1}{48}(x + 1)^4 - \frac{1}{12}(x + 1)^3 + \frac{3}{4}(x + 1)^2 + 1$$

13.15 problem 15

Internal problem ID [1256]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y'x + (2x^2 + 4)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+3*x*diff(y(x),x)+(4+2*x^2)*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='s
```

$$y(x) = 1 - 2x^2 + \frac{3}{2}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{y''[x]+3*x*y'[x]+(4+2*x^2)*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{3x^4}{2} - 2x^2 + 1$$

13.16 problem 19

Internal problem ID [1257]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(4x + 2)y'' - 4y' - (6 + 4x)y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -7]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(2+4*x)*diff(y(x),x$2)-4*diff(y(x),x)-(6+4*x)*y(x)=0,y(0) = 2, D(y)(0) = -7],y(x),t)
```

$$y(x) = 2 - 7x - 4x^2 - \frac{17}{6}x^3 - \frac{3}{4}x^4 - \frac{9}{40}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{(2+4*x)*y'[x]-4*y'[x]-(6+4*x)*y[x]==0,{y[0]==2,y'[0]==-7}},y[x],{x,0
```

$$y(x) \rightarrow -\frac{9x^5}{40} - \frac{3x^4}{4} - \frac{17x^3}{6} - 4x^2 - 7x + 2$$

13.17 problem 20

Internal problem ID [1258]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(1 + 2x)y'' - (1 - 2x)y' - (-2x + 3)y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = -2]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+2*x)*diff(y(x),x$2)-(1-2*x)*diff(y(x),x)-(3-2*x)*y(x)=0,y(1) = 1, D(y)(1) = -2],y
```

$$y(x) = 1 - 2(x - 1) + \frac{1}{2}(x - 1)^2 - \frac{1}{6}(x - 1)^3 + \frac{5}{36}(x - 1)^4 - \frac{73}{1080}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{(1+2*x)*y'[x]-(1-2*x)*y[x]-(3-2*x)*y[x]==0,{y[1]==1,y'[1]==-2}},y[x]
```

$$y(x) \rightarrow -\frac{73(x-1)^5}{1080} + \frac{5}{36}(x-1)^4 - \frac{1}{6}(x-1)^3 + \frac{1}{2}(x-1)^2 - 2(x-1) + 1$$

13.18 problem 21

Internal problem ID [1259]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x + 5)y'' - y' + (x + 5)y = 0$$

With initial conditions

$$[y(-2) = 2, y'(-2) = -1]$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(5+2*x)*diff(y(x),x$2)-diff(y(x),x)+(5+x)*y(x)=0,y(-2) = 2, D(y)(-2) = -1],y(x),type
```

$$y(x) = 2 - (2 + x) - \frac{7}{2}(2 + x)^2 + \frac{4}{3}(2 + x)^3 - \frac{1}{24}(2 + x)^4 + \frac{1}{60}(2 + x)^5 + O((2 + x)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 43

```
AsymptoticDSolveValue[{(5+2*x)*y'[x]-y'[x]+(5+x)*y[x]==0,{y[-2]==2,y'[-2]==-1}},y[x],{x,-2,
```

$$y(x) \rightarrow \frac{1}{60}(x+2)^5 - \frac{1}{24}(x+2)^4 + \frac{4}{3}(x+2)^3 - \frac{7}{2}(x+2)^2 - x$$

13.19 problem 22

Internal problem ID [1260]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 4)y'' - (2x + 4)y' + (x + 6)y = 0$$

With initial conditions

$$[y(-3) = 2, y'(-3) = -2]$$

With the expansion point for the power series method at $x = -3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(4+x)*diff(y(x),x$2)-(4+2*x)*diff(y(x),x)+(6+x)*y(x)=0,y(-3) = 2, D(y)(-3) = -2],y(x),
```

$$y(x) = 2 - 2(x + 3) - (x + 3)^2 + (x + 3)^3 - \frac{11}{12}(x + 3)^4 + \frac{67}{60}(x + 3)^5 + O((x + 3)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[{(4+x)*y''[x]-(4+2*x)*y'[x]+(6+x)*y[x]==0,{y[-3]==2,y'[-3]==-2}},y[x],
```

$$y(x) \rightarrow \frac{67}{60}(x + 3)^5 - \frac{11}{12}(x + 3)^4 + (x + 3)^3 - (x + 3)^2 - 2(x + 3) + 2$$

13.20 problem 23

Internal problem ID [1261]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x + 2)y'' - y'x + 2yx = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([(2+3*x)*diff(y(x),x$2)-x*diff(y(x),x)+2*x*y(x)=0,y(0) = -1, D(y)(0) = 2],y(x),type=''
```

$$y(x) = -1 + 2x + \frac{1}{3}x^3 - \frac{5}{12}x^4 + \frac{2}{5}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 29

```
AsymptoticDSolveValue[{{(2+3*x)*y'[x]-x*y'[x]+2*x*y[x]==0,{y[0]==-1,y'[0]==2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{2x^5}{5} - \frac{5x^4}{12} + \frac{x^3}{3} + 2x - 1$$

13.21 problem 24

Internal problem ID [1262]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x + 3)y'' + 3y' - yx = 0$$

With initial conditions

$$[y(-1) = 2, y'(-1) = -3]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(3+2*x)*diff(y(x),x$2)+3*diff(y(x),x)-x*y(x)=0,y(-1) = 2, D(y)(-1) = -3],y(x),type='
```

$$y(x) = 2 - 3(x + 1) + \frac{7}{2}(x + 1)^2 - 5(x + 1)^3 + \frac{197}{24}(x + 1)^4 - \frac{287}{20}(x + 1)^5 + O((x + 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 44

```
AsymptoticDSolveValue[{(3+2*x)*y'[x]+3*y'[x]-x*y[x]==0,{y[-1]==2,y'[-1]==-3}},y[x],{x,-1,5}
```

$$y(x) \rightarrow -\frac{287}{20}(x + 1)^5 + \frac{197}{24}(x + 1)^4 - 5(x + 1)^3 + \frac{7}{2}(x + 1)^2 - 3(x + 1) + 2$$

13.22 problem 25

Internal problem ID [1263]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x + 3)y'' - 3y' - (2 + x)y = 0$$

With initial conditions

$$[y(-2) = -2, y'(-2) = 3]$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(3+2*x)*diff(y(x),x$2)-3*diff(y(x),x)-(2+x)*y(x)=0,y(-2) = -2, D(y)(-2) = 3],y(x),t
```

$$y(x) = -2 + 3(2 + x) - \frac{9}{2}(2 + x)^2 + \frac{11}{6}(2 + x)^3 + \frac{5}{24}(2 + x)^4 + \frac{7}{20}(2 + x)^5 + O((2 + x)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{(3+2*x)*y'[x]-3*y'[x]-(2+x)*y[x]==0,{y[-2]==-2,y'[-2]==3}},y[x],{x,-
```

$$y(x) \rightarrow \frac{7}{20}(x + 2)^5 + \frac{5}{24}(x + 2)^4 + \frac{11}{6}(x + 2)^3 - \frac{9}{2}(x + 2)^2 + 3(x + 2) - 2$$

13.23 problem 26

Internal problem ID [1264]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(10 - 2x)y'' + (x + 1)y = 0$$

With initial conditions

$$[y(2) = 2, y'(2) = -4]$$

With the expansion point for the power series method at $x = 2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(10-2*x)*diff(y(x),x$2)+(1+x)*y(x)=0,y(2) = 2, D(y)(2) = -4],y(x),type='series',x=2)
```

$$y(x) = 2 - 4(-2 + x) - \frac{1}{2}(-2 + x)^2 + \frac{2}{9}(-2 + x)^3 + \frac{49}{432}(-2 + x)^4 + \frac{23}{1080}(-2 + x)^5 + O((-2 + x)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{(10-2*x)*y'[x]+(1+x)*y[x]==0,{y[2]==2,y'[2]==-4}},y[x],{x,2,5}]
```

$$y(x) \rightarrow \frac{23(x-2)^5}{1080} + \frac{49}{432}(x-2)^4 + \frac{2}{9}(x-2)^3 - \frac{1}{2}(x-2)^2 - 4(x-2) + 2$$

13.24 problem 27

Internal problem ID [1265]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(7+x)y'' + (2x+8)y' + (x+5)y = 0$$

With initial conditions

$$[y(-4) = 1, y'(-4) = 2]$$

With the expansion point for the power series method at $x = -4$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(7+x)*diff(y(x),x$2)+(8+2*x)*diff(y(x),x)+(5+x)*y(x)=0,y(-4) = 1, D(y)(-4) = 2],y(x),{
```

$$y(x) = 1 + 2(x+4) - \frac{1}{6}(x+4)^2 - \frac{10}{27}(x+4)^3 + \frac{19}{648}(x+4)^4 + \frac{13}{324}(x+4)^5 + O((x+4)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{{(7+x)*y'[x]+(8+2*x)*y'[x]+(5+x)*y[x]==0,{y[-4]==1,y'[-4]==2}},y[x],{
```

$$y(x) \rightarrow \frac{13}{324}(x+4)^5 + \frac{19}{648}(x+4)^4 - \frac{10}{27}(x+4)^3 - \frac{1}{6}(x+4)^2 + 2(x+4) + 1$$

13.25 problem 28

Internal problem ID [1266]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(6 + 4x)y'' + (1 + 2x)y = 0$$

With initial conditions

$$[y(-1) = -1, y'(-1) = 2]$$

With the expansion point for the power series method at $x = -1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(6+4*x)*diff(y(x),x$2)+(1+2*x)*y(x)=0,y(-1) = -1, D(y)(-1) = 2],y(x),type='series',x
```

$$y(x) = -1 + 2(x + 1) - \frac{1}{4}(x + 1)^2 + \frac{1}{2}(x + 1)^3 - \frac{65}{96}(x + 1)^4 + \frac{67}{80}(x + 1)^5 + O((x + 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{{(6+4*x)*y'[x]+(1+2*x)*y[x]==0,{y[-1]==-1,y'[-1]==2}},y[x],{x,-1,5}]
```

$$y(x) \rightarrow \frac{67}{80}(x + 1)^5 - \frac{65}{96}(x + 1)^4 + \frac{1}{2}(x + 1)^3 - \frac{1}{4}(x + 1)^2 + 2(x + 1) - 1$$

13.26 problem 29

Internal problem ID [1267]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(\beta x^2 + \alpha x + 1) y'' + (\delta x + \gamma) y' + \epsilon y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 366

Order:=6;

dsolve((1+alpha*x+beta*x^2)*diff(y(x),x\$2)+(gamma+delta*x)*diff(y(x),x)+epsilon*y(x)=0,y(x),

$$\begin{aligned}
 y(x) = & \left(1 - \frac{\epsilon x^2}{2} + \frac{\epsilon(\alpha + \gamma) x^3}{6} + \frac{\epsilon(-\alpha^2 - \frac{3}{2}\alpha\gamma - \frac{1}{2}\gamma^2 + \beta + \delta + \frac{1}{2}\epsilon) x^4}{12} \right. \\
 & \left. - \frac{\epsilon \left(\frac{(\alpha + \frac{\gamma}{2})\epsilon}{3} - \frac{\gamma^3}{12} - \frac{\alpha\gamma^2}{2} + \frac{(-\frac{11\alpha^2}{4} + 2\beta + \frac{5\delta}{4})\gamma}{3} + \alpha \left(-\frac{\alpha^2}{2} + \beta + \frac{3\delta}{4} \right) \right) x^5}{10} \right) y(0) \\
 & + \left(x - \frac{\gamma x^2}{2} + \frac{(\alpha\gamma + \gamma^2 - \delta - \epsilon) x^3}{6} \right. \\
 & \quad \left. + \frac{((2\alpha + 2\gamma)\epsilon - \gamma^3 - 3\alpha\gamma^2 + (-2\alpha^2 + 2\beta + 3\delta)\gamma + 2\delta\alpha) x^4}{24} \right. \\
 & \quad \left. + \frac{(\epsilon^2 + (-6\alpha^2 - 9\alpha\gamma - 3\gamma^2 + 6\beta + 4\delta)\epsilon + \gamma^4 + 6\alpha\gamma^3 + (11\alpha^2 - 8\beta - 6\delta)\gamma^2 - 12\alpha \left(-\frac{\alpha^2}{2} + \beta + \frac{7\delta}{6} \right) \gamma}{120} \right) \\
 & + O(x^6)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 561

AsymptoticDSolveValue[(1+[Alpha]*x+[Beta]*x^2)*y''[x]+([Gamma]+[Delta]*x)*y'[x]+[Epsilon]

$$\begin{aligned}
 y(x) \rightarrow & c_1 \left(-\frac{1}{20}x^5\epsilon(2\alpha\beta - \alpha^3) - \frac{1}{20}x^5\epsilon(\alpha\delta - \gamma(\alpha^2 - \beta)) + \frac{1}{60}\gamma x^5\epsilon(\alpha^2 - \beta) \right. \\
 & + \frac{1}{120}\alpha\gamma^2 x^5\epsilon + \frac{1}{40}\alpha x^5\epsilon(\alpha\gamma - \delta) + \frac{1}{24}\gamma x^5\epsilon(\alpha\gamma - \delta) - \frac{1}{30}\alpha x^5\epsilon^2 + \frac{1}{120}\gamma^3 x^5\epsilon \\
 & - \frac{1}{60}\gamma x^5\epsilon^2 - \frac{1}{12}x^4\epsilon(\alpha^2 - \beta) - \frac{1}{12}x^4\epsilon(\alpha\gamma - \delta) - \frac{1}{24}\alpha\gamma x^4\epsilon - \frac{1}{24}\gamma^2 x^4\epsilon + \frac{x^4\epsilon^2}{24} \\
 & \left. + \frac{1}{6}\alpha x^3\epsilon + \frac{1}{6}\gamma x^3\epsilon - \frac{x^2\epsilon}{2} + 1 \right) \\
 & + c_2 \left(\frac{1}{60}\gamma x^5(\gamma(\alpha^2 - \beta) - \alpha\delta) - \frac{1}{20}\gamma x^5(\alpha\delta - \gamma(\alpha^2 - \beta)) - \frac{1}{20}x^5\epsilon(\alpha^2 - \beta) \right. \\
 & - \frac{1}{20}x^5(\alpha^3(-\gamma) + \alpha^2\delta + 2\alpha\beta\gamma - \beta\delta) + \frac{1}{24}\gamma^2 x^5(\alpha\gamma - \delta) - \frac{1}{120}\gamma^2 x^5(\delta - \alpha\gamma) \\
 & - \frac{1}{40}x^5\epsilon(\alpha\gamma - \delta) + \frac{1}{120}x^5\epsilon(\delta - \alpha\gamma) - \frac{1}{40}x^5(\alpha\gamma - \delta)(\delta - \alpha\gamma) - \frac{1}{24}\alpha\gamma x^5\epsilon + \frac{\gamma^4 x^5}{120} \\
 & - \frac{1}{40}\gamma^2 x^5\epsilon + \frac{x^5\epsilon^2}{120} - \frac{1}{12}x^4(\gamma(\alpha^2 - \beta) - \alpha\delta) - \frac{1}{12}\gamma x^4(\alpha\gamma - \delta) + \frac{1}{24}\gamma x^4(\delta - \alpha\gamma) \\
 & \left. + \frac{1}{12}\alpha x^4\epsilon - \frac{\gamma^3 x^4}{24} + \frac{1}{12}\gamma x^4\epsilon - \frac{1}{6}x^3(\delta - \alpha\gamma) + \frac{\gamma^2 x^3}{6} - \frac{x^3\epsilon}{6} - \frac{\gamma x^2}{2} + x \right)
 \end{aligned}$$

13.27 problem 31(a)

Internal problem ID [1268]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 31(a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(2x^2 + 3x + 1)y'' + (6 + 8x)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve((1+3*x+2*x^2)*diff(y(x),x$2)+(6+8*x)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (30x^5 - 14x^4 + 6x^3 - 2x^2 + 1)y(0) + (31x^5 - 15x^4 + 7x^3 - 3x^2 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 54

```
AsymptoticDSolveValue[(1+3*x+2*x^2)*y'[x]+(6+8*x)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(30x^5 - 14x^4 + 6x^3 - 2x^2 + 1) + c_2(31x^5 - 15x^4 + 7x^3 - 3x^2 + x)$$

13.28 problem 31(b)

Internal problem ID [1269]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 31(b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(6x^2 - 5x + 1)y'' - (10 - 24x)y' + 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 54

Order:=6;

```
dsolve((1-5*x+6*x^2)*diff(y(x),x$2)-(10-24*x)*diff(y(x),x)+12*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = (-390x^5 - 114x^4 - 30x^3 - 6x^2 + 1)y(0) + (211x^5 + 65x^4 + 19x^3 + 5x^2 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 54

```
AsymptoticDSolveValue[(1-5*x+6*x^2)*y'[x]-(10-24*x)*y'[x]+12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(-390x^5 - 114x^4 - 30x^3 - 6x^2 + 1) + c_2(211x^5 + 65x^4 + 19x^3 + 5x^2 + x)$$

13.29 problem 31(c)

Internal problem ID [1270]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 31(c).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(4x^2 - 4x + 1)y'' - (8 - 16x)y' + 8y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=6;
```

```
dsolve((1-4*x+4*x^2)*diff(y(x),x$2)-(8-16*x)*diff(y(x),x)+8*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-128x^5 - 48x^4 - 16x^3 - 4x^2 + 1)y(0) + (80x^5 + 32x^4 + 12x^3 + 4x^2 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 54

```
AsymptoticDSolveValue[(1-4*x+4*x^2)*y'[x]-(8-16*x)*y'[x]+8*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(-128x^5 - 48x^4 - 16x^3 - 4x^2 + 1) + c_2(80x^5 + 32x^4 + 12x^3 + 4x^2 + x)$$

13.30 problem 31(d)

Internal problem ID [1271]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 31(d).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 4x + 4)y'' + (8 + 4x)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

Order:=6;

```
dsolve((4+4*x+x^2)*diff(y(x),x$2)+(8+4*x)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{4}x^2 + \frac{1}{4}x^3 - \frac{3}{16}x^4 + \frac{1}{8}x^5\right) y(0) \\ + \left(x - x^2 + \frac{3}{4}x^3 - \frac{1}{2}x^4 + \frac{5}{16}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 68

```
AsymptoticDSolveValue[(4+4*x+x^2)*y'[x]+(8+4*x)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{8} - \frac{3x^4}{16} + \frac{x^3}{4} - \frac{x^2}{4} + 1 \right) + c_2 \left(\frac{5x^5}{16} - \frac{x^4}{2} + \frac{3x^3}{4} - x^2 + x \right)$$

13.31 problem 31(e)

Internal problem ID [1272]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 31(e).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(3x^2 + 8x + 4)y'' + (16 + 12x)y' + 6y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

Order:=6;

```
dsolve((4+8*x+3*x^2)*diff(y(x),x$2)+(16+12*x)*diff(y(x),x)+6*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{4}x^2 + \frac{3}{2}x^3 - \frac{39}{16}x^4 + \frac{15}{4}x^5\right)y(0) \\ + \left(x - 2x^2 + \frac{13}{4}x^3 - 5x^4 + \frac{121}{16}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 66

```
AsymptoticDSolveValue[(4+8*x+3*x^2)*y'[x]+(16+12*x)*y'[x]+6*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{15x^5}{4} - \frac{39x^4}{16} + \frac{3x^3}{2} - \frac{3x^2}{4} + 1 \right) + c_2 \left(\frac{121x^5}{16} - 5x^4 + \frac{13x^3}{4} - 2x^2 + x \right)$$

13.32 problem 32

Internal problem ID [1273]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 2y'x + (2x^2 + 3)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+2*x*diff(y(x),x)+(3+2*x^2)*y(x)=0,y(0) = 1, D(y)(0) = -2],y(x),type=''
```

$$y(x) = 1 - 2x - \frac{3}{2}x^2 + \frac{5}{3}x^3 + \frac{17}{24}x^4 - \frac{11}{20}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{y''[x]+2*x*y'[x]+(3+2*x^2)*y[x]==0,{y[0]==1,y'[0]==-2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{11x^5}{20} + \frac{17x^4}{24} + \frac{5x^3}{3} - \frac{3x^2}{2} - 2x + 1$$

13.33 problem 33

Internal problem ID [1274]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 3y'x + (2x^2 + 5)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)-3*x*diff(y(x),x)+(5+2*x^2)*y(x)=0,y(0) = 1, D(y)(0) = -2],y(x),type=''
```

$$y(x) = 1 - 2x - \frac{5}{2}x^2 + \frac{2}{3}x^3 - \frac{3}{8}x^4 + \frac{1}{3}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{y''[x]-3*x*y'[x]+(5+2*x^2)*y[x]==0,{y[0]==1,y'[0]==-2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{3} - \frac{3x^4}{8} + \frac{2x^3}{3} - \frac{5x^2}{2} - 2x + 1$$

13.34 problem 34

Internal problem ID [1275]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 5y'x - (-x^2 + 3)y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

```
dsolve([diff(y(x),x$2)+5*x*diff(y(x),x)-(3-x^2)*y(x)=0,y(0) = 6, D(y)(0) = -2],y(x),type='se
```

$$y(x) = 6 - 2x + 9x^2 + \frac{2}{3}x^3 - \frac{23}{4}x^4 - \frac{3}{10}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{y'[x]+5*x*y'[x]-(3-x^2)*y[x]==0,{y[0]==6,y'[0]==-2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{3x^5}{10} - \frac{23x^4}{4} + \frac{2x^3}{3} + 9x^2 - 2x + 6$$

13.35 problem 35

Internal problem ID [1276]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$y'' - 2y'x - (3x^2 + 2)y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -5]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

```
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)-(2+3*x^2)*y(x)=0,y(0) = 2, D(y)(0) = -5],y(x),type=''
```

$$y(x) = 2 - 5x + 2x^2 - \frac{10}{3}x^3 + \frac{3}{2}x^4 - \frac{25}{12}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{y''[x]-2*x*y'[x]-(2+3*x^2)*y[x]==0,{y[0]==2,y'[0]==-5}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{25x^5}{12} + \frac{3x^4}{2} - \frac{10x^3}{3} + 2x^2 - 5x + 2$$

13.36 problem 36

Internal problem ID [1277]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 3y'x + (4x^2 + 2)y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 6]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+3*x*diff(y(x),x)+(2+4*x^2)*y(x)=0,y(0) = 3, D(y)(0) = 6],y(x),type='s
```

$$y(x) = 3 + 6x - 3x^2 - 5x^3 + x^4 + \frac{31}{20}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[{y''[x]+3*x*y'[x]+(2+4*x^2)*y[x]==0,{y[0]==3,y'[0]==6}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{31x^5}{20} + x^4 - 5x^3 - 3x^2 + 6x + 3$$

13.37 problem 37

Internal problem ID [1278]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2y'' + 5y'x + (2x^2 + 4)y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

Order:=6;

```
dsolve([2*diff(y(x),x$2)+5*x*diff(y(x),x)+(4+2*x^2)*y(x)=0,y(0) = 3, D(y)(0) = -2],y(x),type
```

$$y(x) = 3 - 2x - 3x^2 + \frac{3}{2}x^3 + \frac{3}{2}x^4 - \frac{49}{80}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{2*y''[x]+5*x*y'[x]+(4+2*x^2)*y[x]==0,{y[0]==3,y'[0]==-2}},y[x],{x,0,5
```

$$y(x) \rightarrow -\frac{49x^5}{80} + \frac{3x^4}{2} + \frac{3x^3}{2} - 3x^2 - 2x + 3$$

13.38 problem 38

Internal problem ID [1279]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3y'' + 2y'x + (-x^2 + 4)y = 0$$

With initial conditions

$$[y(0) = -2, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([3*diff(y(x),x$2)+2*x*diff(y(x),x)+(4-x^2)*y(x)=0,y(0) = -2, D(y)(0) = 3],y(x),type=''
```

$$y(x) = -2 + 3x + \frac{4}{3}x^2 - x^3 - \frac{19}{54}x^4 + \frac{13}{60}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{3*y''[x]+2*x*y'[x]+(4-x^2)*y[x]==0,{y[0]==-2,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{13x^5}{60} - \frac{19x^4}{54} - x^3 + \frac{4x^2}{3} + 3x - 2$$

13.39 problem 39 (a)

Internal problem ID [1280]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 39 (a).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y'x + (4x^2 + 2)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+4*x*diff(y(x),x)+(2+4*x^2)*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='s
```

$$y(x) = 1 - x^2 + \frac{1}{2}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{y''[x]+4*x*y'[x]+(2+4*x^2)*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^4}{2} - x^2 + 1$$

13.40 problem 39 (b)

Internal problem ID [1281]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 39 (b).

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + 4y'x + (4x^2 + 2)y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+4*x*diff(y(x),x)+(2+4*x^2)*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='s
```

$$y(x) = x - x^3 + \frac{1}{2}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{y'[x]+4*x*y'[x]+(2+4*x^2)*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{2} - x^3 + x$$

13.41 problem 40

Internal problem ID [1282]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + y'x^2 + (1 + 2x)y = 0$$

With initial conditions

$$[y(0) = -2, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+x)*diff(y(x),x$2)+x^2*diff(y(x),x)+(1+2*x)*y(x)=0,y(0) = -2, D(y)(0) = 3],y(x),t
```

$$y(x) = -2 + 3x + x^2 - \frac{1}{6}x^3 - \frac{3}{4}x^4 + \frac{31}{120}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{{(1+x)*y'[x]+x^2*y'[x]+(1+2*x)*y[x]==0,{y[0]==-2,y'[0]==3}},y[x],{x,0
```

$$y(x) \rightarrow \frac{31x^5}{120} - \frac{3x^4}{4} - \frac{x^3}{6} + x^2 + 3x - 2$$

13.42 problem 41

Internal problem ID [1283]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (x^2 + 2x + 1)y' + 2y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+(1+2*x+x^2)*diff(y(x),x)+2*y(x)=0,y(0) = 2, D(y)(0) = 3],y(x),type='s
```

$$y(x) = 2 + 3x - \frac{7}{2}x^2 - \frac{5}{6}x^3 + \frac{41}{24}x^4 + \frac{41}{120}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[{y'[x]+(1+2*x+x^2)*y'[x]+2*y[x]==0,{y[0]==2,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{41x^5}{120} + \frac{41x^4}{24} - \frac{5x^3}{6} - \frac{7x^2}{2} + 3x + 2$$

13.43 problem 42

Internal problem ID [1284]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + (x^2 + 2)y' + yx = 0$$

With initial conditions

$$[y(0) = -3, y'(0) = 5]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

```
dsolve([(1+x^2)*diff(y(x),x$2)+(2+x^2)*diff(y(x),x)+x*y(x)=0,y(0) = -3, D(y)(0) = 5],y(x),ty
```

$$y(x) = -3 + 5x - 5x^2 + \frac{23}{6}x^3 - \frac{23}{12}x^4 + \frac{11}{30}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{(1+x^2)*y''[x]+(2+x^2)*y'[x]+x*y[x]==0,{y[0]==-3,y'[0]==5}},y[x],{x,0
```

$$y(x) \rightarrow \frac{11x^5}{30} - \frac{23x^4}{12} + \frac{23x^3}{6} - 5x^2 + 5x - 3$$

13.44 problem 43

Internal problem ID [1285]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' + (2x^2 - 3x + 1)y' - (x - 4)y = 0$$

With initial conditions

$$[y(1) = -2, y'(1) = 3]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(1+x)*diff(y(x),x$2)+(1-3*x+2*x^2)*diff(y(x),x)-(x-4)*y(x)=0,y(1) = -2, D(y)(1) = 3]`

$$y(x) = -2 + 3(x - 1) + \frac{3}{2}(x - 1)^2 - \frac{17}{12}(x - 1)^3 - \frac{1}{12}(x - 1)^4 + \frac{1}{8}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

`AsymptoticDSolveValue[{(1+x)*y''[x]+(1-3*x+x^2)*y'[x]-(x-4)*y[x]==0,{y[1]==-2,y'[1]==3}},y[x]`

$$y(x) \rightarrow -\frac{13}{240}(x - 1)^5 - \frac{1}{96}(x - 1)^4 - \frac{2}{3}(x - 1)^3 + \frac{9}{4}(x - 1)^2 + 3(x - 1) - 2$$

13.45 problem 44

Internal problem ID [1286]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (3x^2 + 12x + 13)y' + (2x + 5)y = 0$$

With initial conditions

$$[y(-2) = 2, y'(-2) = -3]$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

`Order:=6;`

`dsolve([diff(y(x),x$2)+(13+12*x+3*x^2)*diff(y(x),x)+(5+2*x)*y(x)=0,y(-2) = 2, D(y)(-2) = -3]`

$$y(x) = 2 - 3(2 + x) + \frac{1}{2}(2 + x)^2 - \frac{1}{3}(2 + x)^3 + \frac{31}{24}(2 + x)^4 - \frac{53}{120}(2 + x)^5 + O((2 + x)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

`AsymptoticDSolveValue[{y'[x]+(13+12*x+3*x^2)*y'[x]+(5+2*x)*y[x]==0,{y[-2]==2,y'[-2]==-3}},y`

$$y(x) \rightarrow -\frac{53}{120}(x+2)^5 + \frac{31}{24}(x+2)^4 - \frac{1}{3}(x+2)^3 + \frac{1}{2}(x+2)^2 - 3(x+2) + 2$$

13.46 problem 45

Internal problem ID [1287]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 45.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(3x^2 + 2x + 1)y'' + (-x^2 + 2)y' + (x + 1)y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(1+2*x+3*x^2)*diff(y(x),x$2)+(2-x^2)*diff(y(x),x)+(1+x)*y(x)=0,y(0) = 1, D(y)(0) = -`

$$y(x) = 1 - 2x + \frac{3}{2}x^2 - \frac{11}{6}x^3 + \frac{15}{8}x^4 - \frac{71}{60}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

`AsymptoticDSolveValue[{(1+2*x+3*x^2)*y'[x]+(2-x^2)*y'[x]+(1+x)*y[x]==0,{y[0]==1,y'[0]==-2}]`

$$y(x) \rightarrow -\frac{71x^5}{60} + \frac{15x^4}{8} - \frac{11x^3}{6} + \frac{3x^2}{2} - 2x + 1$$

13.47 problem 46

Internal problem ID [1288]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 46.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 4x + 3)y'' - (-x^2 + 4x + 5)y' - (2 + x)y = 0$$

With initial conditions

$$[y(-2) = 2, y'(-2) = -1]$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

Order:=6;

`dsolve([(3+4*x+x^2)*diff(y(x),x$2)-(5+4*x-x^2)*diff(y(x),x)-(2+x)*y(x)=0,y(-2) = 2, D(y)(-2)=`

$$y(x) = 2 - (2 + x) - \frac{7}{2}(2 + x)^2 - \frac{43}{6}(2 + x)^3 - \frac{203}{24}(2 + x)^4 - \frac{167}{30}(2 + x)^5 + O((2 + x)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 43

`AsymptoticDSolveValue[{(3+4*x+x^2)*y'[x]-(5+4*x-x^2)*y'[x]-(2+x)*y[x]==0,{y[-2]==2,y'[-2]==`

$$y(x) \rightarrow -\frac{167}{30}(x+2)^5 - \frac{203}{24}(x+2)^4 - \frac{43}{6}(x+2)^3 - \frac{7}{2}(x+2)^2 - x$$

13.48 problem 47

Internal problem ID [1289]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 47.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 2x + 1)y'' + (1 - x)y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([(1+2*x+x^2)*diff(y(x),x$2)+(1-x)*y(x)=0,y(0) = 2, D(y)(0) = -1],y(x),type='series',x
```

$$y(x) = 2 - x - x^2 + \frac{7}{6}x^3 - x^4 + \frac{89}{120}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[{{(1+2*x+x^2)*y'[x]+(1-x)*y[x]==0,{y[0]==2,y'[0]==-1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{89x^5}{120} - x^4 + \frac{7x^3}{6} - x^2 - x + 2$$

13.49 problem 48

Internal problem ID [1290]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 48.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(-2x^2 + x)y'' + (-x^2 + 3x + 1)y' + (2 + x)y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

Order:=6;

`dsolve([(x-2*x^2)*diff(y(x),x$2)+(1+3*x-x^2)*diff(y(x),x)+(2+x)*y(x)=0,y(1) = 1, D(y)(1) = 0`

$$y(x) = 1 + \frac{3}{2}(x-1)^2 + \frac{1}{6}(x-1)^3 - \frac{1}{8}(x-1)^5 + O((x-1)^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 32

`AsymptoticDSolveValue[{(x-2*x^2)*y'[x]+(1+3*x-x^2)*y'[x]+(2+x)*y[x]==0,{y[1]==1,y'[1]==0}},`

$$y(x) \rightarrow -\frac{1}{8}(x-1)^5 + \frac{1}{6}(x-1)^3 + \frac{3}{2}(x-1)^2 + 1$$

13.50 problem 49

Internal problem ID [1291]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.3 SERIES SOLUTIONS NEAR AN ORDINARY POINT II. Exercises 7.3. Page 338

Problem number: 49.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(2x^2 - 11x + 16)y'' + (x^2 - 6x + 10)y' - (2 - x)y = 0$$

With initial conditions

$$[y(3) = 1, y'(3) = -2]$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

Order:=6;

`dsolve([(16-11*x+2*x^2)*diff(y(x),x$2)+(10-6*x+x^2)*diff(y(x),x)-(2-x)*y(x)=0,y(3) = 1, D(y`

$$y(x) = 1 - 2(x - 3) + \frac{1}{2}(x - 3)^2 - \frac{1}{6}(x - 3)^3 + \frac{1}{4}(x - 3)^4 - \frac{1}{6}(x - 3)^5 + O((x - 3)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 46

`AsymptoticDSolveValue[{(16-11*x+2*x^2)*y' '[x]+(10-6*x+x^2)*y' [x]-(2-x)*y[x]==0,{y[3]==1,y' [3`

$$y(x) \rightarrow -\frac{1}{6}(x - 3)^5 + \frac{1}{4}(x - 3)^4 - \frac{1}{6}(x - 3)^3 + \frac{1}{2}(x - 3)^2 - 2(x - 3) + 1$$

14 Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

14.1 problem Example 7.5.1 page 353	470
14.2 problem Example 7.5.2 page 354	471
14.3 problem Example 7.5.3 page 356	472
14.4 problem 1	473
14.5 problem 2	474
14.6 problem 3	475
14.7 problem 4	476
14.8 problem 5	477
14.9 problem 6	478
14.10 problem 7	479
14.11 problem 8	480
14.12 problem 9	481
14.13 problem 10	482
14.14 problem 14	483
14.15 problem 15	484
14.16 problem 16	485
14.17 problem 17	486
14.18 problem 18	487
14.19 problem 19	488
14.20 problem 20	489
14.21 problem 21	490
14.22 problem 22	491
14.23 problem 23	492
14.24 problem 24	493
14.25 problem 25	494
14.26 problem 28	495
14.27 problem 29	496
14.28 problem 30	497
14.29 problem 31	498
14.30 problem 32	499
14.31 problem 33	500
14.32 problem 34	501
14.33 problem 35	502
14.34 problem 36	503

14.35problem 37	504
14.36problem 38	505
14.37problem 39	506
14.38problem 40	507
14.39problem 41	508
14.40problem 42	509
14.41problem 43	510
14.42problem 44	511
14.43problem 45	512
14.44problem 46	513
14.45problem 47	514
14.46problem 48	515
14.47problem 49	516
14.48problem 50	517
14.49problem 51	518
14.50problem 61	519
14.51problem 62	520
14.52problem 63	521
14.53problem 64	522
14.54problem 65	523
14.55problem 66	524
14.56problem 67	525
14.57problem 68	526

14.1 problem Example 7.5.1 page 353

Internal problem ID [1292]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: Example 7.5.1 page 353.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + x + 1)y'' + x(11x^2 + 11x + 9)y' + (7x^2 + 10x + 6)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 45

Order:=6;

```
dsolve(2*x^2*(1+x+x^2)*diff(y(x),x$2)+x*(9+11*x+11*x^2)*diff(y(x),x)+(6+10*x+7*x^2)*y(x)=0,y
```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{1}{8}x^4 + \frac{1}{30}x^5 + O(x^6)\right)}{x^2} + \frac{c_2 \left(1 - \frac{1}{3}x + \frac{2}{5}x^2 - \frac{5}{21}x^3 + \frac{7}{135}x^4 + \frac{76}{1155}x^5 + O(x^6)\right)}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 83

```
AsymptoticDSolveValue[2*x^2*(1+x+x^2)*y'[x]+x*(9+11*x+11*x^2)*y'[x]+(6+10*x+7*x^2)*y[x]==0,
```

$$y(x) \rightarrow \frac{c_2 \left(\frac{x^5}{30} + \frac{x^4}{8} - \frac{x^3}{3} + \frac{x^2}{2} + 1\right)}{x^2} + \frac{c_1 \left(\frac{76x^5}{1155} + \frac{7x^4}{135} - \frac{5x^3}{21} + \frac{2x^2}{5} - \frac{x}{3} + 1\right)}{x^{3/2}}$$

14.2 problem Example 7.5.2 page 354

Internal problem ID [1293]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: Example 7.5.2 page 354.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(x+3)y'' + 5x(x+1)y' - (-4x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

Order:=6;

`dsolve(x^2*(3+x)*diff(y(x),x$2)+5*x*(1+x)*diff(y(x),x)-(1-4*x)*y(x)=0,y(x),type='series',x=0`

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{7}{9}x + \frac{35}{81}x^2 - \frac{455}{2187}x^3 + \frac{1820}{19683}x^4 - \frac{6916}{177147}x^5 + O(x^6)\right) + c_1 \left(1 + x - x^2 + \frac{3}{5}x^3 - \frac{3}{10}x^4 + \frac{3}{22}x^5 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 82

`AsymptoticDSolveValue[x^2*(3+x)*y'[x]+5*x*(1+x)*y'[x]-(1-4*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{6916x^5}{177147} + \frac{1820x^4}{19683} - \frac{455x^3}{2187} + \frac{35x^2}{81} - \frac{7x}{9} + 1 \right) + \frac{c_2 \left(\frac{3x^5}{22} - \frac{3x^4}{10} + \frac{3x^3}{5} - x^2 + x + 1 \right)}{x}$$

14.3 problem Example 7.5.3 page 356

Internal problem ID [1294]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: Example 7.5.3 page 356.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-x^2 + 2)y'' - x(4x^2 + 3)y' + (2 + 2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

Order:=6;

```
dsolve(x^2*(2-x^2)*diff(y(x),x$2)-x*(3+4*x^2)*diff(y(x),x)+(2+2*x)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 \sqrt{x} \left(1 + 2x - \frac{9}{8}x^2 + \frac{7}{4}x^3 - \frac{607}{640}x^4 + \frac{13347}{11200}x^5 + O(x^6) \right) \\ + c_2 x^2 \left(1 - \frac{2}{5}x + \frac{27}{35}x^2 - \frac{34}{105}x^3 + \frac{584}{1155}x^4 - \frac{768}{3575}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 86

```
AsymptoticDSolveValue[x^2*(2-x^2)*y'[x]-x*(3+4*x^2)*y'[x]+(2+2*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{768x^5}{3575} + \frac{584x^4}{1155} - \frac{34x^3}{105} + \frac{27x^2}{35} - \frac{2x}{5} + 1 \right) x^2 \\ + c_2 \left(\frac{13347x^5}{11200} - \frac{607x^4}{640} + \frac{7x^3}{4} - \frac{9x^2}{8} + 2x + 1 \right) \sqrt{x}$$

14.4 problem 1

Internal problem ID [1295]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + x + 1)y'' + x(5x^2 + 3x + 3)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 1243

Order:=6;

```
dsolve(2*x^2*(1+x+x^2)*diff(y(x),x$2)+x*(3+3*x+5*x^2)*diff(y(x),x)+y(x)=0,y(x),type='series')
```

$y(x)$

$$= \frac{c_2 x^{\frac{i\sqrt{7}}{4}} \left(1 + \frac{1}{2+i\sqrt{7}}x + \frac{1}{4} \frac{11-i\sqrt{7}}{(2+i\sqrt{7})(i\sqrt{7}+4)} x^2 - \frac{1}{12} \frac{49i\sqrt{7}+89}{(2+i\sqrt{7})(i\sqrt{7}+4)(i\sqrt{7}+6)} x^3 + \frac{1}{48} \frac{395i\sqrt{7}-1553}{(2+i\sqrt{7})(i\sqrt{7}+4)(i\sqrt{7}+6)(i\sqrt{7}+8)} x^4 \right)}{1}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 4838

```
AsymptoticDSolveValue[2*x^2*(1+x+x^2)*y'[x]+x*(3+3*x+5*x^2)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

Too large to display

14.5 problem 2

Internal problem ID [1296]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear]`,

$$3x^2y'' + 2x(-2x^2 + x + 1)y' + (-8x^2 + 2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 44

Order:=6;

```
dsolve(3*x^2*diff(y(x),x$2)+2*x*(1+x-2*x^2)*diff(y(x),x)+(2*x-8*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 x^{\frac{1}{3}} \left(1 - \frac{2}{3}x + \frac{8}{9}x^2 - \frac{40}{81}x^3 + \frac{92}{243}x^4 - \frac{664}{3645}x^5 + O(x^6) \right) \\ + c_2 \left(1 - x + \frac{6}{5}x^2 - \frac{4}{5}x^3 + \frac{32}{55}x^4 - \frac{24}{77}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 83

```
AsymptoticDSolveValue[3*x^2*y''[x]+2*x*(1+x-2*x^2)*y'[x]+(2*x-8*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{664x^5}{3645} + \frac{92x^4}{243} - \frac{40x^3}{81} + \frac{8x^2}{9} - \frac{2x}{3} + 1 \right) \\ + c_2 \left(-\frac{24x^5}{77} + \frac{32x^4}{55} - \frac{4x^3}{5} + \frac{6x^2}{5} - x + 1 \right)$$

14.6 problem 3

Internal problem ID [1297]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 3x + 3)y'' + x(7x^2 + 8x + 5)y' - (-9x^2 - 2x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 43

Order:=6;

```
dsolve(x^2*(3+3*x+x^2)*diff(y(x),x$2)+x*(5+8*x+7*x^2)*diff(y(x),x)-(1-2*x-9*x^2)*y(x)=0,y(x)
```

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{4}{7}x - \frac{7}{45}x^2 + \frac{970}{2457}x^3 - \frac{5707}{22680}x^4 + \frac{13568}{300105}x^5 + O(x^6)\right) + c_1 \left(1 - x^2 + \frac{2}{3}x^3 - \frac{10}{33}x^5 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 74

```
AsymptoticDSolveValue[x^2*(3+3*x+x^2)*y'[x]+x*(5+8*x+7*x^2)*y'[x]-(1-2*x-9*x^2)*y[x]==0,y[x]
```

$$y(x) \rightarrow \frac{c_2 \left(-\frac{10x^5}{33} + \frac{2x^3}{3} - x^2 + 1\right)}{x} + c_1 \sqrt[3]{x} \left(\frac{13568x^5}{300105} - \frac{5707x^4}{22680} + \frac{970x^3}{2457} - \frac{7x^2}{45} - \frac{4x}{7} + 1\right)$$

14.7 problem 4

Internal problem ID [1298]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + x(4x^2 + 2x + 7)y' - (-7x^2 - 4x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(4*x^2*diff(y(x),x$2)+x*(7+2*x+4*x^2)*diff(y(x),x)-(1-4*x-7*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_2 x^{\frac{5}{4}} \left(1 - \frac{1}{2}x - \frac{19}{104}x^2 + \frac{1571}{10608}x^3 + \frac{3225}{198016}x^4 - \frac{752183}{29702400}x^5 + O(x^6)\right) + c_1 \left(1 + 2x - \frac{11}{6}x^2 - \frac{1}{7}x^3 + \frac{895}{1848}x^4 - \dots\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 86

```
AsymptoticDSolveValue[4*x^2*y''[x]+x*(7+2*x+4*x^2)*y'[x]-(1-4*x-7*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(-\frac{752183x^5}{29702400} + \frac{3225x^4}{198016} + \frac{1571x^3}{10608} - \frac{19x^2}{104} - \frac{x}{2} + 1 \right) + \frac{c_2 \left(-\frac{499x^5}{13860} + \frac{895x^4}{1848} - \frac{x^3}{7} - \frac{11x^2}{6} + 2x + 1 \right)}{x}$$

14.8 problem 5

Internal problem ID [1299]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$12x^2(x+1)y'' + x(3x^2 + 35x + 11)y' - (-5x^2 - 10x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 47

Order:=6;

`dsolve(12*x^2*(1+x)*diff(y(x),x$2)+x*(11+35*x+3*x^2)*diff(y(x),x)-(1-10*x-5*x^2)*y(x)=0,y(x))`

$$y(x) = \frac{c_1 \left(1 - x + \frac{7}{8}x^2 - \frac{19}{24}x^3 + \frac{283}{384}x^4 - \frac{1339}{1920}x^5 + O(x^6)\right)}{x^{\frac{1}{4}}} + c_2 x^{\frac{1}{3}} \left(1 - x + \frac{28}{31}x^2 - \frac{1111}{1333}x^3 + \frac{57493}{73315}x^4 - \frac{3668716}{4912105}x^5 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 86

`AsymptoticDSolveValue[12*x^2*(1+x)*y''[x]+x*(11+35*x+3*x^2)*y'[x]-(1-10*x-5*x^2)*y[x]==0,y[x]`

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{3668716x^5}{4912105} + \frac{57493x^4}{73315} - \frac{1111x^3}{1333} + \frac{28x^2}{31} - x + 1 \right) + \frac{c_2 \left(-\frac{1339x^5}{1920} + \frac{283x^4}{384} - \frac{19x^3}{24} + \frac{7x^2}{8} - x + 1 \right)}{\sqrt[4]{x}}$$

14.9 problem 6

Internal problem ID [1300]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear]`,

$$x^2(10x^2 + x + 5)y'' + x(48x^2 + 3x + 4)y' + (36x^2 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 44

Order:=6;

```
dsolve(x^2*(5+x+10*x^2)*diff(y(x),x$2)+x*(4+3*x+48*x^2)*diff(y(x),x)+(x+36*x^2)*y(x)=0,y(x),
```

$$y(x) = c_1 x^{\frac{1}{5}} \left(1 - \frac{6}{25}x - \frac{1217}{625}x^2 + \frac{41972}{46875}x^3 + \frac{1447799}{390625}x^4 - \frac{375253322}{146484375}x^5 + O(x^6) \right) \\ + c_2 \left(1 - \frac{1}{4}x - \frac{35}{18}x^2 + \frac{11}{12}x^3 + \frac{632}{171}x^4 - \frac{2671}{1026}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x^2*(5+x+10*x^2)*y'[x]+x*(4+3*x+48*x^2)*y'[x]+(x+36*x^2)*y[x]==0,y[x]
```

$$y(x) \rightarrow c_1 \sqrt[5]{x} \left(-\frac{375253322x^5}{146484375} + \frac{1447799x^4}{390625} + \frac{41972x^3}{46875} - \frac{1217x^2}{625} - \frac{6x}{25} + 1 \right) \\ + c_2 \left(-\frac{2671x^5}{1026} + \frac{632x^4}{171} + \frac{11x^3}{12} - \frac{35x^2}{18} - \frac{x}{4} + 1 \right)$$

14.10 problem 7

Internal problem ID [1301]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$8x^2y'' - 2x(-x^2 - 4x + 3)y' + (x^2 + 6x + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(8*x^2*diff(y(x),x$2)-2*x*(3-4*x-x^2)*diff(y(x),x)+(3+6*x+x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 + 4x - \frac{131}{24}x^2 + \frac{39}{14}x^3 - \frac{19865}{29568}x^4 + \frac{4421}{110880}x^5 + O(x^6) \right) \\ + c_2 x^{\frac{3}{2}} \left(1 - x + \frac{11}{26}x^2 - \frac{109}{1326}x^3 + \frac{5}{12376}x^4 + \frac{229}{71400}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 86

```
AsymptoticDSolveValue[8*x^2*y'[x]-2*x*(3-4*x-x^2)*y'[x]+(3+6*x+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{4421x^5}{110880} - \frac{19865x^4}{29568} + \frac{39x^3}{14} - \frac{131x^2}{24} + 4x + 1 \right) \sqrt[4]{x} \\ + c_1 \left(\frac{229x^5}{71400} + \frac{5x^4}{12376} - \frac{109x^3}{1326} + \frac{11x^2}{26} - x + 1 \right) x^{3/2}$$

14.11 problem 8

Internal problem ID [1302]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$18x^2(x+1)y'' + 3x(x^2 + 11x + 5)y' - (-5x^2 - 2x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

Order:=6;

```
dsolve(18*x^2*(1+x)*diff(y(x),x$2)+3*x*(5+11*x+x^2)*diff(y(x),x)-(1-2*x-5*x^2)*y(x)=0,y(x),t
```

$$y(x) = \frac{c_2 \sqrt{x} \left(1 - \frac{1}{3}x + \frac{2}{15}x^2 - \frac{5}{63}x^3 + \frac{23}{405}x^4 - \frac{458}{10395}x^5 + O(x^6)\right) + c_1 \left(1 - \frac{1}{12}x^2 + \frac{1}{18}x^3 - \frac{11}{288}x^4 + \frac{31}{1080}x^5 + O(x^6)\right)}{x^{\frac{1}{6}}}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 85

```
AsymptoticDSolveValue[18*x^2*(1+x)*y'[x]+3*x*(5+11*x+x^2)*y'[x]-(1-2*x-5*x^2)*y[x]==0,y[x],
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{458x^5}{10395} + \frac{23x^4}{405} - \frac{5x^3}{63} + \frac{2x^2}{15} - \frac{x}{3} + 1 \right) + \frac{c_2 \left(\frac{31x^5}{1080} - \frac{11x^4}{288} + \frac{x^3}{18} - \frac{x^2}{12} + 1 \right)}{\sqrt[6]{x}}$$

14.12 problem 9

Internal problem ID [1303]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x(x^2 + x + 3)y'' + (-x^2 + x + 4)y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 42

```
Order:=6;
```

```
dsolve(x*(3+x*x^2)*diff(y(x),x$2)+(4+x-x^2)*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{18}x - \frac{71}{405}x^2 + \frac{719}{34992}x^3 - \frac{1678}{1082565}x^4 - \frac{513547}{992023200}x^5 + O(x^6)\right)}{x^{\frac{1}{3}}} + c_2 \left(1 - \frac{1}{14}x^2 + \frac{1}{105}x^3 - \frac{1}{3640}x^4 - \frac{23}{54600}x^5 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 80

```
AsymptoticDSolveValue[x*(3+x*x^2)*y'[x]+(4+x-x^2)*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{23x^5}{54600} - \frac{x^4}{3640} + \frac{x^3}{105} - \frac{x^2}{14} + 1 \right) + \frac{c_2 \left(-\frac{513547x^5}{992023200} - \frac{1678x^4}{1082565} + \frac{719x^3}{34992} - \frac{71x^2}{405} - \frac{x}{18} + 1 \right)}{\sqrt[3]{x}}$$

14.13 problem 10

Internal problem ID [1304]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$10x^2(2x^2 + x + 1)y'' + x(66x^2 + 13x + 13)y' - (10x^2 + 4x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 47

Order:=6;

```
dsolve(10*x^2*(1+x+2*x^2)*diff(y(x),x$2)+x*(13+13*x+66*x^2)*diff(y(x),x)-(1+4*x+10*x^2)*y(x)
```

$$y(x) = \frac{c_1 \left(1 + x + \frac{14}{13}x^2 - \frac{556}{897}x^3 - \frac{5314}{9867}x^4 + \frac{2092186}{2121405}x^5 + O(x^6)\right)}{\sqrt{x}} + c_2 x^{\frac{1}{5}} \left(1 + \frac{3}{17}x - \frac{7}{153}x^2 - \frac{547}{5661}x^3 + \frac{26942}{266067}x^4 + \frac{200432}{3991005}x^5 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 86

```
AsymptoticDSolveValue[10*x^2*(1+x+2*x^2)*y'[x]+x*(13+13*x+66*x^2)*y'[x]-(1+4*x+10*x^2)*y[x]
```

$$y(x) \rightarrow c_1 \sqrt[5]{x} \left(\frac{200432x^5}{3991005} + \frac{26942x^4}{266067} - \frac{547x^3}{5661} - \frac{7x^2}{153} + \frac{3x}{17} + 1 \right) + \frac{c_2 \left(\frac{2092186x^5}{2121405} - \frac{5314x^4}{9867} - \frac{556x^3}{897} + \frac{14x^2}{13} + x + 1 \right)}{\sqrt{x}}$$

14.14 problem 14

Internal problem ID [1305]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' + x(2x + 3)y' - (1 - x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(2*x^2*dif(y(x),x$2)+x*(3+2*x)*dif(y(x),x)-(1-x)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{c_2 x^{\frac{3}{2}} \left(1 - \frac{2}{5}x + \frac{4}{35}x^2 - \frac{8}{315}x^3 + \frac{16}{3465}x^4 - \frac{32}{45045}x^5 + O(x^6) \right) + c_1 \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + O(x^6) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 86

```
AsymptoticDSolveValue[2*x^2*y'[x]+x*(3+2*x)*y'[x]-(1-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{32x^5}{45045} + \frac{16x^4}{3465} - \frac{8x^3}{315} + \frac{4x^2}{35} - \frac{2x}{5} + 1 \right) + \frac{c_2 \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right)}{x}$$

14.15 problem 15

Internal problem ID [1306]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(x+3)y'' + x(4x+5)y' - (1-2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

Order:=6;

```
dsolve(x^2*(3+x)*diff(y(x),x$2)+x*(5+4*x)*diff(y(x),x)-(1-2*x)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{4}{9}x + \frac{14}{81}x^2 - \frac{140}{2187}x^3 + \frac{455}{19683}x^4 - \frac{1456}{177147}x^5 + O(x^6) \right) + c_1(1 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 53

```
AsymptoticDSolveValue[x^2*(3+x)*y'[x]+x*(5+4*x)*y'[x]-(1-2*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{1456x^5}{177147} + \frac{455x^4}{19683} - \frac{140x^3}{2187} + \frac{14x^2}{81} - \frac{4x}{9} + 1 \right) + \frac{c_2}{x}$$

14.16 problem 16

Internal problem ID [1307]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' + x(x+5)y' - (-3x+2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(2*x^2*dif(y(x),x$2)+x*(5+x)*dif(y(x),x)-(2-3*x)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{c_2 x^{\frac{5}{2}} \left(1 - \frac{1}{2}x + \frac{1}{8}x^2 - \frac{1}{48}x^3 + \frac{1}{384}x^4 - \frac{1}{3840}x^5 + O(x^6)\right) + c_1 \left(1 + \frac{1}{3}x + \frac{1}{3}x^2 - \frac{1}{3}x^3 + \frac{1}{9}x^4 - \frac{1}{45}x^5 + O(x^6)\right)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 88

```
AsymptoticDSolveValue[2*x^2*y'[x]+x*(5+x)*y'[x]-(2-3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{x^5}{3840} + \frac{x^4}{384} - \frac{x^3}{48} + \frac{x^2}{8} - \frac{x}{2} + 1 \right) + \frac{c_2 \left(-\frac{x^5}{45} + \frac{x^4}{9} - \frac{x^3}{3} + \frac{x^2}{3} + \frac{x}{3} + 1 \right)}{x^2}$$

14.17 problem 17

Internal problem ID [1308]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2y'' + x(x+1)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(3*x^2*dif(y(x),x$2)+x*(1+x)*dif(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{3}x + \frac{1}{18}x^2 - \frac{1}{162}x^3 + \frac{1}{1944}x^4 - \frac{1}{29160}x^5 + O(x^6) \right)}{x^{\frac{1}{3}}} + c_2 x \left(1 - \frac{1}{7}x + \frac{1}{70}x^2 - \frac{1}{910}x^3 + \frac{1}{14560}x^4 - \frac{1}{276640}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 86

```
AsymptoticDSolveValue[3*x^2*y'[x]+x*(1+x)*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(-\frac{x^5}{276640} + \frac{x^4}{14560} - \frac{x^3}{910} + \frac{x^2}{70} - \frac{x}{7} + 1 \right) + \frac{c_2 \left(-\frac{x^5}{29160} + \frac{x^4}{1944} - \frac{x^3}{162} + \frac{x^2}{18} - \frac{x}{3} + 1 \right)}{\sqrt[3]{x}}$$

14.18 problem 18

Internal problem ID [1309]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' - y'x + (1 - 2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(2*x^2*dif(y(x),x$2)-x*dif(y(x),x)+(1-2*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + 2x + \frac{2}{3}x^2 + \frac{4}{45}x^3 + \frac{2}{315}x^4 + \frac{4}{14175}x^5 + O(x^6) \right) \\ + c_2x \left(1 + \frac{2}{3}x + \frac{2}{15}x^2 + \frac{4}{315}x^3 + \frac{2}{2835}x^4 + \frac{4}{155925}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 84

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-2*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1x \left(\frac{4x^5}{155925} + \frac{2x^4}{2835} + \frac{4x^3}{315} + \frac{2x^2}{15} + \frac{2x}{3} + 1 \right) \\ + c_2\sqrt{x} \left(\frac{4x^5}{14175} + \frac{2x^4}{315} + \frac{4x^3}{45} + \frac{2x^2}{3} + 2x + 1 \right)$$

14.19 problem 19

Internal problem ID [1310]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' + 9y'x - (3x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(9*x^2*diff(y(x),x$2)+9*x*diff(y(x),x)-(1+3*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 x^{\frac{2}{3}} \left(1 + \frac{1}{5}x + \frac{1}{80}x^2 + \frac{1}{2640}x^3 + \frac{1}{147840}x^4 + \frac{1}{12566400}x^5 + O(x^6) \right) + c_1 \left(1 + x + \frac{1}{8}x^2 + \frac{1}{168}x^3 + \frac{1}{6720}x^4 + \frac{1}{436800}x^5 \right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 86

```
AsymptoticDSolveValue[9*x^2*y''[x]+9*x*y'[x]-(1+3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{x^5}{12566400} + \frac{x^4}{147840} + \frac{x^3}{2640} + \frac{x^2}{80} + \frac{x}{5} + 1 \right) + \frac{c_2 \left(\frac{x^5}{436800} + \frac{x^4}{6720} + \frac{x^3}{168} + \frac{x^2}{8} + x + 1 \right)}{\sqrt[3]{x}}$$

14.20 problem 20

Internal problem ID [1311]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2y'' + x(x+1)y' - (3x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

Order:=6;

```
dsolve(3*x^2*dif(y(x),x$2)+x*(1+x)*dif(y(x),x)-(1+3*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{10}{3}x - \frac{35}{18}x^2 - \frac{14}{81}x^3 - \frac{7}{3888}x^4 + \frac{7}{320760}x^5 + O(x^6) \right)}{x^{\frac{1}{3}}} + c_2 x \left(1 + \frac{2}{7}x + \frac{1}{70}x^2 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 65

```
AsymptoticDSolveValue[3*x^2*y'[x]+x*(1+x)*y'[x]-(1+3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{x^2}{70} + \frac{2x}{7} + 1 \right) + \frac{c_2 \left(\frac{7x^5}{320760} - \frac{7x^4}{3888} - \frac{14x^3}{81} - \frac{35x^2}{18} - \frac{10x}{3} + 1 \right)}{\sqrt[3]{x}}$$

14.21 problem 21

Internal problem ID [1312]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x+3)y'' + x(1+5x)y' + (x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

```
dsolve(2*x^2*(3+x)*diff(y(x),x$2)+x*(1+5*x)*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = c_1 x^{\frac{1}{3}} \left(1 - \frac{4}{9}x + \frac{14}{81}x^2 - \frac{140}{2187}x^3 + \frac{455}{19683}x^4 - \frac{1456}{177147}x^5 + O(x^6) \right) \\ + c_2 \sqrt{x} \left(1 - \frac{3}{7}x + \frac{15}{91}x^2 - \frac{15}{247}x^3 + \frac{27}{1235}x^4 - \frac{297}{38285}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 90

```
AsymptoticDSolveValue[2*x^2*(3+x)*y'[x]+x*(1+5*x)*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{297x^5}{38285} + \frac{27x^4}{1235} - \frac{15x^3}{247} + \frac{15x^2}{91} - \frac{3x}{7} + 1 \right) \\ + c_2 \sqrt[3]{x} \left(-\frac{1456x^5}{177147} + \frac{455x^4}{19683} - \frac{140x^3}{2187} + \frac{14x^2}{81} - \frac{4x}{9} + 1 \right)$$

14.22 problem 22

Internal problem ID [1313]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+4)y'' - x(-3x+1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(4+x)*diff(y(x),x$2)-x*(1-3*x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{9}{16}x + \frac{117}{512}x^2 - \frac{663}{8192}x^3 + \frac{13923}{524288}x^4 - \frac{69615}{8388608}x^5 + O(x^6) \right) \\ + c_2 x \left(1 - \frac{3}{7}x + \frac{12}{77}x^2 - \frac{4}{77}x^3 + \frac{24}{1463}x^4 - \frac{24}{4807}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 86

```
AsymptoticDSolveValue[x^2*(4+x)*y'[x]-x*(1-3*x)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(-\frac{24x^5}{4807} + \frac{24x^4}{1463} - \frac{4x^3}{77} + \frac{12x^2}{77} - \frac{3x}{7} + 1 \right) \\ + c_2 \sqrt[4]{x} \left(-\frac{69615x^5}{8388608} + \frac{13923x^4}{524288} - \frac{663x^3}{8192} + \frac{117x^2}{512} - \frac{9x}{16} + 1 \right)$$

14.23 problem 23

Internal problem ID [1314]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2y'' + 5y'x + (x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

```
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - x + \frac{1}{6}x^2 - \frac{1}{90}x^3 + \frac{1}{2520}x^4 - \frac{1}{113400}x^5 + O(x^6)\right)}{x} + \frac{c_2 \left(1 - \frac{1}{3}x + \frac{1}{30}x^2 - \frac{1}{630}x^3 + \frac{1}{22680}x^4 - \frac{1}{1247400}x^5 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 86

```
AsymptoticDSolveValue[2*x^2*y'[x]+5*x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{x^5}{1247400} + \frac{x^4}{22680} - \frac{x^3}{630} + \frac{x^2}{30} - \frac{x}{3} + 1\right)}{\sqrt{x}} + \frac{c_2 \left(-\frac{x^5}{113400} + \frac{x^4}{2520} - \frac{x^3}{90} + \frac{x^2}{6} - x + 1\right)}{x}$$

14.24 problem 24

Internal problem ID [1315]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(4x + 3)y'' + x(5 + 18x)y' - (1 - 12x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(x^2*(3+4*x)*diff(y(x),x$2)+x*(5+18*x)*diff(y(x),x)-(1-12*x)*y(x)=0,y(x),type='series')
```

$y(x)$

$$= \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{22}{9}x + \frac{374}{81}x^2 - \frac{17204}{2187}x^3 + \frac{249458}{19683}x^4 - \frac{3492412}{177147}x^5 + O(x^6) \right) + c_1 (1 + 2x - 6x^2 + 12x^3 - 21x^4 + \frac{37}{11}x^5)}{x}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 80

```
AsymptoticDSolveValue[x^2*(3+4*x)*y'[x]+x*(5+18*x)*y'[x]-(1-12*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{3492412x^5}{177147} + \frac{249458x^4}{19683} - \frac{17204x^3}{2187} + \frac{374x^2}{81} - \frac{22x}{9} + 1 \right) + \frac{c_2 \left(\frac{378x^5}{11} - 21x^4 + 12x^3 - 6x^2 + 2x + 1 \right)}{x}$$

14.25 problem 25

Internal problem ID [1316]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$6x^2y'' + x(10 - x)y' - (2 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 37

Order:=6;

```
dsolve(6*x^2*dif(y(x),x$2)+x*(10-x)*dif(y(x),x)-(2+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 + \frac{2}{21}x + \frac{1}{180}x^2 + \frac{1}{4212}x^3 + \frac{1}{124416}x^4 + \frac{1}{4432320}x^5 + O(x^6)\right) + c_1(1 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 53

```
AsymptoticDSolveValue[6*x^2*y''[x]+x*(10-x)*y'[x]-(2+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{x^5}{4432320} + \frac{x^4}{124416} + \frac{x^3}{4212} + \frac{x^2}{180} + \frac{2x}{21} + 1 \right) + \frac{c_2}{x}$$

14.26 problem 28

Internal problem ID [1317]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+8)y'' + x(3x+2)y' + (x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 47

Order:=6;

```
dsolve(x^2*(8+x)*diff(y(x),x$2)+x*(2+3*x)*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{25}{96}x + \frac{675}{14336}x^2 - \frac{38025}{5046272}x^3 + \frac{732615}{645922816}x^4 - \frac{9230949}{56103010304}x^5 + O(x^6) \right) \\ + c_2 \sqrt{x} \left(1 - \frac{9}{40}x + \frac{5}{128}x^2 - \frac{245}{39936}x^3 + \frac{6615}{7241728}x^4 - \frac{7623}{57933824}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 90

```
AsymptoticDSolveValue[x^2*(8+x)*y'[x]+x*(2+3*x)*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{7623x^5}{57933824} + \frac{6615x^4}{7241728} - \frac{245x^3}{39936} + \frac{5x^2}{128} - \frac{9x}{40} + 1 \right) \\ + c_2 \sqrt[4]{x} \left(-\frac{9230949x^5}{56103010304} + \frac{732615x^4}{645922816} - \frac{38025x^3}{5046272} + \frac{675x^2}{14336} - \frac{25x}{96} + 1 \right)$$

14.27 problem 29

Internal problem ID [1318]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(4x + 3)y'' + x(11 + 4x)y' - (4x + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 41

Order:=6;

```
dsolve(x^2*(3+4*x)*diff(y(x),x$2)+x*(11+4*x)*diff(y(x),x)-(3+4*x)*y(x)=0,y(x),type='series',
```

$$y(x) = \frac{c_1 \left(1 + \frac{32}{7}x + \frac{48}{7}x^2 + O(x^6)\right)}{x^3} + c_2 x^{\frac{1}{3}} \left(1 + \frac{32}{117}x - \frac{28}{1053}x^2 + \frac{4480}{540189}x^3 - \frac{15680}{4113747}x^4 + \frac{401408}{185118615}x^5 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 67

```
AsymptoticDSolveValue[x^2*(3+4*x)*y'[x]+x*(11+4*x)*y'[x]-(3+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_2 \left(\frac{48x^2}{7} + \frac{32x}{7} + 1\right)}{x^3} + c_1 \sqrt[3]{x} \left(\frac{401408x^5}{185118615} - \frac{15680x^4}{4113747} + \frac{4480x^3}{540189} - \frac{28x^2}{1053} + \frac{32x}{117} + 1\right)$$

14.28 problem 30

Internal problem ID [1319]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(3x + 2)y'' + x(4 + 11x)y' - (1 - x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

Order:=6;

```
dsolve(2*x^2*(2+3*x)*diff(y(x),x$2)+x*(4+11*x)*diff(y(x),x)-(1-x)*y(x)=0,y(x),type='series',
```

$y(x)$

$$= \frac{c_1 x \left(1 - \frac{5}{8}x + \frac{55}{96}x^2 - \frac{935}{1536}x^3 + \frac{4301}{6144}x^4 - \frac{124729}{147456}x^5 + O(x^6)\right) + c_2 \left(1 - \frac{5}{4}x + \frac{25}{32}x^2 - \frac{275}{384}x^3 + \frac{4675}{6144}x^4 - \frac{21505}{24576}x^5 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 94

```
AsymptoticDSolveValue[2*x^2*(2+3*x)*y'[x]+x*(4+11*x)*y'[x]-(1-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{935x^{7/2}}{6144} + \frac{55x^{5/2}}{384} - \frac{5x^{3/2}}{32} + \frac{\sqrt{x}}{4} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{4301x^{9/2}}{6144} - \frac{935x^{7/2}}{1536} + \frac{55x^{5/2}}{96} - \frac{5x^{3/2}}{8} + \sqrt{x} \right)$$

14.29 problem 31

Internal problem ID [1320]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(2+x)y'' + 5x(1-x)y' - (-8x+2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(2+x)*diff(y(x),x$2)+5*x*(1-x)*diff(y(x),x)-(2-8*x)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \frac{c_2 x^{\frac{5}{2}} \left(1 - \frac{3}{4}x + \frac{5}{96}x^2 + \frac{5}{4224}x^3 + \frac{5}{292864}x^4 - \frac{1}{3514368}x^5 + O(x^6)\right) + c_1(1 + 8x + 60x^2 - 160x^3 + 40x^4 + O(x^5))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 73

```
AsymptoticDSolveValue[x^2*(2+x)*y'[x]+5*x*(1-x)*y'[x]-(2-8*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_2(40x^4 - 160x^3 + 60x^2 + 8x + 1)}{x^2} + c_1\sqrt{x} \left(-\frac{x^5}{3514368} + \frac{5x^4}{292864} + \frac{5x^3}{4224} + \frac{5x^2}{96} - \frac{3x}{4} + 1 \right)$$

14.30 problem 32

Internal problem ID [1321]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+6)y'' + x(11+4x)y' + (1+2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

Order:=6;

`dsolve(x^2*(6+x)*diff(y(x),x$2)+x*(11+4*x)*diff(y(x),x)+(1+2*x)*y(x)=0,y(x),type='series',x=`

$y(x)$

$$= \frac{c_2 \left(1 - \frac{10}{63}x + \frac{200}{7371}x^2 - \frac{17600}{3781323}x^3 + \frac{3872}{4861701}x^4 - \frac{921536}{6782072895}x^5 + O(x^6) \right) x^{\frac{1}{6}} + c_1 \left(1 - \frac{3}{20}x + \frac{9}{352}x^2 - \frac{105}{23936}x^3 \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 90

`AsymptoticDSolveValue[x^2*(6+x)*y'[x]+x*(11+4*x)*y'[x]+(1+2*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow \frac{c_1 \left(-\frac{921536x^5}{6782072895} + \frac{3872x^4}{4861701} - \frac{17600x^3}{3781323} + \frac{200x^2}{7371} - \frac{10x}{63} + 1 \right)}{\sqrt[3]{x}} + \frac{c_2 \left(-\frac{11907x^5}{92889088} + \frac{6615x^4}{8808448} - \frac{105x^3}{23936} + \frac{9x^2}{352} - \frac{3x}{20} + 1 \right)}{\sqrt{x}}$$

14.31 problem 33

Internal problem ID [1322]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$8x^2y'' + x(x^2 + 2)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(8*x^2*diff(y(x),x$2)+x*(2+x^2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{1}{112} x^2 + \frac{3}{17920} x^4 + O(x^6) \right) + c_2 \sqrt{x} \left(1 - \frac{1}{72} x^2 + \frac{5}{19584} x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 52

```
AsymptoticDSolveValue[8*x^2*y'[x]+x*(2+x^2)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{5x^4}{19584} - \frac{x^2}{72} + 1 \right) + c_2 \sqrt[4]{x} \left(\frac{3x^4}{17920} - \frac{x^2}{112} + 1 \right)$$

14.32 problem 34

Internal problem ID [1323]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$8x^2(-x^2 + 1)y'' + 2x(-13x^2 + 1)y' + (-9x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(8*x^2*(1-x^2)*diff(y(x),x$2)+2*x*(1-13*x^2)*diff(y(x),x)+(1-9*x^2)*y(x)=0,y(x),type='
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 + \frac{1}{2}x^2 + \frac{3}{8}x^4 + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{5}{9}x^2 + \frac{65}{153}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 52

```
AsymptoticDSolveValue[8*x^2*(1-x^2)*y'[x]+2*x*(1-13*x^2)*y'[x]+(1-9*x^2)*y[x]==0,y[x],{x,0,
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{65x^4}{153} + \frac{5x^2}{9} + 1 \right) + c_2 \sqrt[4]{x} \left(\frac{3x^4}{8} + \frac{x^2}{2} + 1 \right)$$

14.33 problem 35

Internal problem ID [1324]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - 2x(-x^2 + 2)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
Order:=6;
```

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-2*x*(2-x^2)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^4 (1 - 2x^2 + 3x^4 + O(x^6)) + c_2 x (12 + 12x^2 - 36x^4 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 34

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]-2*x*(2-x^2)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 (-3x^5 + x^3 + x) + c_2 (3x^8 - 2x^6 + x^4)$$

14.34 problem 36

Internal problem ID [1325]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(x^2 + 3)y'' + (-x^2 + 2)y' - 8yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

Order:=6;

```
dsolve(x*(3+x^2)*diff(y(x),x^2)+(2-x^2)*diff(y(x),x)-8*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{3}} \left(1 + \frac{11}{18}x^2 + \frac{55}{648}x^4 + O(x^6) \right) + c_2 \left(1 + \frac{4}{5}x^2 + \frac{8}{55}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 47

```
AsymptoticDSolveValue[x*(3+x^2)*y'[x]+(2-x^2)*y'[x]-8*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{55x^4}{648} + \frac{11x^2}{18} + 1 \right) + c_2 \left(\frac{8x^4}{55} + \frac{4x^2}{5} + 1 \right)$$

14.35 problem 37

Internal problem ID [1326]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(-x^2 + 1)y'' + x(-19x^2 + 7)y' - (14x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(4*x^2*(1-x^2)*diff(y(x),x$2)+x*(7-19*x^2)*diff(y(x),x)-(1+14*x^2)*y(x)=0,y(x),type='s
```

$$y(x) = \frac{c_2 x^{\frac{5}{4}} \left(1 + \frac{9}{13}x^2 + \frac{51}{91}x^4 + O(x^6)\right) + c_1 \left(1 + \frac{1}{2}x^2 + \frac{3}{8}x^4 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 50

```
AsymptoticDSolveValue[4*x^2*(1-x^2)*y'[x]+x*(7-19*x^2)*y'[x]-(1+14*x^2)*y[x]==0,y[x],{x,0,5
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(\frac{51x^4}{91} + \frac{9x^2}{13} + 1 \right) + \frac{c_2 \left(\frac{3x^4}{8} + \frac{x^2}{2} + 1 \right)}{x}$$

14.36 problem 38

Internal problem ID [1327]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2(-x^2 + 2)y'' + x(-11x^2 + 1)y' + (-5x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(3*x^2*(2-x^2)*diff(y(x),x$2)+x*(1-11*x^2)*diff(y(x),x)+(1-5*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 x^{\frac{1}{3}} \left(1 + \frac{4}{11} x^2 + \frac{40}{253} x^4 + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{3}{8} x^2 + \frac{21}{128} x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 52

```
AsymptoticDSolveValue[3*x^2*(2-x^2)*y''[x]+x*(1-11*x^2)*y'[x]+(1-5*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{21x^4}{128} + \frac{3x^2}{8} + 1 \right) + c_2 \sqrt[3]{x} \left(\frac{40x^4}{253} + \frac{4x^2}{11} + 1 \right)$$

14.37 problem 39

Internal problem ID [1328]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + 2)y'' - x(-7x^2 + 12)y' + (3x^2 + 7)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(2*x^2*(2+x^2)*diff(y(x),x$2)-x*(12-7*x^2)*diff(y(x),x)+(7+3*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \sqrt{x} \left(\left(1 - \frac{9}{8}x^2 + \frac{117}{128}x^4 + O(x^6) \right) x^3 c_1 + \left(12 + 9x^2 - \frac{63}{4}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 58

```
AsymptoticDSolveValue[2*x^2*(2+x^2)*y'[x]-x*(12-7*x^2)*y'[x]+(7+3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{21x^{9/2}}{16} + \frac{3x^{5/2}}{4} + \sqrt{x} \right) + c_2 \left(\frac{117x^{15/2}}{128} - \frac{9x^{11/2}}{8} + x^{7/2} \right)$$

14.38 problem 40

Internal problem ID [1329]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + 2)y'' + x(7x^2 + 4)y' - (-3x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(2*x^2*(2+x^2)*diff(y(x),x$2)+x*(4+7*x^2)*diff(y(x),x)-(1-3*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 x \left(1 - \frac{1}{4}x^2 + \frac{7}{80}x^4 + O(x^6)\right) + c_2 \left(1 - \frac{1}{8}x^2 + \frac{5}{128}x^4 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 58

```
AsymptoticDSolveValue[2*x^2*(2+x^2)*y'[x]+x*(4+7*x^2)*y'[x]-(1-3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{5x^{7/2}}{128} - \frac{x^{3/2}}{8} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{7x^{9/2}}{80} - \frac{x^{5/2}}{4} + \sqrt{x} \right)$$

14.39 problem 41

Internal problem ID [1330]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(2x^2 + 1)y'' + 5x(6x^2 + 1)y' - (-40x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

Order:=6;

```
dsolve(2*x^2*(1+2*x^2)*diff(y(x),x$2)+5*x*(1+6*x^2)*diff(y(x),x)-(2-40*x^2)*y(x)=0,y(x),type
```

$$y(x) = \frac{c_2 x^{\frac{5}{2}} \left(1 - 3x^2 + \frac{15}{2}x^4 + O(x^6)\right) + c_1 \left(1 + 2x^2 - \frac{20}{3}x^4 + O(x^6)\right)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 46

```
AsymptoticDSolveValue[2*x^2*(1+2*x^2)*y'[x]+5*x*(1+6*x^2)*y'[x]-(2-40*x^2)*y[x]==0,y[x],{x,
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{15x^4}{2} - 3x^2 + 1 \right) + \frac{c_2 \left(-\frac{20x^4}{3} + 2x^2 + 1 \right)}{x^2}$$

14.40 problem 42

Internal problem ID [1331]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2(x^2 + 1)y'' + 5x(x^2 + 1)y' - (-5x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(3*x^2*(1+x^2)*diff(y(x),x$2)+5*x*(1+x^2)*diff(y(x),x)-(1-5*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{3}{10}x^2 + \frac{39}{320}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{3}{2}x^2 + \frac{15}{32}x^4 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 50

```
AsymptoticDSolveValue[3*x^2*(1+x^2)*y''[x]+5*x*(1+x^2)*y'[x]-(1-5*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{39x^4}{320} - \frac{3x^2}{10} + 1 \right) + \frac{c_2 \left(\frac{15x^4}{32} - \frac{3x^2}{2} + 1 \right)}{x}$$

14.41 problem 43

Internal problem ID [1332]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(x^2 + 1)y'' + (7x^2 + 4)y' + 8yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

Order:=6;

```
dsolve(x*(1+x^2)*diff(y(x),x$2)+(4+7*x^2)*diff(y(x),x)+8*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{4}{5}x^2 + \frac{24}{35}x^4 + O(x^6) \right) + \frac{c_2(12 - 6x^2 + \frac{9}{2}x^4 + O(x^6))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 42

```
AsymptoticDSolveValue[x*(1+x^2)*y''[x]+(4+7*x^2)*y'[x]+8*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^3} + \frac{3x}{8} - \frac{1}{2x} \right) + c_2 \left(\frac{24x^4}{35} - \frac{4x^2}{5} + 1 \right)$$

14.42 problem 44

Internal problem ID [1333]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 2)y'' + x(x^2 + 3)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(x^2*(2+x^2)*diff(y(x),x$2)+x*(3+x^2)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 x^{\frac{3}{2}} \left(1 - \frac{1}{56}x^2 + \frac{25}{9856}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{1}{2}x^2 + \frac{1}{40}x^4 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 50

```
AsymptoticDSolveValue[x^2*(2+x^2)*y'[x]+x*(3+x^2)*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{25x^4}{9856} - \frac{x^2}{56} + 1 \right) + \frac{c_2 \left(\frac{x^4}{40} - \frac{x^2}{2} + 1 \right)}{x}$$

14.43 problem 45

Internal problem ID [1334]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 45.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + 1)y'' + x(8x^2 + 3)y' - (-4x^2 + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

Order:=6;

```
dsolve(2*x^2*(1+x^2)*diff(y(x),x$2)+x*(3+8*x^2)*diff(y(x),x)-(3-4*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{4}x^2 + \frac{5}{32}x^4 + O(x^6)\right)}{x^{\frac{3}{2}}} + c_2 x \left(1 - \frac{2}{3}x^2 + \frac{20}{39}x^4 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 48

```
AsymptoticDSolveValue[2*x^2*(1+x^2)*y'[x]+x*(3+8*x^2)*y'[x]-(3-4*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{20x^4}{39} - \frac{2x^2}{3} + 1 \right) + \frac{c_2 \left(\frac{5x^4}{32} - \frac{x^2}{4} + 1 \right)}{x^{3/2}}$$

14.44 problem 46

Internal problem ID [1335]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 46.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' + 3x(x^2 + 3)y' - (-5x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(9*x^2*dif(y(x),x$2)+3*x*(3+x^2)*dif(y(x),x)-(1-5*x^2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \frac{c_2 x^{\frac{2}{3}} \left(1 - \frac{1}{8}x^2 + \frac{1}{112}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{72}x^4 + O(x^6)\right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 52

```
AsymptoticDSolveValue[9*x^2*y''[x]+3*x*(3+x^2)*y'[x]-(1-5*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{x^4}{112} - \frac{x^2}{8} + 1 \right) + \frac{c_2 \left(\frac{x^4}{72} - \frac{x^2}{6} + 1 \right)}{\sqrt[3]{x}}$$

14.45 problem 47

Internal problem ID [1336]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 47.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$6x^2y'' + x(6x^2 + 1)y' + (9x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(6*x^2*dif(y(x),x$2)+x*(1+6*x^2)*dif(y(x),x)+(1+9*x^2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = c_1x^{\frac{1}{3}}\left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 + O(x^6)\right) + c_2\sqrt{x}\left(1 - \frac{6}{13}x^2 + \frac{36}{325}x^4 + O(x^6)\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 52

```
AsymptoticDSolveValue[6*x^2*y'[x]+x*(1+6*x^2)*y'[x]+(1+9*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x}\left(\frac{36x^4}{325} - \frac{6x^2}{13} + 1\right) + c_2\sqrt[3]{x}\left(\frac{x^4}{8} - \frac{x^2}{2} + 1\right)$$

14.46 problem 48

Internal problem ID [1337]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 48.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(x^2 + 8)y'' + 7x(x^2 + 2)y' - (-9x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(x^2*(8+x^2)*diff(y(x),x$2)+7*x*(2+x^2)*diff(y(x),x)-(2-9*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_2 x^{\frac{5}{4}} \left(1 - \frac{13}{64}x^2 + \frac{273}{8192}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{1}{3}x^2 + \frac{2}{33}x^4 + O(x^6)\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 50

```
AsymptoticDSolveValue[x^2*(8+x^2)*y'[x]+7*x*(2+x^2)*y'[x]-(2-9*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(\frac{273x^4}{8192} - \frac{13x^2}{64} + 1 \right) + \frac{c_2 \left(\frac{2x^4}{33} - \frac{x^2}{3} + 1 \right)}{x}$$

14.47 problem 49

Internal problem ID [1338]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 49.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2(x^2 + 1)y'' + 3x(13x^2 + 3)y' - (-25x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(9*x^2*(1+x^2)*diff(y(x),x$2)+3*x*(3+13*x^2)*diff(y(x),x)-(1-25*x^2)*y(x)=0,y(x),type=
```

$$y(x) = \frac{c_2 x^{\frac{2}{3}} \left(1 - \frac{3}{4}x^2 + \frac{9}{14}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{2}{3}x^2 + \frac{5}{9}x^4 + O(x^6)\right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 52

```
AsymptoticDSolveValue[9*x^2*(1+x^2)*y'[x]+3*x*(3+13*x^2)*y'[x]-(1-25*x^2)*y[x]==0,y[x],{x,0
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(\frac{9x^4}{14} - \frac{3x^2}{4} + 1 \right) + \frac{c_2 \left(\frac{5x^4}{9} - \frac{2x^2}{3} + 1 \right)}{\sqrt[3]{x}}$$

14.48 problem 50

Internal problem ID [1339]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 50.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 1)y'' + 4x(6x^2 + 1)y' - (-25x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(4*x^2*(1+x^2)*diff(y(x),x$2)+4*x*(1+6*x^2)*diff(y(x),x)-(1-25*x^2)*y(x)=0,y(x),type=''
```

$$y(x) = \frac{c_1 \left(1 - \frac{3}{2}x^2 + \frac{15}{8}x^4 + O(x^6)\right) + c_2 \left(1 - 2x^2 + \frac{8}{3}x^4 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 56

```
AsymptoticDSolveValue[4*x^2*(1+x^2)*y''[x]+4*x*(1+6*x^2)*y'[x]-(1-25*x^2)*y[x]==0,y[x],{x,0,
```

$$y(x) \rightarrow c_1 \left(\frac{8x^{7/2}}{3} - 2x^{3/2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{15x^{9/2}}{8} - \frac{3x^{5/2}}{2} + \sqrt{x} \right)$$

14.49 problem 51

Internal problem ID [1340]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 51.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$8x^2(2x^2 + 1)y'' + 2x(34x^2 + 5)y' - (-30x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(8*x^2*(1+2*x^2)*diff(y(x),x$2)+2*x*(5+34*x^2)*diff(y(x),x)-(1-30*x^2)*y(x)=0,y(x),typ
```

$$y(x) = \frac{c_2 x^{\frac{3}{4}} \left(1 - x^2 + \frac{3}{2}x^4 + O(x^6)\right) + c_1 \left(1 - \frac{2}{5}x^2 + \frac{36}{65}x^4 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 50

```
AsymptoticDSolveValue[8*x^2*(1+2*x^2)*y'[x]+2*x*(5+34*x^2)*y'[x]-(1-30*x^2)*y[x]==0,y[x],{x
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(\frac{3x^4}{2} - x^2 + 1 \right) + \frac{c_2 \left(\frac{36x^4}{65} - \frac{2x^2}{5} + 1 \right)}{\sqrt{x}}$$

14.50 problem 61

Internal problem ID [1341]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 61.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x+1)y'' - x(-3x+1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(2*x^2*(1+x)*diff(y(x),x$2)-x*(1-3*x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-x^5 + x^4 - x^3 + x^2 - x + 1) (c_1\sqrt{x} + c_2x) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 58

```
AsymptoticDSolveValue[2*x^2*(1+x)*y'[x]-x*(1-3*x)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1x(-x^5 + x^4 - x^3 + x^2 - x + 1) + c_2\sqrt{x}(-x^5 + x^4 - x^3 + x^2 - x + 1)$$

14.51 problem 62

Internal problem ID [1342]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 62.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$6x^2(2x^2 + 1)y'' + x(50x^2 + 1)y' + (30x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
Order:=6;
```

```
dsolve(6*x^2*(1+2*x^2)*diff(y(x),x$2)+x*(1+50*x^2)*diff(y(x),x)+(1+30*x^2)*y(x)=0,y(x),type=
```

$$y(x) = (4x^4 - 2x^2 + 1)x^{\frac{1}{3}}(c_2x^{\frac{1}{6}} + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 44

```
AsymptoticDSolveValue[6*x^2*(1+2*x^2)*y'[x]+x*(1+50*x^2)*y'[x]+(1+30*x^2)*y[x]==0,y[x],{x,0
```

$$y(x) \rightarrow c_1\sqrt{x}(4x^4 - 2x^2 + 1) + c_2\sqrt[3]{x}(4x^4 - 2x^2 + 1)$$

14.52 problem 63

Internal problem ID [1343]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 63.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$28x^2(-3x + 1)y'' - 7x(5 + 9x)y' + 7(2 + 9x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(28*x^2*(1-3*x)*diff(y(x),x$2)-7*x*(5+9*x)*diff(y(x),x)+7*(2+9*x)*y(x)=0,y(x),type='series')
```

$$y(x) = (243x^5 + 81x^4 + 27x^3 + 9x^2 + 3x + 1) \left(c_1 x^{\frac{1}{4}} + c_2 x^2 \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 68

```
AsymptoticDSolveValue[28*x^2*(1-3*x)*y''[x]-7*x*(5+9*x)*y'[x]+7*(2+9*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 (243x^5 + 81x^4 + 27x^3 + 9x^2 + 3x + 1) x^{\frac{1}{4}} + c_2 (243x^5 + 81x^4 + 27x^3 + 9x^2 + 3x + 1) x^2$$

14.53 problem 64

Internal problem ID [1344]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 64.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2(x+5)y'' + 9x(5+9x)y' - (5-8x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 47

Order:=6;

```
dsolve(9*x^2*(5+x)*diff(y(x),x$2)+9*x*(5+9*x)*diff(y(x),x)-(5-8*x)*y(x)=0,y(x),type='series')
```

$y(x)$

$$= \frac{c_2 x^{\frac{2}{3}} \left(1 - \frac{11}{25}x + \frac{11}{50}x^2 - \frac{1}{10}x^3 + \frac{29}{700}x^4 - \frac{4727}{297500}x^5 + O(x^6) \right) + c_1 \left(1 + x - \frac{1}{2}x^2 + \frac{17}{70}x^3 - \frac{187}{1750}x^4 + \frac{24497}{568750}x^5 \right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 86

```
AsymptoticDSolveValue[9*x^2*(5+x)*y'[x]+9*x*(5+9*x)*y'[x]-(5-8*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{4727x^5}{297500} + \frac{29x^4}{700} - \frac{x^3}{10} + \frac{11x^2}{50} - \frac{11x}{25} + 1 \right) + \frac{c_2 \left(\frac{24497x^5}{568750} - \frac{187x^4}{1750} + \frac{17x^3}{70} - \frac{x^2}{2} + x + 1 \right)}{\sqrt[3]{x}}$$

14.54 problem 65

Internal problem ID [1345]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 65.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$8x^2(-x^2 + 2)y'' + 2x(-21x^2 + 10)y' - (35x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(8*x^2*(2-x^2)*diff(y(x),x$2)+2*x*(10-21*x^2)*diff(y(x),x)-(2+35*x^2)*y(x)=0,y(x),type
```

$$y(x) = \frac{\left(1 + \frac{1}{2}x^2 + \frac{1}{4}x^4\right) \left(x^{\frac{3}{4}}c_2 + c_1\right)}{\sqrt{x}} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 52

```
AsymptoticDSolveValue[8*x^2*(2-x^2)*y'[x]+2*x*(10-21*x^2)*y'[x]-(2+35*x^2)*y[x]==0,y[x],{x,
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(\frac{x^4}{4} + \frac{x^2}{2} + 1 \right) + \frac{c_2 \left(\frac{x^4}{4} + \frac{x^2}{2} + 1 \right)}{\sqrt{x}}$$

14.55 problem 66

Internal problem ID [1346]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 66.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 3x + 1)y'' - 4x(-3x^2 - 3x + 1)y' + 3(x^2 - x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

```
dsolve(4*x^2*(1+3*x+x^2)*diff(y(x),x$2)-4*x*(1-3*x-3*x^2)*diff(y(x),x)+3*(1-x+x^2)*y(x)=0,y(x))
```

$$y(x) = \sqrt{x} \left(x(1 - 3x + 8x^2 - 21x^3 + 55x^4 - 144x^5 + O(x^6)) c_1 \right. \\ \left. + (1 - 6x + 17x^2 - 45x^3 + 118x^4 - 309x^5 + O(x^6)) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 78

```
AsymptoticDSolveValue[4*x^2*(1+3*x+x^2)*y'[x]-4*x*(1-3*x-3*x^2)*y'[x]+3*(1-x+x^2)*y[x]==0,y[x]]
```

$$y(x) \rightarrow c_1(55x^{9/2} - 21x^{7/2} + 8x^{5/2} - 3x^{3/2} \\ + \sqrt{x}) + c_2(55x^{11/2} - 21x^{9/2} + 8x^{7/2} - 3x^{5/2} + x^{3/2})$$

14.56 problem 67

Internal problem ID [1347]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 67.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2(x+1)^2 y'' - x(-11x^2 - 10x + 1) y' + (5x^2 + 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(3*x^2*(1+x)^2*diff(y(x),x$2)-x*(1-10*x-11*x^2)*diff(y(x),x)+(1+5*x^2)*y(x)=0,y(x),typ
```

$$y(x) = (-6x^5 + 5x^4 - 4x^3 + 3x^2 - 2x + 1) \left(c_1 x^{\frac{1}{3}} + c_2 x \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 66

```
AsymptoticDSolveValue[3*x^2*(1+x)^2*y'[x]-x*(1-10*x-11*x^2)*y'[x]+(1+5*x^2)*y[x]==0,y[x],{x
```

$$y(x) \rightarrow c_1 x (-6x^5 + 5x^4 - 4x^3 + 3x^2 - 2x + 1) + c_2 \sqrt[3]{x} (-6x^5 + 5x^4 - 4x^3 + 3x^2 - 2x + 1)$$

14.57 problem 68

Internal problem ID [1348]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.5 THE METHOD OF FROBENIUS I. Exercises 7.5. Page 358

Problem number: 68.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 2x + 3)y'' - x(-15x^2 - 14x + 3)y' + (7x^2 + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(4*x^2*(3+2*x+x^2)*diff(y(x),x$2)-x*(3-14*x-15*x^2)*diff(y(x),x)+(3+7*x^2)*y(x)=0,y(x))
```

$$y(x) = \left(1 - \frac{2}{3}x + \frac{1}{9}x^2 + \frac{4}{27}x^3 - \frac{11}{81}x^4 + \frac{10}{243}x^5\right) \left(c_1x^{\frac{1}{4}} + c_2x\right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 86

```
AsymptoticDSolveValue[4*x^2*(3+2*x+x^2)*y'[x]-x*(3-14*x-15*x^2)*y'[x]+(3+7*x^2)*y[x]==0,y[x],{x,0,6}]
```

$$y(x) \rightarrow c_1x \left(\frac{10x^5}{243} - \frac{11x^4}{81} + \frac{4x^3}{27} + \frac{x^2}{9} - \frac{2x}{3} + 1 \right) + c_2\sqrt[4]{x} \left(\frac{10x^5}{243} - \frac{11x^4}{81} + \frac{4x^3}{27} + \frac{x^2}{9} - \frac{2x}{3} + 1 \right)$$

15 Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

15.1 problem Example 7.6.1 page 367	529
15.2 problem Example 7.6.2 page 369	531
15.3 problem Example 7.6.3 page 370	532
15.4 problem Example 7.6.4 page 372	533
15.5 problem 1	534
15.6 problem 2	536
15.7 problem 3	538
15.8 problem 4	540
15.9 problem 5	542
15.10 problem 6	544
15.11 problem 7	546
15.12 problem 8	548
15.13 problem 9	550
15.14 problem 10	552
15.15 problem 11	554
15.16 problem 12	556
15.17 problem 13	557
15.18 problem 14	558
15.19 problem 15	559
15.20 problem 16	560
15.21 problem 17	562
15.22 problem 18	563
15.23 problem 19	565
15.24 problem 20	566
15.25 problem 21	567
15.26 problem 22	568
15.27 problem 23	569
15.28 problem 24	571
15.29 problem 25	573
15.30 problem 26	575
15.31 problem 27	577
15.32 problem 28	578
15.33 problem 29	579
15.34 problem 30	580

15.35problem 31	581
15.36problem 32	582
15.37problem 33	583
15.38problem 34	584
15.39problem 35	585
15.40problem 36	586
15.41problem 37	587
15.42problem 38	588
15.43problem 39	589
15.44problem 40	590
15.45problem 41	591
15.46problem 42	592
15.47problem 43	593
15.48problem 44	594
15.49problem 45	595
15.50problem 46	596
15.51problem 47	597
15.52problem 48	598
15.53problem 49	599
15.54problem 50	600
15.55problem 51	601
15.56problem 52	602
15.57problem 58	603
15.58problem 59	604
15.59problem 60	605
15.60problem 61	606
15.61problem 62	607
15.62problem 63	608
15.63problem 64	609
15.64problem 65	610

15.1 problem Example 7.6.1 page 367

Internal problem ID [1349]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: Example 7.6.1 page 367.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 - 2x + 1)y'' - x(x + 3)y' + (x + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 69

Order:=6;

```
dsolve(x^2*(1-2*x+x^2)*diff(y(x),x$2)-x*(3+x)*diff(y(x),x)+(4+x)*y(x)=0,y(x),type='series',x
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 + 5x + 17x^2 + \frac{143}{3}x^3 + \frac{355}{3}x^4 + \frac{4043}{15}x^5 + O(x^6) \right) \right. \\ \left. + \left((-3)x - \frac{29}{2}x^2 - \frac{859}{18}x^3 - \frac{4693}{36}x^4 - \frac{285181}{900}x^5 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 118

```
AsymptoticDSolveValue[x^2*(1-2*x+x^2)*y'[x]-x*(3+x)*y'[x]+(4+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{4043x^5}{15} + \frac{355x^4}{3} + \frac{143x^3}{3} + 17x^2 + 5x + 1 \right) x^2$$
$$+ c_2 \left(\left(-\frac{285181x^5}{900} - \frac{4693x^4}{36} - \frac{859x^3}{18} - \frac{29x^2}{2} - 3x \right) x^2 \right.$$
$$\left. + \left(\frac{4043x^5}{15} + \frac{355x^4}{3} + \frac{143x^3}{3} + 17x^2 + 5x + 1 \right) x^2 \log(x) \right)$$

15.2 problem Example 7.6.2 page 369

Internal problem ID [1350]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: Example 7.6.2 page 369.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(2+x)y'' + 5y'x^2 + (x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(2*x^2*(2+x)*diff(y(x),x$2)+5*x^2*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - \frac{3}{4}x + \frac{15}{32}x^2 - \frac{35}{128}x^3 + \frac{315}{2048}x^4 - \frac{693}{8192}x^5 + O(x^6) \right) \right. \\ \left. + \left(\frac{1}{4}x - \frac{13}{64}x^2 + \frac{101}{768}x^3 - \frac{641}{8192}x^4 + \frac{7303}{163840}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 134

```
AsymptoticDSolveValue[2*x^2*(2+x)*y'[x]+5*x^2*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{693x^5}{8192} + \frac{315x^4}{2048} - \frac{35x^3}{128} + \frac{15x^2}{32} - \frac{3x}{4} + 1 \right) \\ + c_2 \left(\sqrt{x} \left(\frac{7303x^5}{163840} - \frac{641x^4}{8192} + \frac{101x^3}{768} - \frac{13x^2}{64} + \frac{x}{4} \right) \right. \\ \left. + \sqrt{x} \left(-\frac{693x^5}{8192} + \frac{315x^4}{2048} - \frac{35x^3}{128} + \frac{15x^2}{32} - \frac{3x}{4} + 1 \right) \log(x) \right)$$

15.3 problem Example 7.6.3 page 370

Internal problem ID [1351]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: Example 7.6.3 page 370.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-x^2 + 2)y'' - 2x(2x^2 + 1)y' + (-2x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(2-x^2)*diff(y(x),x$2)-2*x*(1+2*x^2)*diff(y(x),x)+(2-2*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 + \frac{3}{4}x^2 + \frac{15}{32}x^4 + O(x^6) \right) + \left(-\frac{1}{8}x^2 - \frac{13}{128}x^4 + O(x^6) \right) c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 65

```
AsymptoticDSolveValue[x^2*(2-x^2)*y'[x]-2*x*(1+2*x^2)*y'[x]+(2-2*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{15x^4}{32} + \frac{3x^2}{4} + 1 \right) + c_2 \left(x \left(-\frac{13x^4}{128} - \frac{x^2}{8} \right) + x \left(\frac{15x^4}{32} + \frac{3x^2}{4} + 1 \right) \log(x) \right)$$

15.4 problem Example 7.6.4 page 372

Internal problem ID [1352]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: Example 7.6.4 page 372.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(5-x)y' + (9-4x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

Order:=6;

```
dsolve(x^2*dif(y(x),x$2)-x*(5-x)*dif(y(x),x)+(9-4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) (1 + x + O(x^6)) \right. \\ \left. + \left((-3)x - \frac{1}{4}x^2 + \frac{1}{36}x^3 - \frac{1}{288}x^4 + \frac{1}{2400}x^5 + O(x^6) \right) c_2 \right) x^3$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 62

```
AsymptoticDSolveValue[x^2*y'[x]-x*(5-x)*y'[x]+(9-4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x+1)x^3 + c_2 \left((x+1)x^3 \log(x) + \left(\frac{x^5}{2400} - \frac{x^4}{288} + \frac{x^3}{36} - \frac{x^2}{4} - 3x \right) x^3 \right)$$

15.5 problem 1

Internal problem ID [1353]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(1-x)y' + (-x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 75

```
Order:=8;
```

```
dsolve(x^2*dif(y(x),x$2)-x*(1-x)*dif(y(x),x)+(1-x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - x + \frac{3}{4}x^2 - \frac{13}{36}x^3 + \frac{79}{576}x^4 - \frac{67}{1600}x^5 + \frac{5593}{518400}x^6 - \frac{60859}{25401600}x^7 + O(x^8) \right) + \left(x - x^2 + \frac{65}{108}x^3 - \frac{895}{3456}x^4 + \frac{12547}{144000}x^5 - \frac{41729}{1728000}x^6 + \frac{10121677}{1778112000}x^7 + O(x^8) \right) c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 154

```
AsymptoticDSolveValue[x^2*y''[x]-x*(1-x)*y'[x]+(1-x^2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x \left(-\frac{60859x^7}{25401600} + \frac{5593x^6}{518400} - \frac{67x^5}{1600} + \frac{79x^4}{576} - \frac{13x^3}{36} + \frac{3x^2}{4} - x + 1 \right) \\ + c_2 \left(x \left(\frac{10121677x^7}{1778112000} - \frac{41729x^6}{1728000} + \frac{12547x^5}{144000} - \frac{895x^4}{3456} + \frac{65x^3}{108} - x^2 + x \right) \right. \\ \left. + x \left(-\frac{60859x^7}{25401600} + \frac{5593x^6}{518400} - \frac{67x^5}{1600} + \frac{79x^4}{576} - \frac{13x^3}{36} + \frac{3x^2}{4} - x + 1 \right) \log(x) \right)$$

15.6 problem 2

Internal problem ID [1354]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(2x^2 + x + 1)y'' + x(7x^2 + 6x + 3)y' + (-3x^2 + 6x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 81

Order:=8;

```
dsolve(x^2*(1+x+2*x^2)*diff(y(x),x$2)+x*(3+6*x+7*x^2)*diff(y(x),x)+(1+6*x-3*x^2)*y(x)=0,y(x))
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - 2x + \frac{9}{2}x^2 - \frac{20}{3}x^3 + \frac{173}{24}x^4 - \frac{93}{20}x^5 - \frac{419}{720}x^6 + \frac{6697}{1260}x^7 + O(x^8)\right) + \left(x - \frac{15}{4}x^2 + \frac{133}{18}x^3 - \dots\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 162

AsymptoticDSolveValue[x^2*(1+x+2*x^2)*y'[x]+x*(3+6*x+7*x^2)*y'[x]+(1+6*x-3*x^2)*y[x]==0,y[x]

$$\begin{aligned}
 y(x) \rightarrow & \frac{c_1 \left(\frac{6697x^7}{1260} - \frac{419x^6}{720} - \frac{93x^5}{20} + \frac{173x^4}{24} - \frac{20x^3}{3} + \frac{9x^2}{2} - 2x + 1 \right)}{x} \\
 & + c_2 \left(\frac{-\frac{125221x^7}{29400} - \frac{70949x^6}{14400} + \frac{4217x^5}{400} - \frac{3077x^4}{288} + \frac{133x^3}{18} - \frac{15x^2}{4} + x}{x} \right. \\
 & \left. + \frac{\left(\frac{6697x^7}{1260} - \frac{419x^6}{720} - \frac{93x^5}{20} + \frac{173x^4}{24} - \frac{20x^3}{3} + \frac{9x^2}{2} - 2x + 1 \right) \log(x)}{x} \right)
 \end{aligned}$$

15.7 problem 3

Internal problem ID [1355]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$x^2(x^2 + 2x + 1)y'' + x(4x^2 + 3x + 1)y' - x(1 - 2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 71

Order:=8;

```
dsolve(x^2*(1+2*x+x^2)*diff(y(x),x$2)+x*(1+3*x+4*x^2)*diff(y(x),x)-x*(1-2*x)*y(x)=0,y(x),typ
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + x - x^2 + \frac{1}{3}x^3 + \frac{1}{3}x^4 - \frac{11}{15}x^5 + \frac{37}{45}x^6 - \frac{209}{315}x^7 + O(x^8) \right) \\ + \left((-3)x + \frac{1}{2}x^2 + \frac{31}{18}x^3 - \frac{91}{36}x^4 + \frac{1897}{900}x^5 - \frac{301}{300}x^6 - \frac{3901}{14700}x^7 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 145

```
AsymptoticDSolveValue[x^2*(1+2*x+x^2)*y'[x]+x*(1+3*x+4*x^2)*y'[x]-x*(1-2*x)*y[x]==0,y[x],{x
```

$$y(x) \rightarrow c_1 \left(-\frac{209x^7}{315} + \frac{37x^6}{45} - \frac{11x^5}{15} + \frac{x^4}{3} + \frac{x^3}{3} - x^2 + x + 1 \right) \\ + c_2 \left(-\frac{3901x^7}{14700} - \frac{301x^6}{300} + \frac{1897x^5}{900} - \frac{91x^4}{36} + \frac{31x^3}{18} + \frac{x^2}{2} \right. \\ \left. + \left(-\frac{209x^7}{315} + \frac{37x^6}{45} - \frac{11x^5}{15} + \frac{x^4}{3} + \frac{x^3}{3} - x^2 + x + 1 \right) \log(x) - 3x \right)$$

15.8 problem 4

Internal problem ID [1356]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + x + 1)y'' + 12x^2(x + 1)y' + (3x^2 + 3x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 81

```
Order:=8;
```

```
dsolve(4*x^2*(1+x+x^2)*diff(y(x),x$2)+12*x^2*(1+x)*diff(y(x),x)+(1+3*x+3*x^2)*y(x)=0,y(x),ty
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - 2x + \frac{5}{2}x^2 - 2x^3 + \frac{5}{8}x^4 + \frac{17}{20}x^5 - \frac{121}{80}x^6 + x^7 + O(x^8) \right) \right. \\ \left. + \left(x - \frac{9}{4}x^2 + \frac{17}{6}x^3 - \frac{205}{96}x^4 + \frac{481}{1200}x^5 + \frac{2109}{1600}x^6 - \frac{1063}{560}x^7 + O(x^8) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 156

```
AsymptoticDSolveValue[4*x^2*(1+x+x^2)*y'[x]+12*x^2*(1+x)*y'[x]+(1+3*x+3*x^2)*y[x]==0,y[x],{
```

$$\begin{aligned} y(x) \rightarrow & c_1 \sqrt{x} \left(x^7 - \frac{121x^6}{80} + \frac{17x^5}{20} + \frac{5x^4}{8} - 2x^3 + \frac{5x^2}{2} - 2x + 1 \right) \\ & + c_2 \left(\sqrt{x} \left(-\frac{1063x^7}{560} + \frac{2109x^6}{1600} + \frac{481x^5}{1200} - \frac{205x^4}{96} + \frac{17x^3}{6} - \frac{9x^2}{4} + x \right) \right. \\ & \left. + \sqrt{x} \left(x^7 - \frac{121x^6}{80} + \frac{17x^5}{20} + \frac{5x^4}{8} - 2x^3 + \frac{5x^2}{2} - 2x + 1 \right) \log(x) \right) \end{aligned}$$

15.9 problem 5

Internal problem ID [1357]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + x + 1)y'' - x(-2x^2 - 4x + 1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 75

Order:=8;

```
dsolve(x^2*(1+x+x^2)*diff(y(x),x$2)-x*(1-4*x-2*x^2)*diff(y(x),x)+y(x)=0,y(x),type='series',x
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - 4x + \frac{19}{2}x^2 - \frac{49}{3}x^3 + \frac{515}{24}x^4 - \frac{319}{15}x^5 + \frac{10093}{720}x^6 - \frac{647}{360}x^7 + O(x^8) \right) + \left(3x - \frac{43}{4}x^2 + \frac{208}{9}x^3 - \frac{10379}{288}x^4 + \frac{76321}{1800}x^5 - \frac{172499}{4800}x^6 + \frac{39091}{2400}x^7 + O(x^8) \right) c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 158

```
AsymptoticDSolveValue[x^2*(1+x+x^2)*y'[x]-x*(1-4*x-2*x^2)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 x \left(-\frac{647x^7}{360} + \frac{10093x^6}{720} - \frac{319x^5}{15} + \frac{515x^4}{24} - \frac{49x^3}{3} + \frac{19x^2}{2} - 4x + 1 \right) \\ & + c_2 \left(x \left(\frac{39091x^7}{2400} - \frac{172499x^6}{4800} + \frac{76321x^5}{1800} - \frac{10379x^4}{288} + \frac{208x^3}{9} - \frac{43x^2}{4} + 3x \right) \right. \\ & \left. + x \left(-\frac{647x^7}{360} + \frac{10093x^6}{720} - \frac{319x^5}{15} + \frac{515x^4}{24} - \frac{49x^3}{3} + \frac{19x^2}{2} - 4x + 1 \right) \log(x) \right) \end{aligned}$$

15.10 problem 6

Internal problem ID [1358]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' + 3x(-2x^2 + 3x + 5)y' + (-14x^2 + 12x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 81

Order:=8;

```
dsolve(9*x^2*diff(y(x),x$2)+3*x*(5+3*x-2*x^2)*diff(y(x),x)+(1+12*x-14*x^2)*y(x)=0,y(x),type=
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - x + \frac{5}{6}x^2 - \frac{1}{2}x^3 + \frac{19}{72}x^4 - \frac{43}{360}x^5 + \frac{319}{6480}x^6 - \frac{167}{9072}x^7 + O(x^8)\right) + \left(x - \frac{11}{12}x^2 + \frac{25}{36}x^3 - \frac{1}{2}x^4\right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 168

AsymptoticDSolveValue[9*x^2*y''[x]+3*x*(5+3*x-2*x^2)*y'[x]+(1+12*x-14*x^2)*y[x]==0,y[x],{x,0

$$y(x) \rightarrow \frac{c_1 \left(-\frac{167x^7}{9072} + \frac{319x^6}{6480} - \frac{43x^5}{360} + \frac{19x^4}{72} - \frac{x^3}{2} + \frac{5x^2}{6} - x + 1 \right)}{\sqrt[3]{x}} + c_2 \left(\frac{\frac{126647x^7}{3810240} - \frac{32773x^6}{388800} + \frac{4211x^5}{21600} - \frac{113x^4}{288} + \frac{25x^3}{36} - \frac{11x^2}{12} + x}{\sqrt[3]{x}} + \frac{\left(-\frac{167x^7}{9072} + \frac{319x^6}{6480} - \frac{43x^5}{360} + \frac{19x^4}{72} - \frac{x^3}{2} + \frac{5x^2}{6} - x + 1 \right) \log(x)}{\sqrt[3]{x}} \right)$$

15.11 problem 7

Internal problem ID [1359]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(x^2 + x + 1) y' + x(2 - x) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 71

```
Order:=8;
```

```
dsolve(x^2*diff(y(x),x$2)+x*(1+x*x^2)*diff(y(x),x)+x*(2-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - 2x + \frac{7}{4}x^2 - \frac{7}{9}x^3 + \frac{77}{576}x^4 + \frac{217}{7200}x^5 - \frac{8813}{518400}x^6 + \frac{143}{453600}x^7 + O(x^8) \right) + \left(3x - \frac{15}{4}x^2 + \frac{239}{108}x^3 - \frac{2021}{3456}x^4 - \frac{1241}{54000}x^5 + \frac{93859}{1728000}x^6 - \frac{311177}{42336000}x^7 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 153

```
AsymptoticDSolveValue[x^2*y'[x]+x*(1+x+x^2)*y'[x]+x*(2-x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{143x^7}{453600} - \frac{8813x^6}{518400} + \frac{217x^5}{7200} + \frac{77x^4}{576} - \frac{7x^3}{9} + \frac{7x^2}{4} - 2x + 1 \right) \\ + c_2 \left(-\frac{311177x^7}{42336000} + \frac{93859x^6}{1728000} - \frac{1241x^5}{54000} - \frac{2021x^4}{3456} + \frac{239x^3}{108} - \frac{15x^2}{4} \right. \\ \left. + \left(\frac{143x^7}{453600} - \frac{8813x^6}{518400} + \frac{217x^5}{7200} + \frac{77x^4}{576} - \frac{7x^3}{9} + \frac{7x^2}{4} - 2x + 1 \right) \log(x) + 3x \right)$$

15.12 problem 8

Internal problem ID [1360]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 + 2x)y'' + x(3x^2 + 14x + 5)y' + (12x^2 + 18x + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 79

```
Order:=8;
```

```
dsolve(x^2*(1+2*x)*diff(y(x),x$2)+x*(5+14*x+3*x^2)*diff(y(x),x)+(4+18*x+12*x^2)*y(x)=0,y(x),
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - 2x + \frac{5}{2}x^2 - 3x^3 + \frac{33}{8}x^4 - \frac{129}{20}x^5 + \frac{867}{80}x^6 - \frac{1059}{56}x^7 + O(x^8)\right) + \left(\frac{3}{4}x^2 - \frac{13}{6}x^3 + \frac{407}{96}x^4 - \frac{13}{12}x^5 + \frac{13}{16}x^6 - \frac{1}{2}x^7 + \frac{1}{24}x^8\right)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 157

AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]+x*(5+14*x+3*x^2)*y'[x]+(4+18*x+12*x^2)*y[x]==0,y[x]

$$y(x) \rightarrow \frac{c_1 \left(-\frac{1059x^7}{56} + \frac{867x^6}{80} - \frac{129x^5}{20} + \frac{33x^4}{8} - 3x^3 + \frac{5x^2}{2} - 2x + 1 \right)}{x^2} + c_2 \left(\frac{-\frac{559033x^7}{23520} + \frac{63851x^6}{4800} - \frac{9047x^5}{1200} + \frac{407x^4}{96} - \frac{13x^3}{6} + \frac{3x^2}{4}}{x^2} + \frac{\left(-\frac{1059x^7}{56} + \frac{867x^6}{80} - \frac{129x^5}{20} + \frac{33x^4}{8} - 3x^3 + \frac{5x^2}{2} - 2x + 1 \right) \log(x)}{x^2} \right)$$

15.13 problem 9

Internal problem ID [1361]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 2x(x^2 + x + 4)y' + (3x^2 + 5x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 81

```
Order:=8;
```

```
dsolve(4*x^2*diff(y(x),x$2)+2*x*(4+x+x^2)*diff(y(x),x)+(1+5*x+3*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - x + \frac{1}{4}x^2 + \frac{1}{18}x^3 - \frac{37}{1152}x^4 - \frac{17}{28800}x^5 + \frac{593}{259200}x^6 - \frac{1913}{12700800}x^7 + O(x^8)\right) + \left(\frac{3}{2}x - \frac{13}{16}x^2 - \frac{1}{\sqrt{x}}\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 172

AsymptoticDSolveValue[4*x^2*y'[x]+2*x*(4+x+x^2)*y'[x]+(1+5*x+3*x^2)*y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & \frac{c_1 \left(-\frac{1913x^7}{12700800} + \frac{593x^6}{259200} - \frac{17x^5}{28800} - \frac{37x^4}{1152} + \frac{x^3}{18} + \frac{x^2}{4} - x + 1 \right)}{\sqrt{x}} \\
 & + c_2 \left(\frac{\frac{982189x^7}{889056000} - \frac{98531x^6}{20736000} - \frac{19507x^5}{1728000} + \frac{1103x^4}{13824} + \frac{x^3}{54} - \frac{13x^2}{16} + \frac{3x}{2}}{\sqrt{x}} \right. \\
 & \left. + \frac{\left(-\frac{1913x^7}{12700800} + \frac{593x^6}{259200} - \frac{17x^5}{28800} - \frac{37x^4}{1152} + \frac{x^3}{18} + \frac{x^2}{4} - x + 1 \right) \log(x)}{\sqrt{x}} \right)
 \end{aligned}$$

15.14 problem 10

Internal problem ID [1362]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$16x^2y'' + 4x(2x^2 + x + 6)y' + (18x^2 + 5x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 81

Order:=8;

```
dsolve(16*x^2*diff(y(x),x$2)+4*x*(6+x+2*x^2)*diff(y(x),x)+(1+5*x+18*x^2)*y(x)=0,y(x),type='s'
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{1}{4}x - \frac{7}{32}x^2 + \frac{23}{384}x^3 + \frac{145}{6144}x^4 - \frac{881}{122880}x^5 - \frac{4919}{2949120}x^6 + \frac{47207}{82575360}x^7 + O(x^8)\right) + \left(\frac{1}{4}x + \frac{1}{6}\right)}{x^{\frac{1}{4}}}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 176

AsymptoticDSolveValue[16*x^2*y''[x]+4*x*(6+x+2*x^2)*y'[x]+(1+5*x+18*x^2)*y[x]==0,y[x],{x,0,7}

$$y(x) \rightarrow \frac{c_1 \left(\frac{47207x^7}{82575360} - \frac{4919x^6}{2949120} - \frac{881x^5}{122880} + \frac{145x^4}{6144} + \frac{23x^3}{384} - \frac{7x^2}{32} - \frac{x}{4} + 1 \right)}{\sqrt[4]{x}} + c_2 \left(\frac{-\frac{953509x^7}{1284505600} + \frac{50791x^6}{58982400} + \frac{65017x^5}{7372800} - \frac{841x^4}{73728} - \frac{157x^3}{2304} + \frac{5x^2}{64} + \frac{x}{4}}{\sqrt[4]{x}} + \frac{\left(\frac{47207x^7}{82575360} - \frac{4919x^6}{2949120} - \frac{881x^5}{122880} + \frac{145x^4}{6144} + \frac{23x^3}{384} - \frac{7x^2}{32} - \frac{x}{4} + 1 \right) \log(x)}{\sqrt[4]{x}} \right)$$

15.15 problem 11

Internal problem ID [1363]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2(x+1)y'' + 3x(-x^2 + 11x + 5)y' + (-7x^2 + 16x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 79

Order:=8;

```
dsolve(9*x^2*(1+x)*diff(y(x),x$2)+3*x*(5+11*x-x^2)*diff(y(x),x)+(1+16*x-7*x^2)*y(x)=0,y(x),t
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - x + \frac{7}{6}x^2 - \frac{23}{18}x^3 + \frac{11}{8}x^4 - \frac{1577}{1080}x^5 + \frac{3319}{2160}x^6 - \frac{72853}{45360}x^7 + O(x^8)\right) + \left(-\frac{1}{12}x^2 + \frac{13}{108}x^3 - \dots\right)}{x^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 167

AsymptoticDSolveValue[9*x^2*(1+x)*y'[x]+3*x*(5+11*x-x^2)*y'[x]+(1+16*x-7*x^2)*y[x]==0,y[x],

$$y(x) \rightarrow \frac{c_1 \left(-\frac{72853x^7}{45360} + \frac{3319x^6}{2160} - \frac{1577x^5}{1080} + \frac{11x^4}{8} - \frac{23x^3}{18} + \frac{7x^2}{6} - x + 1 \right)}{\sqrt[3]{x}} + c_2 \left(\frac{\frac{4118557x^7}{19051200} - \frac{76919x^6}{388800} + \frac{11449x^5}{64800} - \frac{131x^4}{864} + \frac{13x^3}{108} - \frac{x^2}{12}}{\sqrt[3]{x}} + \frac{\left(-\frac{72853x^7}{45360} + \frac{3319x^6}{2160} - \frac{1577x^5}{1080} + \frac{11x^4}{8} - \frac{23x^3}{18} + \frac{7x^2}{6} - x + 1 \right) \log(x)}{\sqrt[3]{x}} \right)$$

15.16 problem 12

Internal problem ID [1364]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + (4x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(4*x^2*diff(y(x),x$2)+(1+4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - x + \frac{1}{4}x^2 - \frac{1}{36}x^3 + \frac{1}{576}x^4 - \frac{1}{14400}x^5 + O(x^6) \right) \right. \\ \left. + \left(2x - \frac{3}{4}x^2 + \frac{11}{108}x^3 - \frac{25}{3456}x^4 + \frac{137}{432000}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 128

```
AsymptoticDSolveValue[4*x^2*y''[x]+(1+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right) \\ + c_2 \left(\sqrt{x} \left(\frac{137x^5}{432000} - \frac{25x^4}{3456} + \frac{11x^3}{108} - \frac{3x^2}{4} + 2x \right) \right. \\ \left. + \sqrt{x} \left(-\frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1 \right) \log(x) \right)$$

15.17 problem 13

Internal problem ID [1365]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$36x^2(1 - 2x)y'' + 24x(1 - 9x)y' + (1 - 70x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 69

Order:=6;

`dsolve(36*x^2*(1-2*x)*diff(y(x),x$2)+24*x*(1-9*x)*diff(y(x),x)+(1-70*x)*y(x)=0,y(x),type='series')`

$$y(x) = x^{\frac{1}{6}} \left((c_2 \ln(x) + c_1) \left(1 + \frac{8}{3}x + \frac{56}{9}x^2 + \frac{1120}{81}x^3 + \frac{7280}{243}x^4 + \frac{46592}{729}x^5 + O(x^6) \right) + \left(-\frac{2}{3}x - 2x^2 - \frac{1192}{243}x^3 - \frac{8168}{729}x^4 - \frac{270112}{10935}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 132

`AsymptoticDSolveValue[36*x^2*(1-2*x)*y'[x]+24*x*(1-9*x)*y'[x]+(1-70*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_1 \sqrt[6]{x} \left(\frac{46592x^5}{729} + \frac{7280x^4}{243} + \frac{1120x^3}{81} + \frac{56x^2}{9} + \frac{8x}{3} + 1 \right) + c_2 \left(\sqrt[6]{x} \left(-\frac{270112x^5}{10935} - \frac{8168x^4}{729} - \frac{1192x^3}{243} - 2x^2 - \frac{2x}{3} \right) + \sqrt[6]{x} \left(\frac{46592x^5}{729} + \frac{7280x^4}{243} + \frac{1120x^3}{81} + \frac{56x^2}{9} + \frac{8x}{3} + 1 \right) \log(x) \right)$$

15.18 problem 14

Internal problem ID [1366]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' - x(3-x)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(x^2*(1+x)*diff(y(x),x$2)-x*(3-x)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x^2((c_2 \ln(x) + c_1)(1 - 4x + 9x^2 - 16x^3 + 25x^4 - 36x^5 + O(x^6)) + (4x - 12x^2 + 24x^3 - 40x^4 + 60x^5 + O(x^6))c_2)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 98

```
AsymptoticDSolveValue[x^2*(1+x)*y''[x]-x*(3-x)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(-36x^5 + 25x^4 - 16x^3 + 9x^2 - 4x + 1)x^2 + c_2((60x^5 - 40x^4 + 24x^3 - 12x^2 + 4x)x^2 + (-36x^5 + 25x^4 - 16x^3 + 9x^2 - 4x + 1)x^2 \log(x))$$

15.19 problem 15

Internal problem ID [1367]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 - 2x)y'' - x(5 - 4x)y' + (9 - 4x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

Order:=6;

```
dsolve(x^2*(1-2*x)*diff(y(x),x$2)-x*(5-4*x)*diff(y(x),x)+(9-4*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = x^3((c_2 \ln(x) + c_1)(1 + 4x + 12x^2 + 32x^3 + 80x^4 + 192x^5 + O(x^6)) + ((-2)x - 8x^2 - 24x^3 - 64x^4 - 160x^5 + O(x^6))c_2)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 98

```
AsymptoticDSolveValue[x^2*(1-2*x)*y'[x]-x*(5-4*x)*y'[x]+(9-4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(192x^5 + 80x^4 + 32x^3 + 12x^2 + 4x + 1)x^3 + c_2((-160x^5 - 64x^4 - 24x^3 - 8x^2 - 2x)x^3 + (192x^5 + 80x^4 + 32x^3 + 12x^2 + 4x + 1)x^3 \log(x))$$

15.20 problem 16

Internal problem ID [1368]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$25x^2y'' + x(15 + x)y' + (x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(25*x^2*diff(y(x),x$2)+x*(15+x)*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x^{\frac{1}{5}} \left((c_2 \ln(x) + c_1) \left(1 - \frac{6}{125}x + \frac{33}{31250}x^2 - \frac{88}{5859375}x^3 + \frac{77}{488281250}x^4 - \frac{1001}{762939453125}x^5 + O(x^6) \right) + \left(\frac{7}{125}x - \frac{113}{62500}x^2 + \frac{1091}{35156250}x^3 - \frac{1721}{4687500000}x^4 + \frac{609221}{183105468750000}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 134

```
AsymptoticDSolveValue[25*x^2*y''[x]+x*(15+x)*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \sqrt[5]{x} \left(-\frac{1001x^5}{762939453125} + \frac{77x^4}{488281250} - \frac{88x^3}{5859375} + \frac{33x^2}{31250} - \frac{6x}{125} + 1 \right) \\ & + c_2 \left(\sqrt[5]{x} \left(\frac{609221x^5}{183105468750000} - \frac{1721x^4}{4687500000} + \frac{1091x^3}{35156250} - \frac{113x^2}{62500} + \frac{7x}{125} \right) \right. \\ & \left. + \sqrt[5]{x} \left(-\frac{1001x^5}{762939453125} + \frac{77x^4}{488281250} - \frac{88x^3}{5859375} + \frac{33x^2}{31250} - \frac{6x}{125} + 1 \right) \log(x) \right) \end{aligned}$$

15.21 problem 17

Internal problem ID [1369]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(2+x)y'' + y'x^2 + (1-x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(2*x^2*(2+x)*diff(y(x),x$2)+x^2*diff(y(x),x)+(1-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 + \frac{1}{4}x - \frac{1}{32}x^2 + \frac{1}{128}x^3 - \frac{5}{2048}x^4 + \frac{7}{8192}x^5 + O(x^6) \right) \right. \\ \left. + \left(-\frac{3}{4}x + \frac{3}{64}x^2 - \frac{7}{768}x^3 + \frac{61}{24576}x^4 - \frac{391}{491520}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 134

```
AsymptoticDSolveValue[2*x^2*(2+x)*y'[x]+x^2*y''[x]+(1-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{7x^5}{8192} - \frac{5x^4}{2048} + \frac{x^3}{128} - \frac{x^2}{32} + \frac{x}{4} + 1 \right) \\ + c_2 \left(\sqrt{x} \left(-\frac{391x^5}{491520} + \frac{61x^4}{24576} - \frac{7x^3}{768} + \frac{3x^2}{64} - \frac{3x}{4} \right) \right. \\ \left. + \sqrt{x} \left(\frac{7x^5}{8192} - \frac{5x^4}{2048} + \frac{x^3}{128} - \frac{x^2}{32} + \frac{x}{4} + 1 \right) \log(x) \right)$$

15.22 problem 18

Internal problem ID [1370]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(4x + 9)y'' + 3y'x + (x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(x^2*(9+4*x)*diff(y(x),x$2)+3*x*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x^{\frac{1}{3}} \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{81}x + \frac{25}{26244}x^2 - \frac{3025}{19131876}x^3 + \frac{874225}{24794911296}x^4 - \frac{18498601}{2008387814976}x^5 + O(x^6) \right) + \left(\frac{14}{81}x - \frac{35}{2916}x^2 + \frac{110495}{57395628}x^3 - \frac{62786185}{148769467776}x^4 + \frac{1315043653}{12050326889856}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 134

```
AsymptoticDSolveValue[x^2*(9+4*x)*y'[x]+3*x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \sqrt[3]{x} \left(-\frac{18498601x^5}{2008387814976} + \frac{874225x^4}{24794911296} - \frac{3025x^3}{19131876} + \frac{25x^2}{26244} - \frac{x}{81} + 1 \right) \\ & + c_2 \left(\sqrt[3]{x} \left(\frac{1315043653x^5}{12050326889856} - \frac{62786185x^4}{148769467776} + \frac{110495x^3}{57395628} - \frac{35x^2}{2916} + \frac{14x}{81} \right) \right. \\ & \left. + \sqrt[3]{x} \left(-\frac{18498601x^5}{2008387814976} + \frac{874225x^4}{24794911296} - \frac{3025x^3}{19131876} + \frac{25x^2}{26244} - \frac{x}{81} + 1 \right) \log(x) \right) \end{aligned}$$

15.23 problem 19

Internal problem ID [1371]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(-2x + 3)y' + (3x + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)-x*(3-2*x)*diff(y(x),x)+(4+3*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - 7x + \frac{63}{4}x^2 - \frac{77}{4}x^3 + \frac{1001}{64}x^4 - \frac{3003}{320}x^5 + O(x^6) \right) + \left(12x - \frac{157}{4}x^2 + \frac{2063}{36}x^3 - \frac{59875}{1152}x^4 + \frac{323399}{9600}x^5 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 122

```
AsymptoticDSolveValue[x^2*y''[x]-x*(3-2*x)*y'[x]+(4+3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{3003x^5}{320} + \frac{1001x^4}{64} - \frac{77x^3}{4} + \frac{63x^2}{4} - 7x + 1 \right) x^2 + c_2 \left(\left(\frac{323399x^5}{9600} - \frac{59875x^4}{1152} + \frac{2063x^3}{36} - \frac{157x^2}{4} + 12x \right) x^2 + \left(-\frac{3003x^5}{320} + \frac{1001x^4}{64} - \frac{77x^3}{4} + \frac{63x^2}{4} - 7x + 1 \right) x^2 \log(x) \right)$$

15.24 problem 20

Internal problem ID [1372]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(-4x + 1)y'' + 3x(1 - 6x)y' + (1 - 12x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 69

Order:=6;

```
dsolve(x^2*(1-4*x)*diff(y(x),x$2)+3*x*(1-6*x)*diff(y(x),x)+(1-12*x)*y(x)=0,y(x),type='series
```

$y(x)$

$$= \frac{(c_2 \ln(x) + c_1)(1 + 2x + 6x^2 + 20x^3 + 70x^4 + 252x^5 + O(x^6)) + (2x + 7x^2 + \frac{74}{3}x^3 + \frac{533}{6}x^4 + \frac{1627}{5}x^5 + \dots}{x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 104

```
AsymptoticDSolveValue[x^2*(1-4*x)*y'[x]+3*x*(1-6*x)*y'[x]+(1-12*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1(252x^5 + 70x^4 + 20x^3 + 6x^2 + 2x + 1)}{x} + c_2 \left(\frac{\frac{1627x^5}{5} + \frac{533x^4}{6} + \frac{74x^3}{3} + 7x^2 + 2x}{x} + \frac{(252x^5 + 70x^4 + 20x^3 + 6x^2 + 2x + 1) \log(x)}{x} \right)$$

15.25 problem 21

Internal problem ID [1373]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(1 + 2x)y'' + x(3 + 5x)y' + (1 - 2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 69

Order:=6;

`dsolve(x^2*(1+2*x)*diff(y(x),x$2)+x*(3+5*x)*diff(y(x),x)+(1-2*x)*y(x)=0,y(x),type='series',x`

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 + 3x + \frac{3}{2}x^2 - \frac{1}{2}x^3 + \frac{3}{8}x^4 - \frac{3}{8}x^5 + O(x^6)\right) + ((-5)x - \frac{25}{4}x^2 + \frac{5}{4}x^3 - \frac{25}{32}x^4 + \frac{113}{160}x^5 + C)}{x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 122

`AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]+x*(3+5*x)*y'[x]+(1-2*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow \frac{c_1 \left(-\frac{3x^5}{8} + \frac{3x^4}{8} - \frac{x^3}{2} + \frac{3x^2}{2} + 3x + 1 \right)}{x} + c_2 \left(\frac{\frac{113x^5}{160} - \frac{25x^4}{32} + \frac{5x^3}{4} - \frac{25x^2}{4} - 5x}{x} + \frac{\left(-\frac{3x^5}{8} + \frac{3x^4}{8} - \frac{x^3}{2} + \frac{3x^2}{2} + 3x + 1 \right) \log(x)}{x} \right)$$

15.26 problem 22

Internal problem ID [1374]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x+1)y'' - x(6-x)y' + (8-x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 69

Order:=6;

```
dsolve(2*x^2*(1+x)*diff(y(x),x$2)-x*(6-x)*diff(y(x),x)+(8-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - \frac{5}{2}x + \frac{35}{8}x^2 - \frac{105}{16}x^3 + \frac{1155}{128}x^4 - \frac{3003}{256}x^5 + O(x^6) \right) + \left(\frac{3}{2}x - \frac{57}{16}x^2 + \frac{583}{96}x^3 - \frac{13771}{1536}x^4 + \frac{187339}{15360}x^5 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 128

```
AsymptoticDSolveValue[2*x^2*(1+x)*y'[x]-x*(6-x)*y'[x]+(8-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{3003x^5}{256} + \frac{1155x^4}{128} - \frac{105x^3}{16} + \frac{35x^2}{8} - \frac{5x}{2} + 1 \right) x^2 + c_2 \left(\left(\frac{187339x^5}{15360} - \frac{13771x^4}{1536} + \frac{583x^3}{96} - \frac{57x^2}{16} + \frac{3x}{2} \right) x^2 + \left(-\frac{3003x^5}{256} + \frac{1155x^4}{128} - \frac{105x^3}{16} + \frac{35x^2}{8} - \frac{5x}{2} + 1 \right) x^2 \log(x) \right)$$

15.27 problem 23

Internal problem ID [1375]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 + 2x)y'' + x(5 + 9x)y' + (3x + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 81

```
Order:=8;
```

```
dsolve(x^2*(1+2*x)*diff(y(x),x$2)+x*(5+9*x)*diff(y(x),x)+(4+3*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 + 3x + \frac{3}{2}x^2 - \frac{1}{2}x^3 + \frac{3}{8}x^4 - \frac{3}{8}x^5 + \frac{7}{16}x^6 - \frac{9}{16}x^7 + O(x^8)\right) + \left((-5)x - \frac{25}{4}x^2 + \frac{5}{4}x^3 - \frac{25}{32}x^4\right)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 164

AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]+x*(5+9*x)*y'[x]+(4+3*x)*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow \frac{c_1 \left(-\frac{9x^7}{16} + \frac{7x^6}{16} - \frac{3x^5}{8} + \frac{3x^4}{8} - \frac{x^3}{2} + \frac{3x^2}{2} + 3x + 1 \right)}{x^2} + c_2 \left(\frac{\frac{2123x^7}{2240} - \frac{247x^6}{320} + \frac{113x^5}{160} - \frac{25x^4}{32} + \frac{5x^3}{4} - \frac{25x^2}{4} - 5x}{x^2} + \frac{\left(-\frac{9x^7}{16} + \frac{7x^6}{16} - \frac{3x^5}{8} + \frac{3x^4}{8} - \frac{x^3}{2} + \frac{3x^2}{2} + 3x + 1 \right) \log(x)}{x^2} \right)$$

15.28 problem 24

Internal problem ID [1376]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 - 2x)y'' - x(4x + 5)y' + (4x + 9)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 81

Order:=8;

```
dsolve(x^2*(1-2*x)*diff(y(x),x$2)-x*(5+4*x)*diff(y(x),x)+(9+4*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = \left((c_2 \ln(x) + c_1) (1 + 20x + 180x^2 + 1120x^3 + 5600x^4 + 24192x^5 + 94080x^6 + 337920x^7 + O(x^8)) + \left((-26)x - 324x^2 - \frac{6968}{3}x^3 - \frac{37780}{3}x^4 - 57360x^5 - \frac{694736}{3}x^6 - \frac{2566144}{3}x^7 + O(x^8) \right) c_2 \right) x^3$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 136

```
AsymptoticDSolveValue[x^2*(1-2*x)*y'[x]-x*(5+4*x)*y'[x]+(9+4*x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1(337920x^7 + 94080x^6 + 24192x^5 + 5600x^4 + 1120x^3 + 180x^2 + 20x + 1)x^3 + c_2\left(\left(-\frac{2566144x^7}{3} - \frac{694736x^6}{3} - 57360x^5 - \frac{37780x^4}{3} - \frac{6968x^3}{3} - 324x^2 - 26x\right)x^3 + (337920x^7 + 94080x^6 + 24192x^5 + 5600x^4 + 1120x^3 + 180x^2 + 20x + 1)x^3 \log(x)\right)$$

15.29 problem 25

Internal problem ID [1377]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(4x + 1)y'' - x(-4x + 1)y' + (x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 75

Order:=8;

```
dsolve(x^2*(1+4*x)*diff(y(x),x$2)-x*(1-4*x)*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - 5x + \frac{85}{4}x^2 - \frac{3145}{36}x^3 + \frac{204425}{576}x^4 - \frac{825877}{576}x^5 + \frac{119752165}{20736}x^6 - \frac{23591176505}{1016064}x^7 + O(x^8) \right) + \left(2x - \frac{39}{4}x^2 + \frac{4499}{108}x^3 - \frac{594305}{3456}x^4 + \frac{2420617}{3456}x^5 - \frac{117547073}{41472}x^6 + \frac{162576422327}{14224896}x^7 + O(x^8) \right) c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 158

```
AsymptoticDSolveValue[x^2*(1+4*x)*y'[x]-x*(1-4*x)*y'[x]+(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x \left(-\frac{23591176505x^7}{1016064} + \frac{119752165x^6}{20736} - \frac{825877x^5}{576} + \frac{204425x^4}{576} - \frac{3145x^3}{36} + \frac{85x^2}{4} - 5x + 1 \right) + c_2 \left(x \left(\frac{162576422327x^7}{14224896} - \frac{117547073x^6}{41472} + \frac{2420617x^5}{3456} - \frac{594305x^4}{3456} + \frac{4499x^3}{108} - \frac{39x^2}{4} + 2x \right) + x \left(-\frac{23591176505x^7}{1016064} + \frac{119752165x^6}{20736} - \frac{825877x^5}{576} + \frac{204425x^4}{576} - \frac{3145x^3}{36} + \frac{85x^2}{4} - 5x + 1 \right) \log(x) \right)$$

15.30 problem 26

Internal problem ID [1378]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' + x(1+2x)y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 71

`Order:=8;`

```
dsolve(x^2*(1+x)*diff(y(x),x$2)+x*(1+2*x)*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - x + \frac{3}{4}x^2 - \frac{7}{12}x^3 + \frac{91}{192}x^4 - \frac{637}{1600}x^5 + \frac{19747}{57600}x^6 - \frac{17329}{57600}x^7 + O(x^8) \right) \\ + \left(x - \frac{3}{4}x^2 + \frac{5}{9}x^3 - \frac{499}{1152}x^4 + \frac{16919}{48000}x^5 - \frac{56861}{192000}x^6 + \frac{1027717}{4032000}x^7 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 151

```
AsymptoticDSolveValue[x^2*(1+x)*y'[x]+x*(1+2*x)*y'[x]+x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{17329x^7}{57600} + \frac{19747x^6}{57600} - \frac{637x^5}{1600} + \frac{91x^4}{192} - \frac{7x^3}{12} + \frac{3x^2}{4} - x + 1 \right) \\ + c_2 \left(\frac{1027717x^7}{4032000} - \frac{56861x^6}{192000} + \frac{16919x^5}{48000} - \frac{499x^4}{1152} + \frac{5x^3}{9} - \frac{3x^2}{4} \right. \\ \left. + \left(-\frac{17329x^7}{57600} + \frac{19747x^6}{57600} - \frac{637x^5}{1600} + \frac{91x^4}{192} - \frac{7x^3}{12} + \frac{3x^2}{4} - x + 1 \right) \log(x) + x \right)$$

15.31 problem 27

Internal problem ID [1379]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1-x)y'' + x(7+x)y' + (9-x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 63

Order:=8;

```
dsolve(x^2*(1-x)*diff(y(x),x$2)+x*(7+x)*diff(y(x),x)+(9-x)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{(c_2 \ln(x) + c_1)(1 + 16x + 36x^2 + 16x^3 + x^4 + O(x^8)) + ((-40)x - 150x^2 - \frac{280}{3}x^3 - \frac{25}{3}x^4 + O(x^8))c_2}{x^3}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 83

```
AsymptoticDSolveValue[x^2*(1-x)*y'[x]+x*(7+x)*y'[x]+(9-x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{c_1(x^4 + 16x^3 + 36x^2 + 16x + 1)}{x^3} + c_2 \left(\frac{-\frac{25x^4}{3} - \frac{280x^3}{3} - 150x^2 - 40x}{x^3} + \frac{(x^4 + 16x^3 + 36x^2 + 16x + 1) \log(x)}{x^3} \right)$$

15.32 problem 28

Internal problem ID [1380]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(-x^2 + 1) y' + y(x^2 + 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*dif(y(x),x$2)-x*(1-x^2)*dif(y(x),x)+(1+x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 + O(x^6) \right) + \left(\frac{1}{4}x^2 - \frac{3}{32}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 65

```
AsymptoticDSolveValue[x^2*y'[x]-x*(1-x^2)*y'[x]+(1+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right) + c_2 \left(x \left(\frac{x^2}{4} - \frac{3x^4}{32} \right) + x \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right) \log(x) \right)$$

15.33 problem 29

Internal problem ID [1381]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - 3x(-x^2 + 1)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-3*x*(1-x^2)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) (1 - 2x^2 + 3x^4 + O(x^6)) + \left(\frac{1}{2}x^2 - x^4 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 61

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]-3*x*(1-x^2)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(3x^4 - 2x^2 + 1)x^2 + c_2 \left(\left(\frac{x^2}{2} - x^4 \right) x^2 + (3x^4 - 2x^2 + 1)x^2 \log(x) \right)$$

15.34 problem 30

Internal problem ID [1382]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 2x^3y' + (3x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

Order:=6;

```
dsolve(4*x^2*diff(y(x),x$2)+2*x^3*diff(y(x),x)+(1+3*x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 + O(x^6) \right) + \left(\frac{1}{8}x^2 - \frac{3}{128}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 77

```
AsymptoticDSolveValue[4*x^2*y'[x]+2*x^3*y''[x]+(1+3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^4}{32} - \frac{x^2}{4} + 1 \right) + c_2 \left(\sqrt{x} \left(\frac{x^2}{8} - \frac{3x^4}{128} \right) + \sqrt{x} \left(\frac{x^4}{32} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

15.35 problem 31

Internal problem ID [1383]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - x(-2x^2 + 1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-x*(1-2*x^2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{2}x^2 + \frac{3}{8}x^4 + O(x^6) \right) + \left(-\frac{1}{4}x^2 + \frac{7}{32}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 65

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]-x*(1-2*x^2)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{3x^4}{8} - \frac{x^2}{2} + 1 \right) + c_2 \left(x \left(\frac{7x^4}{32} - \frac{x^2}{4} \right) + x \left(\frac{3x^4}{8} - \frac{x^2}{2} + 1 \right) \log(x) \right)$$

15.36 problem 32

Internal problem ID [1384]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(x^2 + 2)y'' + 7x^3y' + (3x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(2*x^2*(2+x^2)*diff(y(x),x$2)+7*x^3*diff(y(x),x)+(1+3*x^2)*y(x)=0,y(x),type='series',x
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - \frac{3}{8}x^2 + \frac{21}{128}x^4 + O(x^6) \right) + \left(-\frac{1}{16}x^2 + \frac{17}{512}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 77

```
AsymptoticDSolveValue[2*x^2*(2+x^2)*y'[x]+7*x^3*y'[x]+(1+3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{21x^4}{128} - \frac{3x^2}{8} + 1 \right) + c_2 \left(\sqrt{x} \left(\frac{17x^4}{512} - \frac{x^2}{16} \right) + \sqrt{x} \left(\frac{21x^4}{128} - \frac{3x^2}{8} + 1 \right) \log(x) \right)$$

15.37 problem 33

Internal problem ID [1385]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - x(-4x^2 + 1)y' + y(2x^2 + 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-x*(1-4*x^2)*diff(y(x),x)+(1+2*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = x \left((c_2 \ln(x) + c_1) \left(1 - \frac{3}{2}x^2 + \frac{15}{8}x^4 + O(x^6) \right) + \left(\frac{1}{4}x^2 - \frac{13}{32}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 65

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]-x*(1-4*x^2)*y'[x]+(1+2*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{15x^4}{8} - \frac{3x^2}{2} + 1 \right) + c_2 \left(x \left(\frac{x^2}{4} - \frac{13x^4}{32} \right) + x \left(\frac{15x^4}{8} - \frac{3x^2}{2} + 1 \right) \log(x) \right)$$

15.38 problem 34

Internal problem ID [1386]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 4)y'' + 3x(3x^2 + 8)y' + (-9x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(4*x^2*(4+x^2)*diff(y(x),x$2)+3*x*(8+3*x^2)*diff(y(x),x)+(1-9*x^2)*y(x)=0,y(x),type='s
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 + \frac{5}{32}x^2 - \frac{15}{2048}x^4 + O(x^6)\right) + \left(-\frac{13}{64}x^2 + \frac{13}{8192}x^4 + O(x^6)\right) c_2}{x^{\frac{1}{4}}}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 77

```
AsymptoticDSolveValue[4*x^2*(4+x^2)*y'[x]+3*x*(8+3*x^2)*y'[x]+(1-9*x^2)*y[x]==0,y[x],{x,0,5
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{15x^4}{2048} + \frac{5x^2}{32} + 1\right)}{\sqrt[4]{x}} + c_2 \left(\frac{\frac{13x^4}{8192} - \frac{13x^2}{64}}{\sqrt[4]{x}} + \frac{\left(-\frac{15x^4}{2048} + \frac{5x^2}{32} + 1\right) \log(x)}{\sqrt[4]{x}}\right)$$

15.39 problem 35

Internal problem ID [1387]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2(x^2 + 3)y'' + x(11x^2 + 3)y' + (5x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(3*x^2*(3+x^2)*diff(y(x),x$2)+x*(3+11*x^2)*diff(y(x),x)+(1+5*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = x^{\frac{1}{3}} \left((c_2 \ln(x) + c_1) \left(1 - \frac{2}{9}x^2 + \frac{5}{81}x^4 + O(x^6) \right) + \left(-\frac{1}{18}x^2 + \frac{1}{54}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
AsymptoticDSolveValue[3*x^2*(3+x^2)*y'[x]+x*(3+11*x^2)*y'[x]+(1+5*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{1139x^5}{405} + \frac{53x^4}{81} + \frac{7x^3}{9} - \frac{11x^2}{9} + x + 1 \right)}{\sqrt[3]{x}}$$

15.40 problem 36

Internal problem ID [1388]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(4x^2 + 1)y'' + 32x^3y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

Order:=6;

```
dsolve(4*x^2*(1+4*x^2)*diff(y(x),x$2)+32*x^3*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 - \frac{3}{4}x^2 + \frac{105}{64}x^4 + O(x^6) \right) + \left(-\frac{5}{4}x^2 + \frac{389}{128}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 77

```
AsymptoticDSolveValue[4*x^2*(1+4*x^2)*y'[x]+32*x^3*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{105x^4}{64} - \frac{3x^2}{4} + 1 \right) + c_2 \left(\sqrt{x} \left(\frac{389x^4}{128} - \frac{5x^2}{4} \right) + \sqrt{x} \left(\frac{105x^4}{64} - \frac{3x^2}{4} + 1 \right) \log(x) \right)$$

15.41 problem 37

Internal problem ID [1389]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' - 3x(-2x^2 + 7)y' + (2x^2 + 25)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(9*x^2*diff(y(x),x$2)-3*x*(7-2*x^2)*diff(y(x),x)+(25+2*x^2)*y(x)=0,y(x),type='series',
```

$$y(x) = x^{\frac{5}{3}} \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{3}x^2 + \frac{1}{18}x^4 + O(x^6) \right) + \left(\frac{1}{6}x^2 - \frac{1}{24}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 77

```
AsymptoticDSolveValue[9*x^2*y'[x]-3*x*(7-2*x^2)*y'[x]+(25+2*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{18} - \frac{x^2}{3} + 1 \right) x^{5/3} + c_2 \left(\left(\frac{x^2}{6} - \frac{x^4}{24} \right) x^{5/3} + \left(\frac{x^4}{18} - \frac{x^2}{3} + 1 \right) x^{5/3} \log(x) \right)$$

15.42 problem 38

Internal problem ID [1390]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(2x^2 + 1)y'' + x(7x^2 + 3)y' + (-3x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(1+2*x^2)*diff(y(x),x$2)+x*(3+7*x^2)*diff(y(x),x)+(1-3*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 + \frac{3}{2}x^2 - \frac{3}{8}x^4 + O(x^6)\right) + \left(-\frac{7}{4}x^2 - \frac{7}{32}x^4 + O(x^6)\right) c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(1+2*x^2)*y''[x]+x*(3+7*x^2)*y'[x]+(1-3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{3x^4}{8} + \frac{3x^2}{2} + 1\right)}{x} + c_2 \left(\frac{-\frac{7x^4}{32} - \frac{7x^2}{4}}{x} + \frac{\left(-\frac{3x^4}{8} + \frac{3x^2}{2} + 1\right) \log(x)}{x}\right)$$

15.43 problem 39

Internal problem ID [1391]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(x^2 + 1)y'' + x(8x^2 + 3)y' + (12x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*(3+8*x^2)*diff(y(x),x)+(1+12*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{3}{2}x^2 + \frac{15}{8}x^4 + O(x^6)\right) + \left(\frac{1}{4}x^2 - \frac{13}{32}x^4 + O(x^6)\right) c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]+x*(3+8*x^2)*y'[x]+(1+12*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(\frac{15x^4}{8} - \frac{3x^2}{2} + 1\right)}{x} + c_2 \left(\frac{\frac{x^2}{4} - \frac{13x^4}{32}}{x} + \frac{\left(\frac{15x^4}{8} - \frac{3x^2}{2} + 1\right) \log(x)}{x}\right)$$

15.44 problem 40

Internal problem ID [1392]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(-x^2 + 1) y' + y(x^2 + 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*dif(y(x),x$2)-x*(1-x^2)*dif(y(x),x)+(1+x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 + O(x^6) \right) + \left(\frac{1}{4}x^2 - \frac{3}{32}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 65

```
AsymptoticDSolveValue[x^2*y'[x]-x*(1-x^2)*y'[x]+(1+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right) + c_2 \left(x \left(\frac{x^2}{4} - \frac{3x^4}{32} \right) + x \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right) \log(x) \right)$$

15.45 problem 41

Internal problem ID [1393]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-2x^2 + 1)y'' + x(-9x^2 + 5)y' + (-3x^2 + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(1-2*x^2)*diff(y(x),x$2)+x*(5-9*x^2)*diff(y(x),x)+(4-3*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{3}{4}x^2 - \frac{9}{64}x^4 + O(x^6)\right) + \left(\frac{1}{2}x^2 - \frac{21}{128}x^4 + O(x^6)\right) c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(1-2*x^2)*y''[x]+x*(5-9*x^2)*y'[x]+(4-3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{9x^4}{64} - \frac{3x^2}{4} + 1\right)}{x^2} + c_2 \left(\frac{\frac{x^2}{2} - \frac{21x^4}{128}}{x^2} + \frac{\left(-\frac{9x^4}{64} - \frac{3x^2}{4} + 1\right) \log(x)}{x^2}\right)$$

15.46 problem 42

Internal problem ID [1394]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 2)y'' + x(-x^2 + 14)y' + 2(x^2 + 9)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(2+x^2)*diff(y(x),x$2)+x*(14-x^2)*diff(y(x),x)+2*(9+x^2)*y(x)=0,y(x),type='series
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{17}{8}x^2 + \frac{85}{256}x^4 + O(x^6)\right) + \left(\frac{25}{8}x^2 - \frac{471}{512}x^4 + O(x^6)\right) c_2}{x^3}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(2+x^2)*y''[x]+x*(14-x^2)*y'[x]+2*(9+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(\frac{85x^4}{256} - \frac{17x^2}{8} + 1\right)}{x^3} + c_2 \left(\frac{\frac{25x^2}{8} - \frac{471x^4}{512}}{x^3} + \frac{\left(\frac{85x^4}{256} - \frac{17x^2}{8} + 1\right) \log(x)}{x^3}\right)$$

15.47 problem 43

Internal problem ID [1395]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' + x(7x^2 + 3)y' + (8x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*(3+7*x^2)*diff(y(x),x)+(1+8*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{3}{4}x^2 + \frac{45}{64}x^4 + O(x^6)\right) + \left(-\frac{1}{4}x^2 + \frac{33}{128}x^4 + O(x^6)\right) c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]+x*(3+7*x^2)*y'[x]+(1+8*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(\frac{45x^4}{64} - \frac{3x^2}{4} + 1\right)}{x} + c_2 \left(\frac{\frac{33x^4}{128} - \frac{x^2}{4}}{x} + \frac{\left(\frac{45x^4}{64} - \frac{3x^2}{4} + 1\right) \log(x)}{x}\right)$$

15.48 problem 44

Internal problem ID [1396]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 - 2x)y'' + 3y'x + (4x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(1-2*x)*diff(y(x),x$2)+3*x*diff(y(x),x)+(1+4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(c_2 \ln(x) + c_1)(1 + O(x^6)) + ((-6)x + 6x^2 - \frac{8}{3}x^3 + O(x^6))c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 40

```
AsymptoticDSolveValue[x^2*(1-2*x)*y'[x]+3*x*y'[x]+(1+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{-\frac{8x^3}{3} + 6x^2 - 6x}{x} + \frac{\log(x)}{x} \right) + \frac{c_1}{x}$$

15.49 problem 45

Internal problem ID [1397]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 45.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$x(x+1)y'' + (1-x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
Order:=6;
```

```
dsolve(x*(1+x)*diff(y(x),x$2)+(1-x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) (1 - x + O(x^6)) + (4x + O(x^6)) c_2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 27

```
AsymptoticDSolveValue[x*(1+x)*y''[x]+(1-x)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(1-x) + c_2(4x + (1-x)\log(x))$$

15.50 problem 46

Internal problem ID [1398]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 46.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2(1-x)y'' + x(-2x+3)y' + (1+2x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 57

Order:=6;

```
dsolve(x^2*(1-x)*diff(y(x),x$2)+x*(3-2*x)*diff(y(x),x)+(1+2*x)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \frac{(c_2 \ln(x) + c_1)(1 - 2x + x^2 + O(x^6)) + (3x - 3x^2 + \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{30}x^5 + O(x^6))c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 70

```
AsymptoticDSolveValue[x^2*(1-x)*y''[x]+x*(3-2*x)*y'[x]+(1+2*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1(x^2 - 2x + 1)}{x} + c_2 \left(\frac{(x^2 - 2x + 1) \log(x)}{x} + \frac{\frac{x^5}{30} + \frac{x^4}{12} + \frac{x^3}{3} - 3x^2 + 3x}{x} \right)$$

15.51 problem 47

Internal problem ID [1399]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 47.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x+1)y'' + 4y'x^2 + (1-5x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(4*x^2*(1+x)*diff(y(x),x$2)+4*x^2*diff(y(x),x)+(1-5*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left((c_2 \ln(x) + c_1) \left(1 + x - \frac{1}{4}x^2 + \frac{5}{36}x^3 - \frac{55}{576}x^4 + \frac{209}{2880}x^5 + O(x^6) \right) \right. \\ \left. + \left((-3)x + \frac{1}{4}x^2 - \frac{5}{54}x^3 + \frac{175}{3456}x^4 - \frac{2863}{86400}x^5 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 124

```
AsymptoticDSolveValue[4*x^2*(1+x)*y'[x]+4*x^2*y'[x]+(1-5*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{209x^5}{2880} - \frac{55x^4}{576} + \frac{5x^3}{36} - \frac{x^2}{4} + x + 1 \right) \\ + c_2 \left(\sqrt{x} \left(-\frac{2863x^5}{86400} + \frac{175x^4}{3456} - \frac{5x^3}{54} + \frac{x^2}{4} - 3x \right) \right. \\ \left. + \sqrt{x} \left(\frac{209x^5}{2880} - \frac{55x^4}{576} + \frac{5x^3}{36} - \frac{x^2}{4} + x + 1 \right) \log(x) \right)$$

15.52 problem 48

Internal problem ID [1400]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 48.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1-x)y'' - x(3-5x)y' + (4-5x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 61

Order:=6;

```
dsolve(x^2*(1-x)*diff(y(x),x$2)-x*(3-5*x)*diff(y(x),x)+(4-5*x)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \left((c_2 \ln(x) + c_1) (1 - 3x + 3x^2 - x^3 + O(x^6)) \right. \\ \left. + \left(4x - 7x^2 + \frac{11}{3}x^3 - \frac{1}{4}x^4 - \frac{1}{20}x^5 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 84

```
AsymptoticDSolveValue[x^2*(1-x)*y'[x]-x*(3-5*x)*y'[x]+(4-5*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(-x^3 + 3x^2 - 3x + 1)x^2 \\ + c_2 \left((-x^3 + 3x^2 - 3x + 1)x^2 \log(x) + \left(-\frac{x^5}{20} - \frac{x^4}{4} + \frac{11x^3}{3} - 7x^2 + 4x \right) x^2 \right)$$

15.53 problem 49

Internal problem ID [1401]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 49.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - x(9x^2 + 1)y' + (25x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-x*(1+9*x^2)*diff(y(x),x)+(1+25*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = x((c_2 \ln(x) + c_1)(1 - 4x^2 + x^4 + O(x^6)) + (6x^2 - 3x^4 + O(x^6))c_2)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 49

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]-x*(1+9*x^2)*y'[x]+(1+25*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x(x^4 - 4x^2 + 1) + c_2(x(6x^2 - 3x^4) + x(x^4 - 4x^2 + 1) \log(x))$$

15.54 problem 50

Internal problem ID [1402]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 50.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' + 3x(-x^2 + 1)y' + (7x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

Order:=6;

```
dsolve(9*x^2*dif(y(x),x$2)+3*x*(1-x^2)*dif(y(x),x)+(1+7*x^2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = x^{\frac{1}{3}} \left((c_2 \ln(x) + c_1) \left(1 - \frac{1}{6}x^2 + O(x^6) \right) + \left(\frac{1}{4}x^2 - \frac{1}{288}x^4 + O(x^6) \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 63

```
AsymptoticDSolveValue[9*x^2*y'[x]+3*x*(1-x^2)*y'[x]+(1+7*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(1 - \frac{x^2}{6} \right) + c_2 \left(\sqrt[3]{x} \left(1 - \frac{x^2}{6} \right) \log(x) + \sqrt[3]{x} \left(\frac{x^2}{4} - \frac{x^4}{288} \right) \right)$$

15.55 problem 51

Internal problem ID [1403]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 51.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(x^2 + 1)y'' + (-x^2 + 1)y' - 8yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

Order:=6;

```
dsolve(x*(1+x^2)*diff(y(x),x$2)+(1-x^2)*diff(y(x),x)-8*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) (1 + 2x^2 + x^4 + O(x^6)) + \left(-\frac{3}{2}x^2 - \frac{3}{2}x^4 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 48

```
AsymptoticDSolveValue[x*(1+x^2)*y'[x]+(1-x^2)*y'[x]-8*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(x^4 + 2x^2 + 1) + c_2 \left(-\frac{3x^4}{2} - \frac{3x^2}{2} + (x^4 + 2x^2 + 1) \log(x) \right)$$

15.56 problem 52

Internal problem ID [1404]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 52.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 2x(-x^2 + 4)y' + (7x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 51

Order:=6;

```
dsolve(4*x^2*dif(y(x),x$2)+2*x*(4-x^2)*dif(y(x),x)+(1+7*x^2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - \frac{1}{2}x^2 + \frac{1}{32}x^4 + O(x^6)\right) + \left(\frac{5}{8}x^2 - \frac{9}{128}x^4 + O(x^6)\right) c_2}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 77

```
AsymptoticDSolveValue[4*x^2*y'[x]+2*x*(4-x^2)*y'[x]+(1+7*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1 \left(\frac{x^4}{32} - \frac{x^2}{2} + 1\right)}{\sqrt{x}} + c_2 \left(\frac{\frac{5x^2}{8} - \frac{9x^4}{128}}{\sqrt{x}} + \frac{\left(\frac{x^4}{32} - \frac{x^2}{2} + 1\right) \log(x)}{\sqrt{x}} \right)$$

15.57 problem 58

Internal problem ID [1405]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 58.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x+1)y'' + 8y'x^2 + (x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

Order:=6;

```
dsolve(4*x^2*(1+x)*diff(y(x),x$2)+8*x^2*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x}(-x^5 + x^4 - x^3 + x^2 - x + 1)(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 64

```
AsymptoticDSolveValue[4*x^2*(1+x)*y'[x]+8*x^2*y'[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x}(-x^5 + x^4 - x^3 + x^2 - x + 1) + c_2 \sqrt{x}(-x^5 + x^4 - x^3 + x^2 - x + 1) \log(x)$$

15.58 problem 59

Internal problem ID [1406]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 59.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2(x+3)y'' + 3x(3+7x)y' + (4x+3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

Order:=6;

```
dsolve(9*x^2*(3+x)*diff(y(x),x$2)+3*x*(3+7*x)*diff(y(x),x)+(3+4*x)*y(x)=0,y(x),type='series')
```

$$y(x) = x^{\frac{1}{3}} \left(1 - \frac{1}{3}x + \frac{1}{9}x^2 - \frac{1}{27}x^3 + \frac{1}{81}x^4 - \frac{1}{243}x^5 \right) (c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 92

```
AsymptoticDSolveValue[9*x^2*(3+x)*y'[x]+3*x*(3+7*x)*y'[x]+(3+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{x^5}{243} + \frac{x^4}{81} - \frac{x^3}{27} + \frac{x^2}{9} - \frac{x}{3} + 1 \right) + c_2 \sqrt[3]{x} \left(-\frac{x^5}{243} + \frac{x^4}{81} - \frac{x^3}{27} + \frac{x^2}{9} - \frac{x}{3} + 1 \right) \log(x)$$

15.59 problem 60

Internal problem ID [1407]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 60.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-x^2 + 2)y'' - x(3x^2 + 2)y' + (-x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

Order:=6;

```
dsolve(x^2*(2-x^2)*diff(y(x),x$2)-x*(2+3*x^2)*diff(y(x),x)+(2-x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{4}x^4\right) x(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 46

```
AsymptoticDSolveValue[x^2*(2-x^2)*y'[x]-x*(2+3*x^2)*y'[x]+(2-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(\frac{x^4}{4} + \frac{x^2}{2} + 1 \right) + c_2 x \left(\frac{x^4}{4} + \frac{x^2}{2} + 1 \right) \log(x)$$

15.60 problem 61

Internal problem ID [1408]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 61.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$16x^2(x^2 + 1)y'' + 8x(9x^2 + 1)y' + (49x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 37

Order:=6;

```
dsolve(16*x^2*(1+x^2)*diff(y(x),x$2)+8*x*(1+9*x^2)*diff(y(x),x)+(1+49*x^2)*y(x)=0,y(x),type=
```

$$y(x) = x^{\frac{1}{4}}(x^4 - x^2 + 1)(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 42

```
AsymptoticDSolveValue[16*x^2*(1+x^2)*y'[x]+8*x*(1+9*x^2)*y'[x]+(1+49*x^2)*y[x]==0,y[x],{x,0
```

$$y(x) \rightarrow c_1 \sqrt[4]{x}(x^4 - x^2 + 1) + c_2 \sqrt[4]{x}(x^4 - x^2 + 1) \log(x)$$

15.61 problem 62

Internal problem ID [1409]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 62.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(3x + 4)y'' - x(4 - 3x)y' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(4+3*x)*diff(y(x),x$2)-x*(4-3*x)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{4}x + \frac{9}{16}x^2 - \frac{27}{64}x^3 + \frac{81}{256}x^4 - \frac{243}{1024}x^5\right) x(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 84

```
AsymptoticDSolveValue[x^2*(4+3*x)*y''[x]-x*(4-3*x)*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x \left(-\frac{243x^5}{1024} + \frac{81x^4}{256} - \frac{27x^3}{64} + \frac{9x^2}{16} - \frac{3x}{4} + 1 \right) + c_2 x \left(-\frac{243x^5}{1024} + \frac{81x^4}{256} - \frac{27x^3}{64} + \frac{9x^2}{16} - \frac{3x}{4} + 1 \right) \log(x)$$

15.62 problem 63

Internal problem ID [1410]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 63.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 3x + 1)y'' + 8x^2(2x + 3)y' + (9x^2 + 3x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(4*x^2*(1+3*x+x^2)*diff(y(x),x$2)+8*x^2*(3+2*x)*diff(y(x),x)+(1+3*x+9*x^2)*y(x)=0,y(x)
```

$$y(x) = \sqrt{x}(-144x^5 + 55x^4 - 21x^3 + 8x^2 - 3x + 1)(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 72

```
AsymptoticDSolveValue[4*x^2*(1+3*x+x^2)*y''[x]+8*x^2*(3+2*x)*y'[x]+(1+3*x+9*x^2)*y[x]==0,y[x]
```

$$y(x) \rightarrow c_1 \sqrt{x}(-144x^5 + 55x^4 - 21x^3 + 8x^2 - 3x + 1) + c_2 \sqrt{x}(-144x^5 + 55x^4 - 21x^3 + 8x^2 - 3x + 1) \log(x)$$

15.63 problem 64

Internal problem ID [1411]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 64.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1-x)^2 y'' - x(-3x^2 + 2x + 1) y' + y(x^2 + 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(x^2*(1-x)^2*diff(y(x),x$2)-x*(1+2*x-3*x^2)*diff(y(x),x)+(1+x^2)*y(x)=0,y(x),type='ser
```

$$y(x) = (6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1) x(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 64

```
AsymptoticDSolveValue[x^2*(1-x)^2*y'[x]-x*(1+2*x-3*x^2)*y'[x]+(1+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x(6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1) + c_2 x(6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1) \log(x)$$

15.64 problem 65

Internal problem ID [1412]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS II. Exercises 7.6. Page 374

Problem number: 65.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`

$$9x^2(x^2 + x + 1)y'' + 3x(13x^2 + 7x + 1)y' + (25x^2 + 4x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

Order:=6;

```
dsolve(9*x^2*(1+x+x^2)*diff(y(x),x$2)+3*x*(1+7*x+13*x^2)*diff(y(x),x)+(1+4*x+25*x^2)*y(x)=0,
```

$$y(x) = x^{\frac{1}{3}}(-x^4 + x^3 - x + 1)(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 48

```
AsymptoticDSolveValue[9*x^2*(1+x+x^2)*y'[x]+3*x*(1+7*x+13*x^2)*y'[x]+(1+4*x+25*x^2)*y[x]==0,
```

$$y(x) \rightarrow c_1 \sqrt[3]{x}(-x^4 + x^3 - x + 1) + c_2 \sqrt[3]{x}(-x^4 + x^3 - x + 1) \log(x)$$

16 Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

16.1	problem Example 7.7.1 page 381	613
16.2	problem Example 7.7.2 page 383	614
16.3	problem Example 7.7.3 page 385	615
16.4	problem Example 7.7.4 page 387	616
16.5	problem 1	617
16.6	problem 2	618
16.7	problem 3	619
16.8	problem 4	620
16.9	problem 5	621
16.10	problem 6	622
16.11	problem 7	623
16.12	problem 8	624
16.13	problem 9	625
16.14	problem 10	626
16.15	problem 11	627
16.16	problem 12	628
16.17	problem 13	629
16.18	problem 14	630
16.19	problem 15	631
16.20	problem 16	632
16.21	problem 17	633
16.22	problem 18	634
16.23	problem 19	635
16.24	problem 20	636
16.25	problem 21	637
16.26	problem 22	638
16.27	problem 23	639
16.28	problem 24	640
16.29	problem 25	641
16.30	problem 26	642
16.31	problem 27	643
16.32	problem 28	644
16.33	problem 29	645
16.34	problem 30	646

16.35problem 31	647
16.36problem 32	648
16.37problem 33	649
16.38problem 34	650
16.39problem 35	651
16.40problem 36	652
16.41problem 37	653
16.42problem 38	654
16.43problem 39	655
16.44problem 40	656

16.1 problem Example 7.7.1 page 381

Internal problem ID [1413]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: Example 7.7.1 page 381.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(2+x)y'' - x(4-7x)y' - (5-3x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 65

Order:=6;

`dsolve(2*x^2*(2+x)*diff(y(x),x$2)-x*(4-7*x)*diff(y(x),x)-(5-3*x)*y(x)=0,y(x),type='series',x`

$$y(x) = \frac{c_1 x^3 \left(1 - \frac{7}{4}x + \frac{63}{32}x^2 - \frac{231}{128}x^3 + \frac{3003}{2048}x^4 - \frac{9009}{8192}x^5 + O(x^6)\right) + c_2 (\ln(x) \left(-\frac{45}{32}x^3 + \frac{315}{128}x^4 - \frac{2835}{1024}x^5 + O(x^6)\right))}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 98

`AsymptoticDSolveValue[2*x^2*(2+x)*y'[x]-x*(4-7*x)*y'[x]-(5-3*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_2 \left(\frac{3003x^{13/2}}{2048} - \frac{231x^{11/2}}{128} + \frac{63x^{9/2}}{32} - \frac{7x^{7/2}}{4} + x^{5/2} \right) + c_1 \left(\frac{15}{512}(7x-4)x^{5/2} \log(x) + \frac{809x^4 - 548x^3 + 96x^2 + 128x + 1024}{1024\sqrt{x}} \right)$$

16.2 problem Example 7.7.2 page 383

Internal problem ID [1414]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: Example 7.7.2 page 383.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 - 2x)y'' + x(8 - 9x)y' + (6 - 3x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(1-2*x)*diff(y(x),x$2)+x*(8-9*x)*diff(y(x),x)+(6-3*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{3}x - \frac{1}{14}x^2 - \frac{1}{28}x^3 - \frac{25}{1008}x^4 - \frac{1}{48}x^5 + O(x^6)\right)}{x} + \frac{c_2(2880 - 23760x + 71280x^2 - 83160x^3 + 62370x^5 + O(x^6))}{x^6}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 61

```
AsymptoticDSolveValue[x^2*(1-2*x)*y'[x]+x*(8-9*x)*y'[x]+(6-3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{25x^3}{1008} - \frac{x^2}{28} - \frac{x}{14} + \frac{1}{x} - \frac{1}{3} \right) + c_1 \left(\frac{1}{x^6} - \frac{33}{4x^5} + \frac{99}{4x^4} - \frac{231}{8x^3} \right)$$

16.3 problem Example 7.7.3 page 385

Internal problem ID [1415]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: Example 7.7.3 page 385.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' + x(10x^2 + 3)y' - (-14x^2 + 15)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*(3+10*x^2)*diff(y(x),x)-(15-14*x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 x^3 \left(1 - \frac{5}{2}x^2 + \frac{35}{8}x^4 + O(x^6) \right) + \frac{c_2(-203212800 + 101606400x^2 - 25401600x^4 + O(x^6))}{x^5}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 46

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]+x*(3+10*x^2)*y'[x]-(15-14*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^5} - \frac{1}{2x^3} + \frac{1}{8x} \right) + c_2 \left(\frac{35x^7}{8} - \frac{5x^5}{2} + x^3 \right)$$

16.4 problem Example 7.7.4 page 387

Internal problem ID [1416]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: Example 7.7.4 page 387.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(-2x^2 + 1)y'' + x(-13x^2 + 7)y' - 14x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

Order:=6;

```
dsolve(x^2*(1-2*x^2)*diff(y(x),x$2)+x*(7-13*x^2)*diff(y(x),x)-14*x^2*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 \left(1 + \frac{7}{8}x^2 + \frac{77}{80}x^4 + O(x^6) \right) + \frac{c_2(-86400 + 216000x^2 - 54000x^4 + O(x^6))}{x^6}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 44

```
AsymptoticDSolveValue[x^2*(1-2*x^2)*y'[x]+x*(7-13*x^2)*y'[x]-14*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{77x^4}{80} + \frac{7x^2}{8} + 1 \right) + c_1 \left(\frac{1}{x^6} - \frac{5}{2x^4} + \frac{5}{8x^2} \right)$$

16.5 problem 1

Internal problem ID [1417]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - 3y'x + (4x + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 61

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+(3+4*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(c_1 x^2 \left(1 - \frac{4}{3}x + \frac{2}{3}x^2 - \frac{8}{45}x^3 + \frac{4}{135}x^4 - \frac{16}{4725}x^5 + O(x^6) \right) + c_2 \left(\ln(x) \left(16x^2 - \frac{64}{3}x^3 + \frac{32}{3}x^4 - \frac{128}{45}x^5 + O(x^6) \right) + \left(-2 - 8x + \frac{256}{9}x^3 - \frac{200}{9}x^4 + \frac{5024}{675}x^5 + O(x^6) \right) \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^2*y''[x]-3*x*y'[x]+(3+4*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{9}x(124x^4 - 176x^3 + 36x^2 + 36x + 9) - \frac{8}{3}x^3(2x^2 - 4x + 3) \log(x) \right) + c_2 \left(\frac{4x^7}{135} - \frac{8x^6}{45} + \frac{2x^5}{3} - \frac{4x^4}{3} + x^3 \right)$$

16.6 problem 2

Internal problem ID [1418]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [[_Emden, _Fowler]]

$$xy'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{144}x^3 + \frac{1}{2880}x^4 - \frac{1}{86400}x^5 + O(x^6) \right) \\ + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 + \frac{1}{144}x^4 - \frac{1}{2880}x^5 + O(x^6) \right) \right. \\ \left. + \left(1 - \frac{3}{4}x^2 + \frac{7}{36}x^3 - \frac{35}{1728}x^4 + \frac{101}{86400}x^5 + O(x^6) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{144}x(x^3 - 12x^2 + 72x - 144) \log(x) \right. \\ \left. + \frac{-47x^4 + 480x^3 - 2160x^2 + 1728x + 1728}{1728} \right) + c_2 \left(\frac{x^5}{2880} - \frac{x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

16.7 problem 3

Internal problem ID [1419]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x+1)y'' + 4x(1+2x)y' - (3x+1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 51

Order:=6;

```
dsolve(4*x^2*(1+x)*diff(y(x),x$2)+4*x*(1+2*x)*diff(y(x),x)-(1+3*x)*y(x)=0,y(x),type='series')
```

$y(x)$

$$= \frac{c_1(1 + O(x^6))x + \ln(x)(x + O(x^6))c_2 + (1 - x - x^2 + \frac{1}{2}x^3 - \frac{1}{3}x^4 + \frac{1}{4}x^5 + O(x^6))c_2}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 53

```
AsymptoticDSolveValue[4*x^2*(1+x)*y'[x]+4*x*(1+2*x)*y'[x]-(1+3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\sqrt{x} \log(x) - \frac{2x^4 - 3x^3 + 6x^2 + 6x - 6}{6\sqrt{x}} \right) + c_2 \sqrt{x}$$

16.8 problem 4

Internal problem ID [1420]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x(x+1)y'' + y'x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 60

```
Order:=6;
```

```
dsolve(x*(1+x)*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x \left(1 - x + \frac{5}{6}x^2 - \frac{25}{36}x^3 + \frac{85}{144}x^4 - \frac{221}{432}x^5 + O(x^6) \right) \\ & + \left(-x + x^2 - \frac{5}{6}x^3 + \frac{25}{36}x^4 - \frac{85}{144}x^5 + O(x^6) \right) \ln(x) c_2 \\ & + \left(1 - x + \frac{1}{2}x^2 - \frac{7}{18}x^3 + \frac{145}{432}x^4 - \frac{257}{864}x^5 + O(x^6) \right) c_2 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x*(1+x)*y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left(\frac{1}{36}x(25x^3 - 30x^2 + 36x - 36) \log(x) \right. \\ & \left. + \frac{1}{432}(-455x^4 + 552x^3 - 648x^2 + 432x + 432) \right) + c_2 \left(\frac{85x^5}{144} - \frac{25x^4}{36} + \frac{5x^3}{6} - x^2 + x \right) \end{aligned}$$

16.9 problem 5

Internal problem ID [1421]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$2x^2(3x + 2)y'' + x(4 + 21x)y' - (1 - 9x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 69

Order:=6;

`dsolve(2*x^2*(2+3*x)*diff(y(x),x$2)+x*(4+21*x)*diff(y(x),x)-(1-9*x)*y(x)=0,y(x),type='series`

$$y(x) = \frac{c_1 x \left(1 - \frac{9}{4}x + \frac{135}{32}x^2 - \frac{945}{128}x^3 + \frac{25515}{2048}x^4 - \frac{168399}{8192}x^5 + O(x^6)\right) + c_2 (\ln(x) \left(-\frac{3}{4}x + \frac{27}{16}x^2 - \frac{405}{128}x^3 + \frac{2835}{512}x^4 - \frac{168399}{8192}x^5 + O(x^6)\right))}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 108

`AsymptoticDSolveValue[2*x^2*(2+3*x)*y'[x]+x*(4+21*x)*y'[x]-(1-9*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_2 \left(\frac{25515x^{9/2}}{2048} - \frac{945x^{7/2}}{128} + \frac{135x^{5/2}}{32} - \frac{9x^{3/2}}{4} + \sqrt{x} \right) + c_1 \left(\frac{3}{512} \sqrt{x} (945x^3 - 540x^2 + 288x - 128) \log(x) + \frac{8613x^4 - 5076x^3 + 2880x^2 - 1536x + 1024}{1024\sqrt{x}} \right)$$

16.10 problem 6

Internal problem ID [1422]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(2+x)y' - (-3x+2)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 63

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*(2+x)*diff(y(x),x)-(2-3*x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^3 \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + O(x^6)\right) + c_2 (\ln(x) \left((-6)x^3 + 6x^4 - 3x^5 + O(x^6)\right) + (12 + 6x^2))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 74

```
AsymptoticDSolveValue[x^2*y''[x]+x*(2+x)*y'[x]-(2-3*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4 - 3x^3 + 2x^2 + 2x + 4}{4x^2} + \frac{1}{2}(x-1)x \log(x) \right) + c_2 \left(\frac{x^5}{24} - \frac{x^4}{6} + \frac{x^3}{2} - x^2 + x \right)$$

16.11 problem 7

Internal problem ID [1423]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 4y'x - (9 - x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 63

Order:=6;

`dsolve(4*x^2*diff(y(x),x$2)+4*x*diff(y(x),x)-(9-x)*y(x)=0,y(x),type='series',x=0);`

$$y(x) = \frac{c_1 x^3 \left(1 - \frac{1}{16}x + \frac{1}{640}x^2 - \frac{1}{46080}x^3 + \frac{1}{5160960}x^4 - \frac{1}{825753600}x^5 + O(x^6)\right) + c_2 (\ln(x) \left(-\frac{1}{64}x^3 + \frac{1}{1024}x^4 - \frac{1}{40960}x^5\right) + \frac{1}{x^{\frac{3}{2}}}}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 96

`AsymptoticDSolveValue[4*x^2*y''[x]+4*x*y'[x]-(9-x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_2 \left(\frac{x^{11/2}}{5160960} - \frac{x^{9/2}}{46080} + \frac{x^{7/2}}{640} - \frac{x^{5/2}}{16} + x^{3/2} \right) + c_1 \left(\frac{(x-16)x^{3/2} \log(x)}{12288} - \frac{19x^4 - 64x^3 - 2304x^2 - 18432x - 147456}{147456x^{3/2}} \right)$$

16.12 problem 8

Internal problem ID [1424]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + 10y'x + (x + 14)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 59

Order:=6;

```
dsolve(x^2*diff(y(x),x$2)+10*x*diff(y(x),x)+(14+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x + \frac{1}{84}x^2 - \frac{1}{2016}x^3 + \frac{1}{72576}x^4 - \frac{1}{3628800}x^5 + O(x^6)\right) x^5 + c_2 (\ln(x) (-x^5 + O(x^6))) + (2880 + 7200 \ln(x))}{x^7}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 68

```
AsymptoticDSolveValue[x^2*y'[x]+10*x*y'[x]+(14+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^2}{72576} + \frac{1}{x^2} - \frac{x}{2016} - \frac{1}{6x} + \frac{1}{84} \right) + c_1 \left(\frac{1}{x^7} + \frac{1}{4x^6} + \frac{1}{24x^5} + \frac{1}{144x^4} + \frac{1}{576x^3} \right)$$

16.13 problem 9

Internal problem ID [1425]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x+1)y'' + 4x(3+8x)y' - (5-49x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 65

Order:=6;

```
dsolve(4*x^2*(1+x)*diff(y(x),x$2)+4*x*(3+8*x)*diff(y(x),x)-(5-49*x)*y(x)=0,y(x),type='series
```

$$y(x) = \frac{c_1 x^3 (1 - 4x + 10x^2 - 20x^3 + 35x^4 - 56x^5 + O(x^6)) + c_2 (\ln(x) ((-36)x^3 + 144x^4 - 360x^5 + O(x^6)) + x^{\frac{5}{2}})}{x^{\frac{5}{2}}}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 86

```
AsymptoticDSolveValue[4*x^2*(1+x)*y'[x]+4*x*(3+8*x)*y'[x]-(5-49*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 (35x^{9/2} - 20x^{7/2} + 10x^{5/2} - 4x^{3/2} + \sqrt{x}) + c_1 \left(\frac{62x^4 - 20x^3 + 2x^2 + x + 2}{2x^{5/2}} + 3\sqrt{x}(4x - 1) \log(x) \right)$$

16.14 problem 10

Internal problem ID [1426]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' - x(3+10x)y' + 30yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 48

Order:=6;

```
dsolve(x^2*(1+x)*diff(y(x),x$2)-x*(3+10*x)*diff(y(x),x)+30*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^4 \left(1 - \frac{2}{5}x + O(x^6) \right) + \ln(x) (43200x^4 - 17280x^5 + O(x^6)) c_2 \\ + (-144 - 1440x - 7200x^2 - 28800x^3 - 90720x^4 + 82944x^5 + O(x^6)) c_2$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 48

```
AsymptoticDSolveValue[x^2*(1+x)*y''[x]-x*(3+10*x)*y'[x]+30*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x^4 - \frac{2x^5}{5} \right) + c_1 (745x^4 - 300x^4 \log(x) + 200x^3 + 50x^2 + 10x + 1)$$

16.15 problem 11

Internal problem ID [1427]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(x+1)y' - 3y(x+3) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 37

Order:=6;

```
dsolve(x^2*diff(y(x),x$2)+x*(1+x)*diff(y(x),x)-3*(3+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^3 (1 + O(x^6)) + \frac{c_2 (-86400 + 103680x - 64800x^2 + 28800x^3 - 10800x^4 + 4320x^5 + O(x^6))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 39

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1+x)*y'[x]-3*(3+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x^3 + c_1 \left(\frac{1}{x^3} - \frac{6}{5x^2} + \frac{x}{8} + \frac{3}{4x} - \frac{1}{3} \right)$$

16.16 problem 12

Internal problem ID [1428]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(1 - 2x) y' - (x + 4) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 61

Order:=6;

```
dsolve(x^2*dif(y(x),x$2)+x*(1-2*x)*dif(y(x),x)-(4+x)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{c_1 x^4 \left(1 + x + \frac{7}{12} x^2 + \frac{1}{4} x^3 + \frac{11}{128} x^4 + \frac{143}{5760} x^5 + O(x^6)\right) + c_2 (\ln(x) (9x^4 + 9x^5 + O(x^6))) + (-144 - 144x)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 75

```
AsymptoticDSolveValue[x^2*y'[x]+x*(1-2*x)*y'[x]-(4+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{3x^4 - 16x^3 + 48x^2 + 192x + 192}{192x^2} - \frac{1}{16} x^2 \log(x) \right) + c_2 \left(\frac{11x^6}{128} + \frac{x^5}{4} + \frac{7x^4}{12} + x^3 + x^2 \right)$$

16.17 problem 13

Internal problem ID [1429]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x(x+1)y'' - 4y' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 40

```
Order:=6;  
dsolve(x*(1+x)*diff(y(x),x$2)-4*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^5 (1 - 3x + 6x^2 - 10x^3 + 15x^4 - 21x^5 + O(x^6)) \\ + c_2 (2880 - 1440x + 480x^2 - 480x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 48

```
AsymptoticDSolveValue[x*(1+x)*y''[x]-4*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^2}{6} - \frac{x}{2} + 1 \right) + c_2 (15x^9 - 10x^8 + 6x^7 - 3x^6 + x^5)$$

16.18 problem 14

Internal problem ID [1430]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 + 2x)y'' + x(9 + 13x)y' + (7 + 5x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

`Order:=6;`

`dsolve(x^2*(1+2*x)*diff(y(x),x$2)+x*(9+13*x)*diff(y(x),x)+(7+5*x)*y(x)=0,y(x),type='series',`

$$y(x) = \frac{c_1 \left(1 + \frac{4}{7}x - \frac{5}{28}x^2 + \frac{5}{42}x^3 - \frac{5}{48}x^4 + \frac{7}{66}x^5 + O(x^6)\right)}{x} + \frac{c_2(-86400 - 449280x - 617760x^2 + O(x^6))}{x^7}$$

✓ Solution by Mathematica

Time used: 0.059 (sec). Leaf size: 54

`AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]+x*(9+13*x)*y'[x]+(7+5*x)*y[x]==0,y[x],{x,0,5}]`

$$y(x) \rightarrow c_2 \left(-\frac{5x^3}{48} + \frac{5x^2}{42} - \frac{5x}{28} + \frac{1}{x} + \frac{4}{7} \right) + c_1 \left(\frac{1}{x^7} + \frac{26}{5x^6} + \frac{143}{20x^5} \right)$$

16.19 problem 15

Internal problem ID [1431]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(1+2x)y'' - 2x(-x+4)y' - (7+5x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 41

Order:=6;

```
dsolve(4*x^2*(1+2*x)*diff(y(x),x$2)-2*x*(4-x)*diff(y(x),x)-(7+5*x)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{18}{5}x + \frac{39}{4}x^2 - \frac{663}{28}x^3 + \frac{13923}{256}x^4 - \frac{7735}{64}x^5 + O(x^6)\right) + c_2 \left(-144 - \frac{405}{8}x^4 + \frac{729}{4}x^5 + O(x^6)\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 67

```
AsymptoticDSolveValue[4*x^2*(1+2*x)*y'[x]-2*x*(4-x)*y'[x]-(7+5*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{\sqrt{x}} - \frac{35x^{7/2}}{128} \right) + c_2 \left(\frac{13923x^{15/2}}{256} - \frac{663x^{13/2}}{28} + \frac{39x^{11/2}}{4} - \frac{18x^{9/2}}{5} + x^{7/2} \right)$$

16.20 problem 16

Internal problem ID [1432]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3x^2(x+3)y'' - x(15+x)y' - 20y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

`Order:=6;`

```
dsolve(3*x^2*(3+x)*diff(y(x),x$2)-x*(15+x)*diff(y(x),x)-20*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{4}{9}x + \frac{13}{81}x^2 - \frac{832}{15309}x^3 + \frac{2470}{137781}x^4 - \frac{21736}{3720087}x^5 + O(x^6)\right) + c_2 \left(-144 - \frac{64}{3}x + \frac{16}{27}x^2 - \frac{112}{6561}x^4 + \frac{448}{59049}x^5\right)}{x^{\frac{2}{3}}}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 85

```
AsymptoticDSolveValue[3*x^2*(3+x)*y'[x]-x*(15+x)*y'[x]-20*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{7x^{10/3}}{59049} - \frac{x^{4/3}}{243} + \frac{1}{x^{2/3}} + \frac{4\sqrt[3]{x}}{27} \right) + c_2 \left(\frac{2470x^{22/3}}{137781} - \frac{832x^{19/3}}{15309} + \frac{13x^{16/3}}{81} - \frac{4x^{13/3}}{9} + x^{10/3} \right)$$

16.21 problem 17

Internal problem ID [1433]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' + x(1-10x)y' - (9-10x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 45

Order:=6;

```
dsolve(x^2*(1+x)*diff(y(x),x$2)+x*(1-10*x)*diff(y(x),x)-(9-10*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = c_1 x^3 \left(1 + 2x + \frac{9}{4}x^2 + \frac{5}{3}x^3 + \frac{5}{6}x^4 + \frac{3}{11}x^5 + O(x^6) \right) + \frac{c_2(-86400 - 898560x - 4043520x^2 - 9884160x^3 - 12355200x^4 + O(x^6))}{x^3}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 64

```
AsymptoticDSolveValue[x^2*(1+x)*y'[x]+x*(1-10*x)*y'[x]-(9-10*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^3} + \frac{52}{5x^2} + 143x + \frac{234}{5x} + \frac{572}{5} \right) + c_2 \left(\frac{5x^7}{6} + \frac{5x^6}{3} + \frac{9x^5}{4} + 2x^4 + x^3 \right)$$

16.22 problem 18

Internal problem ID [1434]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' + 3y'x^2 - (6-x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 39

```
Order:=6;
```

```
dsolve(x^2*(1+x)*diff(y(x),x$2)+3*x^2*diff(y(x),x)-(6-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^3 \left(1 - \frac{8}{3}x + \frac{100}{21}x^2 - \frac{50}{7}x^3 + \frac{175}{18}x^4 - \frac{112}{9}x^5 + O(x^6) \right) + \frac{c_2(2880 + 720x + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 53

```
AsymptoticDSolveValue[x^2*(1+x)*y''[x]+3*x^2*y'[x]-(6-x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^2} + \frac{1}{4x} \right) + c_2 \left(\frac{175x^7}{18} - \frac{50x^6}{7} + \frac{100x^5}{21} - \frac{8x^4}{3} + x^3 \right)$$

16.23 problem 19

Internal problem ID [1435]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1+2x)y'' - 2x(3+14x)y' + (6+100x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

Order:=6;

```
dsolve(x^2*(1+2*x)*diff(y(x),x$2)-2*x*(3+14*x)*diff(y(x),x)+(6+100*x)*y(x)=0,y(x),type='series')
```

$$y(x) = c_1 x^6 \left(1 + \frac{4}{3}x + \frac{8}{7}x^2 + \frac{4}{7}x^3 + \frac{8}{63}x^4 + O(x^6) \right) + c_2 x (2880 + 51840x + 414720x^2 + 1935360x^3 + 5806080x^4 + 11612160x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 64

```
AsymptoticDSolveValue[x^2*(1+2*x)*y''[x]-2*x*(3+14*x)*y'[x]+(6+100*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 (2016x^5 + 672x^4 + 144x^3 + 18x^2 + x) + c_2 \left(\frac{8x^{10}}{63} + \frac{4x^9}{7} + \frac{8x^8}{7} + \frac{4x^7}{3} + x^6 \right)$$

16.24 problem 20

Internal problem ID [1436]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x+1)y'' - x(6+11x)y' + (6+32x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

Order:=6;

```
dsolve(x^2*(1+x)*diff(y(x),x$2)-x*(6+11*x)*diff(y(x),x)+(6+32*x)*y(x)=0,y(x),type='series',x
```

$$y(x) = c_1 x^6 \left(1 + \frac{2}{3}x + \frac{1}{7}x^2 + O(x^6) \right) \\ + c_2 x (2880 + 15120x + 30240x^2 + 25200x^3 - 15120x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 51

```
AsymptoticDSolveValue[x^2*(1+x)*y'[x]-x*(6+11*x)*y'[x]+(6+32*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^8}{7} + \frac{2x^7}{3} + x^6 \right) + c_1 \left(\frac{35x^4}{4} + \frac{21x^3}{2} + \frac{21x^2}{4} + x \right)$$

16.25 problem 21

Internal problem ID [1437]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x+1)y'' + 4x(4x+1)y' - (49+27x)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

Order:=6;

```
dsolve(4*x^2*(1+x)*diff(y(x),x$2)+4*x*(1+4*x)*diff(y(x),x)-(49+27*x)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 x^7 (1 - 2x + 3x^2 - 4x^3 + 5x^4 - 6x^5 + O(x^6)) + c_2 (3628800 - 3024000x + 2419200x^2 - 1814400x^3 + \dots)}{x^{\frac{7}{2}}}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 86

```
AsymptoticDSolveValue[4*x^2*(1+x)*y'[x]+4*x*(1+4*x)*y'[x]-(49+27*x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{2}{3x^{3/2}} - \frac{5}{6x^{5/2}} + \frac{1}{x^{7/2}} + \frac{\sqrt{x}}{3} - \frac{1}{2\sqrt{x}} \right) + c_2 (5x^{15/2} - 4x^{13/2} + 3x^{11/2} - 2x^{9/2} + x^{7/2})$$

16.26 problem 22

Internal problem ID [1438]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$x^2(1 + 2x)y'' - x(9 + 8x)y' - 12yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
Order:=6;
```

```
dsolve(x^2*(1+2*x)*diff(y(x),x$2)-x*(9+8*x)*diff(y(x),x)-12*x*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = c_1 x^{10} (1 - 8x + 40x^2 - 160x^3 + 560x^4 - 1792x^5 + O(x^6)) \\ + c_2 (-1316818944000 + 1755758592000x - 2194698240000x^2 \\ + 2508226560000x^3 - 2508226560000x^4 + 2006581248000x^5 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 62

```
AsymptoticDSolveValue[x^2*(1+2*x)*y'[x]-x*(9+8*x)*y'[x]-12*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{40x^4}{21} - \frac{40x^3}{21} + \frac{5x^2}{3} - \frac{4x}{3} + 1 \right) + c_2 (560x^{14} - 160x^{13} + 40x^{12} - 8x^{11} + x^{10})$$

16.27 problem 23

Internal problem ID [1439]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - x(-2x^2 + 7)y' + 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-x*(7-2*x^2)*diff(y(x),x)+12*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \left(c_1 x^4 \left(1 - \frac{7}{2}x^2 + \frac{63}{8}x^4 + O(x^6) \right) + c_2 \left(\ln(x) (1080x^4 + O(x^6)) + (-144 - 216x^2 + 2106x^4 + O(x^6)) \right) \right) x^2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 57

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]-x*(7-2*x^2)*y'[x]+12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{63x^{10}}{8} - \frac{7x^8}{2} + x^6 \right) + c_1 \left(-\frac{15}{2}x^6 \log(x) - \frac{1}{4}(31x^4 - 6x^2 - 4)x^2 \right)$$

16.28 problem 24

Internal problem ID [1440]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(-x^2 + 7) y' + 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(x^2*dif(y(x),x$2)-x*(7-x^2)*dif(y(x),x)+12*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(c_1 x^4 \left(1 - \frac{1}{2} x^2 + \frac{1}{8} x^4 + O(x^6) \right) + c_2 (\ln(x) (72x^4 + O(x^6)) + (-144 - 72x^2 + 54x^4 + O(x^6))) \right) x^2$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 55

```
AsymptoticDSolveValue[x^2*y''[x]-x*(7-x^2)*y'[x]+12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^{10}}{8} - \frac{x^8}{2} + x^6 \right) + c_1 \left(-\frac{1}{2} x^6 \log(x) - \frac{1}{4} (x^4 - 2x^2 - 4) x^2 \right)$$

16.29 problem 25

Internal problem ID [1441]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Lienard]

$$xy'' - 5y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
Order:=6;  
dsolve(x*diff(y(x),x$2)-5*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^6 \left(1 - \frac{1}{16} x^2 + \frac{1}{640} x^4 + O(x^6) \right) + c_2 (-86400 - 10800x^2 - 1350x^4 + O(x^6))$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 44

```
AsymptoticDSolveValue[x*y''[x]-5*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{64} + \frac{x^2}{8} + 1 \right) + c_2 \left(\frac{x^{10}}{640} - \frac{x^8}{16} + x^6 \right)$$

16.30 problem 26

Internal problem ID [1442]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(2x^2 + 1) y' - (-10x^2 + 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(x^2*dif(y(x),x$2)+x*(1+2*x^2)*dif(y(x),x)-(1-10*x^2)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{3}{2} x^2 + x^4 + O(x^6)\right) + c_2 (\ln(x) (8x^2 - 12x^4 + O(x^6)) + (-2 + 2x^2 + 4x^4 + O(x^6)))}{x}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 52

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1+2*x^2)*y'[x]-(1-10*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x^5 - \frac{3x^3}{2} + x \right) + c_1 \left(2x(3x^2 - 2) \log(x) - \frac{5x^4 - x^2 - 1}{x} \right)$$

16.31 problem 27

Internal problem ID [1443]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - y' x - (-x^2 + 3) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(x^2*dif(y(x),x^2)-x*dif(y(x),x)-(3-x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{1}{12} x^2 + \frac{1}{384} x^4 + O(x^6)\right) + c_2 (\ln(x) (9x^4 + O(x^6)) + (-144 - 36x^2 + O(x^6)))}{x}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 52

```
AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]-(3-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{(x^2 + 8)^2}{64x} - \frac{1}{16} x^3 \log(x) \right) + c_2 \left(\frac{x^7}{384} - \frac{x^5}{12} + x^3 \right)$$

16.32 problem 28

Internal problem ID [1444]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 2x(x^2 + 8)y' + (3x^2 + 5)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

```
Order:=6;
```

```
dsolve(4*x^2*diff(y(x),x$2)+2*x*(8+x^2)*diff(y(x),x)+(5+3*x^2)*y(x)=0,y(x),type='series',x=0
```

$y(x)$

$$= \frac{c_1 \left(1 - \frac{1}{16}x^2 + \frac{1}{256}x^4 + O(x^6)\right) x^2 + c_2 \left(\ln(x) \left(-\frac{1}{2}x^2 + \frac{1}{32}x^4 + O(x^6)\right) + \left(-2 + \frac{1}{2}x^2 - \frac{3}{128}x^4 + O(x^6)\right)\right)}{x^{\frac{5}{2}}}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 72

```
AsymptoticDSolveValue[4*x^2*y''[x]+2*x*(8+x^2)*y'[x]+(5+3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^{7/2}}{256} - \frac{x^{3/2}}{16} + \frac{1}{\sqrt{x}} \right) + c_1 \left(\frac{5x^4 - 96x^2 + 256}{256x^{5/2}} - \frac{(x^2 - 16) \log(x)}{64\sqrt{x}} \right)$$

16.33 problem 29

Internal problem ID [1445]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$x^2 y'' + x(x^2 + 1) y' - (-3x^2 + 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

Order:=6;

```
dsolve(x^2*diff(y(x),x$2)+x*(1+x^2)*diff(y(x),x)-(1-3*x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{2} x^2 + \frac{1}{8} x^4 + O(x^6)\right) + c_2 (\ln(x) (2x^2 - x^4 + O(x^6)) + (-2 + x^2 + O(x^6)))}{x}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 51

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1+x^2)*y'[x]-(1-3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{8} - \frac{x^3}{2} + x \right) + c_1 \left(\frac{1}{2} x (x^2 - 2) \log(x) - \frac{x^4 - 4}{4x} \right)$$

16.34 problem 30

Internal problem ID [1446]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(-2x^2 + 1) y' - 4y(2x^2 + 1) = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 49

```
Order:=6;
```

```
dsolve(x^2*dif(y(x),x$2)+x*(1-2*x^2)*dif(y(x),x)-4*(1+2*x^2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \frac{c_1 x^4 (1 + x^2 + \frac{1}{2} x^4 + O(x^6)) + c_2 (\ln(x) (288x^4 + O(x^6)) + (-144 + 144x^2 + 216x^4 + O(x^6)))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 45

```
AsymptoticDSolveValue[x^2*y'[x]+x*(1-2*x^2)*y'[x]-4*(1+2*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-2x^2 \log(x) - \frac{x^4 + x^2 - 1}{x^2} \right) + c_2 \left(\frac{x^6}{2} + x^4 + x^2 \right)$$

16.35 problem 31

Internal problem ID [1447]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 31.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2y'' + 8y'x - (-x^2 + 35)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

Order:=6;

```
dsolve(4*x^2*diff(y(x),x$2)+8*x*diff(y(x),x)-(35-x^2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 x^6 \left(1 - \frac{1}{64}x^2 + \frac{1}{10240}x^4 + O(x^6)\right) + c_2 \left(-86400 - 2700x^2 - \frac{675}{8}x^4 + O(x^6)\right)}{x^{\frac{7}{2}}}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 58

```
AsymptoticDSolveValue[4*x^2*y''[x]+8*x*y'[x]-(35-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{32x^{3/2}} + \frac{1}{x^{7/2}} + \frac{\sqrt{x}}{1024} \right) + c_2 \left(\frac{x^{13/2}}{10240} - \frac{x^{9/2}}{64} + x^{5/2} \right)$$

16.36 problem 32

Internal problem ID [1448]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 32.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$9x^2y'' - 3x(2x^2 + 11)y' + (10x^2 + 13)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

Order:=6;

```
dsolve(9*x^2*diff(y(x),x$2)-3*x*(11+2*x^2)*diff(y(x),x)+(13+10*x^2)*y(x)=0,y(x),type='series
```

$$y(x) = x^{\frac{1}{3}} \left(\left(1 + \frac{4}{27}x^2 + \frac{7}{486}x^4 + O(x^6) \right) x^4 c_1 + c_2 \left(\ln(x) \left(-\frac{32}{9}x^4 + O(x^6) \right) + \left(-144 - 32x^2 - \frac{8}{3}x^4 + O(x^6) \right) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 62

```
AsymptoticDSolveValue[9*x^2*y'[x]-3*x*(11+2*x^2)*y'[x]+(13+10*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^{25/3}}{486} + \frac{4x^{19/3}}{27} + x^{13/3} \right) + c_1 \left(\frac{2}{81}x^{13/3} \log(x) + \frac{1}{81}(x^2 + 9)^2 \sqrt[3]{x} \right)$$

16.37 problem 33

Internal problem ID [1449]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[_2nd_order, _with_linear_symmetries]`, `[_2nd_order, _linear,`

$$x^2 y'' + x(-2x^2 + 1) y' - 4(1 - x^2) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*(1-2*x^2)*diff(y(x),x)-4*(1-x^2)*y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= c_1 x^2 (1 + O(x^6)) + \frac{c_2 (\ln(x) (288x^4 + O(x^6)) + (-144 - 288x^2 - 216x^4 + O(x^6)))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 37

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1-2*x^2)*y'[x]-4*(1-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x^2 + c_1 \left(\frac{2x^4 + 2x^2 + 1}{x^2} - 2x^2 \log(x) \right)$$

16.38 problem 34

Internal problem ID [1450]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + x(-3x^2 + 1) y' - 4(-3x^2 + 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x*(1-3*x^2)*diff(y(x),x)-4*(1-3*x^2)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = c_1 x^2 \left(1 - \frac{1}{2} x^2 + O(x^6) \right) + \frac{c_2 (\ln(x) (1944x^4 + O(x^6)) + (-144 - 648x^2 - 810x^4 + O(x^6)))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 50

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1-3*x^2)*y'[x]-4*(1-3*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x^2 - \frac{x^4}{2} \right) + c_1 \left(\frac{18x^4 + 9x^2 + 2}{2x^2} - \frac{27}{2} x^2 \log(x) \right)$$

16.39 problem 35

Internal problem ID [1451]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 35.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' + x(11x^2 + 5)y' + 24x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*(5+11*x^2)*diff(y(x),x)+24*x^2*y(x)=0,y(x),type='series')
```

$$y(x) = c_1(1 - 2x^2 + 3x^4 + O(x^6)) + \frac{c_2(-144 + 432x^4 + O(x^6))}{x^4}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 27

```
AsymptoticDSolveValue[x^2*(1+x^2)*y''[x]+x*(5+11*x^2)*y'[x]+24*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\left(\frac{1}{x^4} - 1\right) + c_2(3x^4 - 2x^2 + 1)$$

16.40 problem 36

Internal problem ID [1452]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 36.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 1)y'' + 8y'x - (-x^2 + 35)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(4*x^2*(1+x^2)*diff(y(x),x$2)+8*x*diff(y(x),x)-(35-x^2)*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = \frac{c_1 x^6 \left(1 - \frac{1}{4}x^2 + \frac{1}{10}x^4 + O(x^6)\right) + c_2 \left(-86400 - 172800x^2 - 86400x^4 + O(x^6)\right)}{x^{\frac{7}{2}}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 52

```
AsymptoticDSolveValue[4*x^2*(1+x^2)*y'[x]+8*x*y'[x]-(35-x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{2}{x^{3/2}} + \frac{1}{x^{7/2}} + \sqrt{x} \right) + c_2 \left(\frac{x^{13/2}}{10} - \frac{x^{9/2}}{4} + x^{5/2} \right)$$

16.41 problem 37

Internal problem ID [1453]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 37.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' - x(-x^2 + 5)y' - (25x^2 + 7)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)-x*(5-x^2)*diff(y(x),x)-(7+25*x^2)*y(x)=0,y(x),type='series
```

$$y(x) = c_1 x^7 \left(1 - \frac{6}{5}x^2 + \frac{7}{5}x^4 + O(x^6) \right) + \frac{c_2(-203212800 + 406425600x^2 - 609638400x^4 + O(x^6))}{x}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 40

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]-x*(5-x^2)*y'[x]-(7+25*x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(3x^3 - 2x + \frac{1}{x} \right) + c_2 \left(\frac{7x^{11}}{5} - \frac{6x^9}{5} + x^7 \right)$$

16.42 problem 38

Internal problem ID [1454]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(x^2 + 1)y'' + x(2x^2 + 5)y' - 21y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

Order:=6;

```
dsolve(x^2*(1+x^2)*diff(y(x),x$2)+x*(5+2*x^2)*diff(y(x),x)-21*y(x)=0,y(x),type='series',x=0)
```

$$y(x) = c_1 x^3 \left(1 - \frac{1}{2}x^2 + \frac{15}{56}x^4 + O(x^6) \right) + \frac{c_2(-1316818944000 - 3456649728000x^2 - 2880541440000x^4 + O(x^6))}{x^7}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 46

```
AsymptoticDSolveValue[x^2*(1+x^2)*y'[x]+x*(5+2*x^2)*y'[x]-21*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{x^7} + \frac{21}{8x^5} + \frac{35}{16x^3} \right) + c_2 \left(\frac{15x^7}{56} - \frac{x^5}{2} + x^3 \right)$$

16.43 problem 39

Internal problem ID [1455]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2(1 + 2x^2)y'' - x(x^2 + 3)y' - 2yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 54

Order:=6;

```
dsolve(x^2*(1+2*x^2)*diff(y(x),x$2)-x*(3+x^2)*diff(y(x),x)-2*x*y(x)=0,y(x),type='series',x=0
```

$$\begin{aligned} y(x) = & c_1 x^4 \left(1 + \frac{2}{5}x - \frac{8}{5}x^2 - \frac{86}{105}x^3 + \frac{445}{168}x^4 + \frac{9571}{6300}x^5 + O(x^6) \right) \\ & + c_2 \left(\ln(x) \left(24x^4 + \frac{48}{5}x^5 + O(x^6) \right) \right. \\ & \left. + \left(-144 + 96x - 48x^2 + 210x^4 + \frac{1812}{25}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x^2*(1+2*x^2)*y'[x]-x*(3+x^2)*y'[x]-2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{12}(3x^4 + 4x^2 - 8x + 12) - \frac{1}{6}x^4 \log(x) \right) + c_2 \left(\frac{445x^8}{168} - \frac{86x^7}{105} - \frac{8x^6}{5} + \frac{2x^5}{5} + x^4 \right)$$

16.44 problem 40

Internal problem ID [1456]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 7 Series Solutions of Linear Second Equations. 7.6 THE METHOD OF FROBENIUS III. Exercises 7.7. Page 389

Problem number: 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$4x^2(x^2 + 1)y'' + 4x(x^2 + 2)y' - (x^2 + 15)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

Order:=6;

```
dsolve(4*x^2*(1+x^2)*diff(y(x),x$2)+4*x*(2+x^2)*diff(y(x),x)-(15+x^2)*y(x)=0,y(x),type='series')
```

$$y(x) = \frac{c_1 x^4 \left(1 - \frac{1}{6}x^2 + \frac{1}{16}x^4 + O(x^6)\right) + c_2 (-144 - 216x^2 - 54x^4 + O(x^6))}{x^{\frac{5}{2}}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 58

```
AsymptoticDSolveValue[4*x^2*(1+x^2)*y'[x]+4*x*(2+x^2)*y'[x]-(15+x^2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{3x^{3/2}}{8} + \frac{1}{x^{5/2}} + \frac{3}{2\sqrt{x}} \right) + c_2 \left(\frac{x^{11/2}}{16} - \frac{x^{7/2}}{6} + x^{3/2} \right)$$

17 Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

17.1	problem section 9.1, problem 2	658
17.2	problem section 9.1, problem 3	659
17.3	problem section 9.1, problem 5(b) 1	660
17.4	problem section 9.1, problem 5(b) 2	661
17.5	problem section 9.1, problem 5(b) 3	662
17.6	problem section 9.1, problem 5(b)	663
17.7	problem section 9.1, problem 6(a)	664
17.8	problem section 9.1, problem 6(b)	665

17.1 problem section 9.1, problem 2

Internal problem ID [1457]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 2.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' - x^2 y'' - 2y'x + 6y = 0$$

With initial conditions

$$[y(-1) = -4, y'(-1) = -14, y''(-1) = -20]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+6*y(x)=0,y(-1) = -4, D(y)(-1)
```

$$y(x) = 3x^3 + \frac{22x^2}{3} + \frac{25}{3x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 24

```
DSolve[{x^3*y'''[x]-x^2*y''[x]-2*x*y'[x]+6*y[x]==0,{y[-1]==-4,y'[-1]==-14,y''[-1]==-20}},y[x]
```

$$y(x) \rightarrow \frac{9x^4 + 22x^3 + 25}{3x}$$

17.2 problem section 9.1, problem 3

Internal problem ID [1458]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 3.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y''' - 7y'' - y' + 6y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = -6, y''(0) = 10, y'''(0) = -36]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$4)+diff(y(x),x$3)-7*diff(y(x),x$2)-diff(y(x),x)+6*y(x)=0,y(0) = 5, D(y)
```

$$y(x) = (-e^{5x} + 2e^{4x} + 3e^{2x} + 1) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 30

```
DSolve[{y''''[x]+y'''[x]-7*y''[x]-y'[x]+6*y[x]==0,{y[0]==5,y'[0]==-6,y''[0]==10,y'''[0]==-36
```

$$y(x) \rightarrow e^{-3x} + 3e^{-x} + 2e^x - e^{2x}$$

17.3 problem section 9.1, problem 5(b) 1

Internal problem ID [1459]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 5(b) 1.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' - x^2 y'' - 2y'x + 6y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0, y''(1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([x^3*dif(y(x),x$3)-x^2*dif(y(x),x$2)-2*x*dif(y(x),x)+6*y(x)=0,y(1) = 1, D(y)(1) =
```

$$y(x) = -\frac{x^3}{2} + x^2 + \frac{1}{2x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 23

```
DSolve[{x^3*y'''[x]-x^2*y''[x]-2*x*y'[x]+6*y[x]==0,{y[1]==1,y'[1]==0,y''[1]==0}},y[x],x,Incl
```

$$y(x) \rightarrow -\frac{x^3}{2} + x^2 + \frac{1}{2x}$$

17.4 problem section 9.1, problem 5(b) 2

Internal problem ID [1460]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 5(b) 2.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' - x^2 y'' - 2y'x + 6y = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1, y''(1) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+6*y(x)=0,y(1) = 0, D(y)(1) =
```

$$y(x) = \frac{x^3 - 1}{3x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[{x^3*y'''[x]-x^2*y''[x]-2*x*y'[x]+6*y[x]==0,{y[1]==0,y'[1]==1,y''[1]==0}},y[x],x,Incl
```

$$y(x) \rightarrow \frac{x^3 - 1}{3x}$$

17.5 problem section 9.1, problem 5(b) 3

Internal problem ID [1461]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 5(b) 3.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' - x^2 y'' - 2y'x + 6y = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 0, y''(1) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 20

```
dsolve([x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+6*y(x)=0,y(1) = 0, D(y)(1) =
```

$$y(x) = \frac{3x^4 - 4x^3 + 1}{12x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 24

```
DSolve[{x^3*y'''[x]-x^2*y''[x]-2*x*y'[x]+6*y[x]==0,{y[1]==0,y'[1]==0,y''[1]==1}},y[x],x,Incl
```

$$y(x) \rightarrow \frac{3x^4 - 4x^3 + 1}{12x}$$

17.6 problem section 9.1, problem 5(b)

Internal problem ID [1462]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 5(b).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' - x^2 y'' - 2y'x + 6y = 0$$

With initial conditions

$$[y(1) = k_0, y'(1) = k_1, y''(1) = k_2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 40

```
dsolve([x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+6*y(x)=0,y(1) = k__0, D(y)(1)
```

$$y(x) = \frac{3(-2k_0 + k_2)x^4 + 4(k_1 + 3k_0 - k_2)x^3 + 6k_0 - 4k_1 + k_2}{12x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 47

```
DSolve[{x^3*y'''[x]-x^2*y''[x]-2*x*y'[x]+6*y[x]==0,{y[1]==k0,y'[1]==k1,y''[1]==k2}},y[x],x,D
```

$$y(x) \rightarrow \frac{-6k_0(x^4 - 2x^3 - 1) + 4k_1(x^3 - 1) + 3k_2x^4 - 4k_2x^3 + k_2}{12x}$$

17.7 problem section 9.1, problem 6(a)

Internal problem ID [1463]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 6(a).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + y'' - y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$3)+diff(y(x),x$2)-diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_3x + c_2)e^{-x} + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]+y''[x]-y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2x + c_3e^{2x} + c_1)$$

17.8 problem section 9.1, problem 6(b)

Internal problem ID [1464]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.1. Page 471

Problem number: section 9.1, problem 6(b).

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 3y'' + 7y' - 5y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+7*diff(y(x),x)-5*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x(c_1 + \sin(2x)c_2 + \cos(2x)c_3)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]-3*y''[x]+7*y'[x]-5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(2x) + c_1 \sin(2x) + c_3)$$

18 Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient.

Page 483

18.1	problem section 9.2, problem 1	667
18.2	problem section 9.2, problem 2	668
18.3	problem section 9.2, problem 3	669
18.4	problem section 9.2, problem 4	670
18.5	problem section 9.2, problem 5	671
18.6	problem section 9.2, problem 6	672
18.7	problem section 9.2, problem 7	673
18.8	problem section 9.2, problem 8	674
18.9	problem section 9.2, problem 9	675
18.10	problem section 9.2, problem 10	676
18.11	problem section 9.2, problem 11	677
18.12	problem section 9.2, problem 12	678
18.13	problem section 9.2, problem 13	679
18.14	problem section 9.2, problem 14	680
18.15	problem section 9.2, problem 15	681
18.16	problem section 9.2, problem 16	682
18.17	problem section 9.2, problem 17	683
18.18	problem section 9.2, problem 18	684
18.19	problem section 9.2, problem 19	685
18.20	problem section 9.2, problem 20	686
18.21	problem section 9.2, problem 21	687
18.22	problem section 9.2, problem 22	688
18.23	problem section 9.2, problem 23	689
18.24	problem section 9.2, problem 24	690
18.25	problem section 9.2, problem 25	691
18.26	problem section 9.2, problem 26	692
18.27	problem section 9.2, problem 27	693
18.28	problem section 9.2, problem 43(a)	694
18.29	problem section 9.2, problem 43(b)	695
18.30	problem section 9.2, problem 43(c)	696
18.31	problem section 9.2, problem 43(d)	697
18.32	problem section 9.2, problem 43(e)	698
18.33	problem section 9.2, problem 43(g)	699

18.1 problem section 9.2, problem 1

Internal problem ID [1465]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 1.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 3y'' + 3y' - y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+3*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x (c_3 x^2 + c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 21

```
DSolve[y'''[x]-3*y''[x]+3*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (x(c_3 x + c_2) + c_1)$$

18.2 problem section 9.2, problem 2

Internal problem ID [1466]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 2.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 8y'' - 9y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)+8*diff(y(x),x$2)-9*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + c_2e^x + c_3 \sin(3x) + c_4 \cos(3x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y''''[x]+8*y''[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3e^{-x} + c_4e^x + c_1 \cos(3x) + c_2 \sin(3x)$$

18.3 problem section 9.2, problem 3

Internal problem ID [1467]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 3.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y'' + 16y' - 16y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)+16*diff(y(x),x)-16*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + \sin(4x) c_2 + c_3 \cos(4x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]-y''[x]+16*y'[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 e^x + c_1 \cos(4x) + c_2 \sin(4x)$$

18.4 problem section 9.2, problem 4

Internal problem ID [1468]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 4.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$2y''' + 3y'' - 2y' - 3y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(2*diff(y(x),x$3)+3*diff(y(x),x$2)-2*diff(y(x),x)-3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(c_3 e^{\frac{5x}{2}} + c_2 e^{\frac{x}{2}} + c_1 \right) e^{-\frac{3x}{2}}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[2*y'''[x]+3*y''[x]-2*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-3x/2} + c_2 e^{-x} + c_3 e^x$$

18.5 problem section 9.2, problem 5

Internal problem ID [1469]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 5.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 5y'' + 9y' + 5y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$3)+5*diff(y(x),x$2)+9*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + c_2e^{-2x} \sin(x) + c_3e^{-2x} \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 28

```
DSolve[y'''[x]+5*y''[x]+9*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_3e^x + c_2 \cos(x) + c_1 \sin(x))$$

18.6 problem section 9.2, problem 6

Internal problem ID [1470]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 6.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$4y''' - 8y'' + 5y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(4*diff(y(x),x$3)-8*diff(y(x),x$2)+5*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_3x + c_2) e^{\frac{x}{2}} + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[4*y'''[x]-8*y''[x]+5*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2}(c_2x + c_3e^{x/2} + c_1)$$

18.7 problem section 9.2, problem 7

Internal problem ID [1471]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 7.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$27y''' + 27y'' + 9y' + y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(27*diff(y(x),x$3)+27*diff(y(x),x$2)+9*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{3}}(c_3x^2 + c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[27*y'''[x]+27*y''[x]+9*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/3}(x(c_3x + c_2) + c_1)$$

18.8 problem section 9.2, problem 8

Internal problem ID [1472]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 8.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y'' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$4)+diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2x + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 24

```
DSolve[y''''[x]+y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_4x - c_1 \cos(x) - c_2 \sin(x) + c_3$$

18.9 problem section 9.2, problem 9

Internal problem ID [1473]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 9.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-16*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} + e^{-2x} c_2 + c_3 \sin(2x) + c_4 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y''''[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x} + c_3 e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

18.10 problem section 9.2, problem 10

Internal problem ID [1474]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 10.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 12y'' + 36y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)+12*diff(y(x),x$2)+36*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_4x + c_2) \cos(\sqrt{6}x) + \sin(\sqrt{6}x) (c_3x + c_1)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 38

```
DSolve[y''''[x]+12*y''[x]+36*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (c_2x + c_1) \cos(\sqrt{6}x) + (c_4x + c_3) \sin(\sqrt{6}x)$$

18.11 problem section 9.2, problem 11

Internal problem ID [1475]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 11.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$16y'''' - 72y'' + 81y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(16*diff(y(x),x$4)-72*diff(y(x),x$2)+81*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{3x}{2}}(c_2x + c_1) + e^{\frac{3x}{2}}(c_4x + c_3)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 37

```
DSolve[16*y''''[x]-72*y''[x]+81*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x/2}(c_3e^{3x} + x(c_4e^{3x} + c_2) + c_1)$$

18.12 problem section 9.2, problem 12

Internal problem ID [1476]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 12.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$6y'''' + 5y''' + 7y'' + 5y' + y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(6*diff(y(x),x$4)+5*diff(y(x),x$3)+7*diff(y(x),x$2)+5*diff(y(x),x)+y(x)=0,y(x), singularities)
```

$$y(x) = c_1 e^{-\frac{x}{2}} + c_2 e^{-\frac{x}{3}} + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 37

```
DSolve[6*y''''[x]+5*y'''[x]+7*y''[x]+5*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} (c_3 e^{x/6} + c_4) + c_1 \cos(x) + c_2 \sin(x)$$

18.13 problem section 9.2, problem 13

Internal problem ID [1477]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 13.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$4y'''' + 12y''' + 3y'' - 13y' - 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(4*diff(y(x),x$4)+12*diff(y(x),x$3)+3*diff(y(x),x$2)-13*diff(y(x),x)-6*y(x)=0,y(x), si
```

$$y(x) = \left(c_4 e^{3x} + c_3 e^{\frac{3x}{2}} + c_1 e^{\frac{x}{2}} + c_2 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
DSolve[4*y''''[x]+12*y'''[x]+3*y''[x]-13*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow e^{-2x} (c_1 e^{x/2} + c_2 e^{3x/2} + c_4 e^{3x} + c_3)$$

18.14 problem section 9.2, problem 14

Internal problem ID [1478]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 14.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 4y''' + 7y'' - 6y' + 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+7*diff(y(x),x$2)-6*diff(y(x),x)+2*y(x))=0,y(x), singso
```

$$y(x) = e^x(c_1 + c_2x + c_3 \sin(x) + c_4 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 26

```
DSolve[y''''[x]-4*y'''[x]+7*y''[x]-6*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^x(c_4x + c_2 \cos(x) + c_1 \sin(x) + c_3)$$

18.15 problem section 9.2, problem 15

Internal problem ID [1479]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 15.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 2y'' + 4y' - 8y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = -2, y''(0) = 2]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$3)-2*diff(y(x),x$2)+4*diff(y(x),x)-8*y(x)=0,y(0) = 2, D(y)(0) = -2, (D@@
```

$$y(x) = \frac{5e^{2x}}{4} - \frac{9\sin(2x)}{4} + \frac{3\cos(2x)}{4}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 21

```
DSolve[{y'''[x]-2*y''[x]+4*y'[x]-8*y[x]==0,{y[0]==2,y'[0]==-2,y''[0]==0}},y[x],x,IncludeSing
```

$$y(x) \rightarrow e^{2x} - 2\sin(2x) + \cos(2x)$$

18.16 problem section 9.2, problem 16

Internal problem ID [1480]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 16.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 3y'' - y' - 3y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 14, y''(0) = -40]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$3)+3*diff(y(x),x$2)-diff(y(x),x)-3*y(x)=0,y(0) = 0, D(y)(0) = 14, (D@@2)
```

$$y(x) = (2e^{4x} + 3e^{2x} - 5)e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[{y'''[x]+3*y''[x]-y'[x]-3*y[x]==0,{y[0]==0,y'[0]==14,y''[0]==-40}},y[x],x,IncludeSing
```

$$y(x) \rightarrow -5e^{-3x} + 3e^{-x} + 2e^x$$

18.17 problem section 9.2, problem 17

Internal problem ID [1481]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 17.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y'' - y' + y = 0$$

With initial conditions

$$[y(0) = -2, y'(0) = 9, y''(0) = 4]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$3)-diff(y(x),x$2)-diff(y(x),x)+y(x)=0,y(0) = -2, D(y)(0) = 9, (D@@2)(y)(0) = 4],y(x),x)
```

$$y(x) = -4e^{-x} + (3x + 2)e^x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 22

```
DSolve[{y'''[x]-y''[x]-y'[x]+y[x]==0,{y[0]==-2,y'[0]==9,y''[0]==4}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^x(3x + 2) - 4e^{-x}$$

18.18 problem section 9.2, problem 18

Internal problem ID [1482]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 18.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 2y' - 4y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = 3, y''(0) = 22]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$3)-2*diff(y(x),x)-4*y(x)=0,y(0) = 6, D(y)(0) = 3, (D@@2)(y)(0) = 22],y(x)
```

$$y(x) = 4e^{2x} + (-3\sin(x) + 2\cos(x))e^{-x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 27

```
DSolve[{y'''[x]-2*y'[x]-4*y[x]==0,{y[0]==6,y'[0]==3,y''[0]==22}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow e^{-x}(4e^{3x} - 3\sin(x) + 2\cos(x))$$

18.19 problem section 9.2, problem 19

Internal problem ID [1483]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 19.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$3y''' - y'' - 7y' + 5y = 0$$

With initial conditions

$$\left[y(0) = \frac{14}{5}, y'(0) = 0, y''(0) = 10 \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([3*diff(y(x),x$3)-diff(y(x),x$2)-7*diff(y(x),x)+5*y(x)=0,y(0) = 14/5, D(y)(0) = 0, D
```

$$y(x) = e^{-\frac{5x}{3}} \left(\frac{9}{5} + (2x + 1) e^{\frac{8x}{3}} \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 26

```
DSolve[{3*y'''[x]-y''[x]-7*y'[x]+5*y[x]==0,{y[0]==14/5,y'[0]==0,y''[0]==10}},y[x],x,IncludeS
```

$$y(x) \rightarrow e^x(2x + 1) + \frac{9}{5}e^{-5x/3}$$

18.20 problem section 9.2, problem 20

Internal problem ID [1484]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 20.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 6y'' + 12y' - 8y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1, y''(0) = -4]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$3)-6*diff(y(x),x$2)+12*diff(y(x),x)-8*y(x)=0,y(0) = 1, D(y)(0) = -1, (D
```

$$y(x) = e^{2x}(2x^2 - 3x + 1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 21

```
DSolve[{y'''[x]-6*y''[x]+12*y'[x]-8*y[x]==0,{y[0]==1,y'[0]==-1,y''[0]==-4}},y[x],x,IncludeSi
```

$$y(x) \rightarrow e^{2x}(2x^2 - 3x + 1)$$

18.21 problem section 9.2, problem 21

Internal problem ID [1485]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 21.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$2y''' - 11y'' + 12y' + 9y = 0$$

With initial conditions

$$[y(0) = 6, y'(0) = 3, y''(0) = 13]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve([2*diff(y(x),x$3)-11*diff(y(x),x$2)+12*diff(y(x),x)+9*y(x)=0,y(0) = 6, D(y)(0) = 3, (
```

$$y(x) = 4e^{-\frac{x}{2}} + (2 - x)e^{3x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[{2*y'''[x]-11*y''[x]+12*y'[x]+9*y[x]==0,{y[0]==6,y'[0]==3,y''[0]==13}},y[x],x,Include
```

$$y(x) \rightarrow 4e^{-x/2} - e^{3x}(x - 2)$$

18.22 problem section 9.2, problem 22

Internal problem ID [1486]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 22.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$8y''' - 4y'' - 2y' + y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = -3, y''(0) = -1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve([8*diff(y(x),x$3)-4*diff(y(x),x$2)-2*diff(y(x),x)+y(x)=0,y(0) = 4, D(y)(0) = -3, (D@@
```

$$y(x) = 3e^{-\frac{x}{2}} + e^{\frac{x}{2}} - 2e^{\frac{x}{2}}x$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 57

```
DSolve[{8*y'''[x]-4*y''[x]-2*y'[x]-2*y[x]==0,{y[0]==4,y'[0]==-3,y''[0]==-1}},y[x],x,IncludeS
```

$$y(x) \rightarrow -\frac{2}{21}e^{-x/4} \left(9e^{5x/4} + 13\sqrt{3} \sin\left(\frac{\sqrt{3}x}{4}\right) - 51 \cos\left(\frac{\sqrt{3}x}{4}\right) \right)$$

18.23 problem section 9.2, problem 23

Internal problem ID [1487]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 23.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 2, y''(0) = -2, y'''(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$4)-16*y(x)=0,y(0) = 2, D(y)(0) = 2, (D@@2)(y)(0) = -2, (D@@3)(y)(0) = 0],y(x),x)
```

$$y(x) = \frac{5e^{2x}}{8} + \frac{e^{-2x}}{8} + \frac{\sin(2x)}{2} + \frac{5\cos(2x)}{4}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[{y''''[x]-16*y[x]==0,{y[0]==2,y'[0]==2,y''[0]==-2,y'''[0]==0}},y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{1}{8}(e^{-2x} + 5e^{2x} + 4\sin(2x) + 10\cos(2x))$$

18.24 problem section 9.2, problem 24

Internal problem ID [1488]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 24.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 6y'''' + 7y'' + 6y' - 8y = 0$$

With initial conditions

$$[y(0) = -2, y'(0) = -8, y''(0) = -14, y'''(0) = -62]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 25

```
dsolve([diff(y(x),x$4)-6*diff(y(x),x$3)+7*diff(y(x),x$2)+6*diff(y(x),x)-8*y(x)=0,y(0) = -2,
```

$$y(x) = e^{2x} - e^{4x} + 2e^{-x} - 4e^x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[{y''''[x]-6*y''''[x]+7*y''[x]+6*y'[x]-8*y[x]==0,{y[0]==-2,y'[0]==-8,y''[0]==-14,y'''[0]
```

$$y(x) \rightarrow 2e^{-x} - 4e^x + e^{2x} - e^{4x}$$

18.25 problem section 9.2, problem 25

Internal problem ID [1489]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 25.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$4y'''' - 13y'' + 9y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 3, y''(0) = 1, y'''(0) = 3]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve([4*diff(y(x),x$4)-13*diff(y(x),x$2)+9*y(x)=0,y(0) = 1, D(y)(0) = 3, (D@@2)(y)(0) = 1,
```

$$y(x) = -e^{-x} + 2e^x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 18

```
DSolve[{4*y''''[x]-13*y''[x]+9*y[x]==0,{y[0]==1,y'[0]==3,y''[0]==1,y'''[0]==3}},y[x],x,Inclu
```

$$y(x) \rightarrow 2e^x - e^{-x}$$

18.26 problem section 9.2, problem 26

Internal problem ID [1490]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 26.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 2y''' - 2y'' - 8y' - 8y = 0$$

With initial conditions

$$[y(0) = 5, y'(0) = -2, y''(0) = 6, y'''(0) = 8]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve([diff(y(x),x$4)+2*diff(y(x),x$3)-2*diff(y(x),x$2)-8*diff(y(x),x)-8*y(x))=0,y(0) = 5, D
```

$$y(x) = (1 + (\sin(x) + 3 \cos(x)) e^x + e^{4x}) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 31

```
DSolve[{y''''[x]+2*y'''[x]-2*y''[x]-8*y'[x]-8*y[x]==0,{y[0]==5,y'[0]==-2,y''[0]==6,y'''[0]==
```

$$y(x) \rightarrow e^{-2x} (e^{4x} + e^x \sin(x) + 3e^x \cos(x) + 1)$$

18.27 problem section 9.2, problem 27

Internal problem ID [1491]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 27.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$4y'''' + 8y''' + 19y'' + 32y' + 12y = 0$$

With initial conditions

$$\left[y(0) = 3, y'(0) = -3, y''(0) = -\frac{7}{2}, y'''(0) = \frac{31}{4} \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve([4*diff(y(x),x$4)+8*diff(y(x),x$3)+19*diff(y(x),x$2)+32*diff(y(x),x)+12*y(x)=0,y(0) =
```

$$y(x) = 2e^{-\frac{x}{2}} - \sin(2x) + \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 31

```
DSolve[{y''''[x]+2*y'''[x]-2*y''[x]-8*y'[x]-8*y[x]==0,{y[0]==5,y'[0]==-2,y''[0]==6,y'''[0]==
```

$$y(x) \rightarrow e^{-2x}(e^{4x} + e^x \sin(x) + 3e^x \cos(x) + 1)$$

18.28 problem section 9.2, problem 43(a)

Internal problem ID [1492]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(a).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$4)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}c_1 + c_2e^x + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 30

```
DSolve[y''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x + c_3e^{-x} + c_2 \cos(x) + c_4 \sin(x)$$

18.29 problem section 9.2, problem 43(b)

Internal problem ID [1493]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(b).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 61

```
dsolve(diff(y(x),x$4)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(-e^{-\frac{\sqrt{2}x}{2}} c_1 - e^{\frac{\sqrt{2}x}{2}} c_2\right) \sin\left(\frac{\sqrt{2}x}{2}\right) + \left(c_3 e^{-\frac{\sqrt{2}x}{2}} + c_4 e^{\frac{\sqrt{2}x}{2}}\right) \cos\left(\frac{\sqrt{2}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 65

```
DSolve[y''''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{x}{\sqrt{2}}}\left(\left(c_1 e^{\sqrt{2}x} + c_2\right) \cos\left(\frac{x}{\sqrt{2}}\right) + \left(c_4 e^{\sqrt{2}x} + c_3\right) \sin\left(\frac{x}{\sqrt{2}}\right)\right)$$

18.30 problem section 9.2, problem 43(c)

Internal problem ID [1494]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(c).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 64y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(diff(y(x),x$4)+64*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_2 e^{2x} + c_4 e^{-2x}) \cos(2x) + \sin(2x) (c_1 e^{2x} + c_3 e^{-2x})$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 44

```
DSolve[y''''[x]+64*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} ((c_4 e^{4x} + c_1) \cos(2x) + (c_3 e^{4x} + c_2) \sin(2x))$$

18.31 problem section 9.2, problem 43(d)

Internal problem ID [1495]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(d).

ODE order: 6.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(6)} - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve(diff(y(x),x$6)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(\left(e^{\frac{x}{2}} c_6 + e^{\frac{3x}{2}} c_4 \right) \cos \left(\frac{\sqrt{3}x}{2} \right) + \left(e^{\frac{x}{2}} c_5 + c_3 e^{\frac{3x}{2}} \right) \sin \left(\frac{\sqrt{3}x}{2} \right) + c_2 e^{2x} + c_1 \right) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 78

```
DSolve[y''''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_1 e^{2x} + e^{x/2} (c_2 e^x + c_3) \cos \left(\frac{\sqrt{3}x}{2} \right) + e^{x/2} (c_6 e^x + c_5) \sin \left(\frac{\sqrt{3}x}{2} \right) + c_4 \right)$$

18.32 problem section 9.2, problem 43(e)

Internal problem ID [1496]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(e).

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 64y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(diff(y(x),x$4)+64*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_2 e^{2x} + c_4 e^{-2x}) \cos(2x) + \sin(2x) (c_1 e^{2x} + c_3 e^{-2x})$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 44

```
DSolve[y''''[x]+64*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} ((c_4 e^{4x} + c_1) \cos(2x) + (c_3 e^{4x} + c_2) \sin(2x))$$

18.33 problem section 9.2, problem 43(g)

Internal problem ID [1497]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.2. constant coefficient. Page 483

Problem number: section 9.2, problem 43(g).

ODE order: 5.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y^{(5)} + y'''' + y''' + y'' + y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 53

```
dsolve(diff(y(x),x$5)+diff(y(x),x$4)+diff(y(x),x$3)+diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(x),
```

$$y(x) = e^{-x} \left(\left(e^{\frac{x}{2}} c_5 + c_3 e^{\frac{3x}{2}} \right) \cos \left(\frac{\sqrt{3}x}{2} \right) + \left(c_2 e^{\frac{3x}{2}} + c_4 e^{\frac{x}{2}} \right) \sin \left(\frac{\sqrt{3}x}{2} \right) + c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 70

```
DSolve[y''''''[x]+y''''[x]+y'''[x]+y''[x]+y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow e^{-x} \left(e^{x/2} (c_3 e^x + c_2) \cos \left(\frac{\sqrt{3}x}{2} \right) + e^{x/2} (c_4 e^x + c_1) \sin \left(\frac{\sqrt{3}x}{2} \right) + c_5 \right)$$

19 Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

19.1	problem section 9.3, problem 1	703
19.2	problem section 9.3, problem 2	704
19.3	problem section 9.3, problem 3	705
19.4	problem section 9.3, problem 4	706
19.5	problem section 9.3, problem 5	707
19.6	problem section 9.3, problem 6	708
19.7	problem section 9.3, problem 7	709
19.8	problem section 9.3, problem 8	710
19.9	problem section 9.3, problem 9	711
19.10	problem section 9.3, problem 10	712
19.11	problem section 9.3, problem 11	713
19.12	problem section 9.3, problem 12	714
19.13	problem section 9.3, problem 13	715
19.14	problem section 9.3, problem 14	716
19.15	problem section 9.3, problem 15	717
19.16	problem section 9.3, problem 16	718
19.17	problem section 9.3, problem 17	719
19.18	problem section 9.3, problem 18	720
19.19	problem section 9.3, problem 19	721
19.20	problem section 9.3, problem 20	722
19.21	problem section 9.3, problem 21	723
19.22	problem section 9.3, problem 22	724
19.23	problem section 9.3, problem 23	725
19.24	problem section 9.3, problem 24	726
19.25	problem section 9.3, problem 25	727
19.26	problem section 9.3, problem 26	728
19.27	problem section 9.3, problem 27	729
19.28	problem section 9.3, problem 28	730
19.29	problem section 9.3, problem 29	731
19.30	problem section 9.3, problem 30	732
19.31	problem section 9.3, problem 31	733
19.32	problem section 9.3, problem 32	734
19.33	problem section 9.3, problem 33	735

19.34	problem section 9.3, problem 34	736
19.35	problem section 9.3, problem 35	737
19.36	problem section 9.3, problem 36	738
19.37	problem section 9.3, problem 37	739
19.38	problem section 9.3, problem 38	740
19.39	problem section 9.3, problem 39	741
19.40	problem section 9.3, problem 40	742
19.41	problem section 9.3, problem 41	743
19.42	problem section 9.3, problem 42	744
19.43	problem section 9.3, problem 43	745
19.44	problem section 9.3, problem 44	746
19.45	problem section 9.3, problem 45	747
19.46	problem section 9.3, problem 46	748
19.47	problem section 9.3, problem 47	749
19.48	problem section 9.3, problem 48	750
19.49	problem section 9.3, problem 49	751
19.50	problem section 9.3, problem 50	752
19.51	problem section 9.3, problem 51	753
19.52	problem section 9.3, problem 52	754
19.53	problem section 9.3, problem 53	755
19.54	problem section 9.3, problem 54	756
19.55	problem section 9.3, problem 55	757
19.56	problem section 9.3, problem 56	758
19.57	problem section 9.3, problem 57	759
19.58	problem section 9.3, problem 58	760
19.59	problem section 9.3, problem 59	761
19.60	problem section 9.3, problem 60	762
19.61	problem section 9.3, problem 61	763
19.62	problem section 9.3, problem 62	764
19.63	problem section 9.3, problem 63	765
19.64	problem section 9.3, problem 64	766
19.65	problem section 9.3, problem 65	767
19.66	problem section 9.3, problem 66	768
19.67	problem section 9.3, problem 67	769
19.68	problem section 9.3, problem 68	770
19.69	problem section 9.3, problem 69	771
19.70	problem section 9.3, problem 70	772
19.71	problem section 9.3, problem 71	773
19.72	problem section 9.3, problem 72	774

19.73	problem section 9.3, problem 73	775
19.74	problem section 9.3, problem 74	776

19.1 problem section 9.3, problem 1

Internal problem ID [1498]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 1.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 6y'' + 11y' - 6y = -e^x(-24x^2 + 76x + 4)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+11*diff(y(x),x)-6*y(x)=-exp(x)*(4+76*x-24*x^2),y(x),
```

$$y(x) = e^x(c_3e^{2x} + 4x^3 + c_2e^x - x^2 + c_1 - 17x)$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 47

```
DSolve[y'''[x]-6*y''[x]+11*y'[x]-6*y[x]==-Exp[x]*(4+76*x-24*x^2),y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{1}{2}e^x(8x^3 - 2x^2 - 34x + 2c_2e^x + 2c_3e^{2x} - 49 + 2c_1)$$

19.2 problem section 9.3, problem 2

Internal problem ID [1499]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 2.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 2y'' - 5y' + 6y = e^{-3x}(6x^2 - 23x + 32)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)-2*diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=exp(-3*x)*(32-23*x+6*x^2),y(x),
```

$$y(x) = \frac{(4c_3e^{6x} + 4e^{4x}c_1 + 4c_2e^x - x^2 + x - 3)e^{-3x}}{4}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 45

```
DSolve[y'''[x]-2*y''[x]-5*y'[x]+6*y[x]==Exp[-3*x]*(32-23*x+6*x^2),y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow -\frac{1}{4}e^{-3x}(x^2 - x + 3) + c_1e^{-2x} + c_2e^x + c_3e^{3x}$$

19.3 problem section 9.3, problem 3

Internal problem ID [1500]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 3.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$4y''' + 8y'' - y' - 2y = -e^x(6x^2 + 45x + 4)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(4*diff(y(x),x$3)+8*diff(y(x),x$2)-diff(y(x),x)-2*y(x)=-exp(x)*(4+45*x+6*x^2),y(x), si
```

$$y(x) = \frac{\left(-18x^2e^{3x} - 27xe^{3x} + 149e^{3x} + 27c_3e^{\frac{5x}{2}} + 27c_2e^{\frac{3x}{2}} + 27c_1\right)e^{-2x}}{27}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 52

```
DSolve[4*y'''[x]+8*y''[x]-y'[x]-2*y[x]==-Exp[x]*(4+45*x+6*x^2),y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow e^x \left(-\frac{2x^2}{3} - x + \frac{149}{27} \right) + c_1 e^{-x/2} + c_2 e^{x/2} + c_3 e^{-2x}$$

19.4 problem section 9.3, problem 4

Internal problem ID [1501]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 4.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 3y'' - y' - 3y = e^{-2x}(3x^2 - 17x + 2)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)-diff(y(x),x)-3*y(x)=exp(-2*x)*(2-17*x+3*x^2),y(x), si
```

$$y(x) = (c_3 e^{2x} + e^{4x} c_1 + (x^2 - 5x + 1) e^x + c_2) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 42

```
DSolve[y'''[x]+3*y''[x]-y'[x]-3*y[x]==Exp[-2*x]*(2-17*x+3*x^2),y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow e^{-3x}(e^x(x^2 - 5x + 1) + c_2 e^{2x} + c_3 e^{4x} + c_1)$$

19.5 problem section 9.3, problem 5

Internal problem ID [1502]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 5.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 3y'' - y' - 3y = e^x(16x^3 + 24x^2 + 2x - 1)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)-diff(y(x),x)-3*y(x)=exp(x)*(-1+2*x+24*x^2+16*x^3),y(x)
```

$$y(x) = \frac{((x^4 - x^3 + x^2 + 2c_1 - x)e^{4x} + 2c_3e^{2x} + 2c_2)e^{-3x}}{2}$$

✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 53

```
DSolve[y'''[x]+3*y''[x]-y'[x]-3*y[x]==Exp[x]*(-1+2*x+24*x^2+16*x^3),y[x],x,IncludeSingularSo
```

$$y(x) \rightarrow \frac{1}{4}e^x(2x^4 - 2x^3 + 2x^2 - 2x + 1 + 4c_3) + c_1e^{-3x} + c_2e^{-x}$$

19.6 problem section 9.3, problem 6

Internal problem ID [1503]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 6.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + y'' - 2y = e^x(15x^2 + 34x + 14)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$3)+diff(y(x),x$2)-2*y(x)=exp(x)*(14+34*x+15*x^2),y(x), singsol=all)
```

$$y(x) = (c_2 \cos(x) + c_3 \sin(x)) e^{-x} + e^x(x^3 + x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.159 (sec). Leaf size: 49

```
DSolve[y'''[x]+y''[x]-2*y[x]==Exp[x]*(14+34*x+15*x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5} e^{-x} (e^{2x} (5x^3 + 5x^2 - 2 + 5c_3) + 5c_2 \cos(x) + 5c_1 \sin(x))$$

19.7 problem section 9.3, problem 7

Internal problem ID [1504]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 7.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$4y''' + 8y'' - y' - 2y = -e^{-2x}(1 - 15x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(4*diff(y(x),x$3)+8*diff(y(x),x$2)-diff(y(x),x)-2*y(x)=-exp(-2*x)*(1-15*x),y(x),sings
```

$$y(x) = \frac{\left(2c_3e^{\frac{5x}{2}} + 2c_2e^{\frac{3x}{2}} + x^2 + 2c_1 + 2x\right)e^{-2x}}{2}$$

✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 53

```
DSolve[4*y'''[x]+8*y''[x]-y'[x]-2*y[x]==-Exp[-2*x]*(1-15*x),y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{10}e^{-2x}(5x^2 + 10x + 2(5c_1e^{3x/2} + 5c_2e^{5x/2} + 4 + 5c_3))$$

19.8 problem section 9.3, problem 8

Internal problem ID [1505]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 8.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' - y' + y = -e^x(7 + 6x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$3)-diff(y(x),x$2)-diff(y(x),x)+y(x)=-exp(x)*(7+6*x),y(x), singsol=all)
```

$$y(x) = e^{-x}c_2 - \frac{e^x(x^3 - 2c_3x + 2x^2 - 2c_1)}{2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 41

```
DSolve[y'''[x]-y''[x]-y'[x]+y[x]==-Exp[x]*(7+6*x),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(-\frac{x^3}{2} - x^2 + x + c_3x - \frac{1}{2} + c_2 \right) + c_1 e^{-x}$$

19.9 problem section 9.3, problem 9

Internal problem ID [1506]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 9.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$2y''' - 7y'' + 4y' + 4y = e^{2x}(17 + 30x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(2*diff(y(x),x$3)-7*diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=exp(2*x)*(17+30*x),y(x),sing
```

$$y(x) = e^{-\frac{x}{2}}c_2 + e^{2x}\left(x^3 + \frac{1}{2}x^2 + c_1 + c_3x\right)$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 46

```
DSolve[2*y'''[x]-7*y''[x]+4*y'[x]+4*y[x]==Exp[2*x]*(17+30*x),y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow e^{2x}\left(x^3 + \frac{x^2}{2} + \left(-\frac{2}{5} + c_3\right)x + \frac{4}{25} + c_2\right) + c_1e^{-x/2}$$

19.10 problem section 9.3, problem 10

Internal problem ID [1507]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 10.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 5y'' + 3y' + 9y = 2e^{3x}(11 - 24x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$3)-5*diff(y(x),x$2)+3*diff(y(x),x)+9*y(x)=2*exp(3*x)*(11-24*x),y(x),sing
```

$$y(x) = \frac{(-8x^3 + 4c_3x + 17x^2 + 4c_2)e^{3x}}{4} + e^{-x}c_1$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 46

```
DSolve[y'''[x]-5*y''[x]+3*y'[x]+9*y[x]==2*Exp[3*x]*(11-24*x),y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow e^{3x} \left(-2x^3 + \frac{17x^2}{4} + \left(-\frac{17}{8} + c_3 \right) x + \frac{17}{32} + c_2 \right) + c_1 e^{-x}$$

19.11 problem section 9.3, problem 11

Internal problem ID [1508]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 11.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[_3rd_order, _linear, _nonhomogeneous]`

$$y''' - 7y'' + 8y' + 16y = 2e^{4x}(13 + 15x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$3)-7*diff(y(x),x$2)+8*diff(y(x),x)+16*y(x)=2*exp(4*x)*(13+15*x),y(x), sin
```

$$y(x) = (x^3 + c_3x + 2x^2 + c_2) e^{4x} + e^{-x}c_1$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 42

```
DSolve[y'''[x]-7*y''[x]+8*y'[x]+16*y[x]==2*Exp[4*x]*(13+15*x),y[x],x,IncludeSingularSolution
```

$$y(x) \rightarrow e^{4x} \left(x^3 + 2x^2 + \left(-\frac{4}{5} + c_3 \right) x + \frac{4}{25} + c_2 \right) + c_1 e^{-x}$$

19.12 problem section 9.3, problem 12

Internal problem ID [1509]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 12.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$8y''' - 12y'' + 6y' - y = e^{\frac{x}{2}}(4x + 1)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve(8*diff(y(x),x$3)-12*diff(y(x),x$2)+6*diff(y(x),x)-y(x)=exp(x/2)*(1+4*x),y(x), singsol
```

$$y(x) = \frac{(x^4 + x^3 + (48c_2 + \frac{3}{16})x^2 + 48c_3x + 48c_1)e^{\frac{x}{2}}}{48}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 39

```
DSolve[8*y'''[x]-12*y''[x]+6*y'[x]-y[x]==Exp[x/2]*(1+4*x),y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{48}e^{x/2}(x^4 + x^3 + 48c_3x^2 + 48c_2x + 48c_1)$$

19.13 problem section 9.3, problem 13

Internal problem ID [1510]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 13.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 3y''' - 3y'' - 7y' + 6y = -3e^{-x}(-8x^2 + 8x + 12)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$4)+3*diff(y(x),x$3)-3*diff(y(x),x$2)-7*diff(y(x),x)+6*y(x))=-3*exp(-x)*(12+8*x-8*x^2),y(x),x,IncludeS
```

$$y(x) = 3 \left((x-1)^2 e^{2x} + \frac{(c_4 x + c_1) e^{4x}}{3} + \frac{c_3 e^x}{3} + \frac{c_2}{3} \right) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 45

```
DSolve[y''''[x]+3*y'''[x]-3*y''[x]-7*y'[x]+6*y[x]==-3*Exp[-x]*(12+8*x-8*x^2),y[x],x,IncludeS
```

$$y(x) \rightarrow e^{-3x} (3e^{2x}(x-1)^2 + c_2 e^x + e^{4x}(c_4 x + c_3) + c_1)$$

19.14 problem section 9.3, problem 14

Internal problem ID [1511]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 14.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 3y''' + y'' - 3y' - 2y = -3e^{2x}(11 + 12x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$4)+3*diff(y(x),x$3)+diff(y(x),x$2)-3*diff(y(x),x)-2*y(x))=-3*exp(2*x)*(11+
```

$$y(x) = (-e^{4x}x + e^{4x} + c_1e^{3x} + c_4xe^x + c_3e^x + c_2)e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 43

```
DSolve[y''''[x]+3*y'''[x]+y''[x]-3*y'[x]-2*y[x]==-3*Exp[2*x]*(11+12*x),y[x],x,IncludeSingular
```

$$y(x) \rightarrow e^{-2x}(-e^{4x}(x - 1) + e^x(c_3x + c_2) + c_4e^{3x} + c_1)$$

19.15 problem section 9.3, problem 15

Internal problem ID [1512]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 15.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' + 8y''' + 24y'' + 32y' = -16e^{-2x}(-x^3 + x^2 + x + 1)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve(diff(y(x),x$4)+8*diff(y(x),x$3)+24*diff(y(x),x$2)+32*diff(y(x),x)=-16*exp(-2*x)*(1+x+
```

$$y(x) = \frac{((-c_2 - c_3) \cos(2x) + (c_2 - c_3) \sin(2x) - 4x^3 + 4x^2 + 4x + 4) e^{-2x}}{4} - \frac{e^{-4x} c_1}{4} + c_4$$

✓ Solution by Mathematica

Time used: 0.712 (sec). Leaf size: 64

```
DSolve[y''''[x]+8*y'''[x]+24*y''[x]+32*y'[x]==-16*Exp[-2*x]*(1+x+x^2-x^3),y[x],x,IncludeSing
```

$$y(x) \rightarrow \frac{1}{4}e^{-2x}(-4x^3 + 4x^2 + 4x - c_3e^{-2x} - (c_1 + c_2) \cos(2x) + (c_2 - c_1) \sin(2x) + 4) + c_4$$

19.16 problem section 9.3, problem 16

Internal problem ID [1513]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 16.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$4y'''' - 11y'' - 9y' - 2y = -e^x(1 - 6x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 40

```
dsolve(4*diff(y(x),x$4)-11*diff(y(x),x$2)-9*diff(y(x),x)-2*y(x)=-exp(x)*(1-6*x),y(x), singular
```

$$y(x) = e^{-x} \left((c_4 x + c_3) e^{\frac{x}{2}} - \frac{e^{2x} x}{3} + c_2 e^{3x} + c_1 + \frac{e^{2x}}{3} \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 47

```
DSolve[4*y''''[x]-11*y''[x]-9*y'[x]-2*y[x]==-Exp[x]*(1-6*x),y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow -\frac{1}{3}e^x(x-1) + e^{-x/2}(c_2 x + c_1) + c_3 e^{-x} + c_4 e^{2x}$$

19.17 problem section 9.3, problem 17

Internal problem ID [1514]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 17.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 2y''' + 3y' - y = e^x(x^2 + 4x + 3)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 101

```
dsolve(diff(y(x),x$4)-2*diff(y(x),x$3)+0*diff(y(x),x$2)+3*diff(y(x),x)-y(x)=exp(x)*(3+4*x+x^2),y(x),x)
```

$$y(x) = c_1 e^{\text{RootOf}(-Z^4 - 2Z^3 + 3Z - 1, \text{index}=1)x} + c_2 e^{\text{RootOf}(-Z^4 - 2Z^3 + 3Z - 1, \text{index}=2)x} \\ + c_3 e^{\text{RootOf}(-Z^4 - 2Z^3 + 3Z - 1, \text{index}=3)x} \\ + c_4 e^{\text{RootOf}(-Z^4 - 2Z^3 + 3Z - 1, \text{index}=4)x} + (x + 1)^2 e^x$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 123

```
DSolve[y''''[x]-2*y'''[x]+0*y''[x]+3*y'[x]-y[x]==Exp[x]*(3+4*x+x^2),y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 \exp(x \text{Root}[\#1^4 - 2\#1^3 + 3\#1 - 1 \&, 2]) \\ + c_3 \exp(x \text{Root}[\#1^4 - 2\#1^3 + 3\#1 - 1 \&, 3]) \\ + c_4 \exp(x \text{Root}[\#1^4 - 2\#1^3 + 3\#1 - 1 \&, 4]) \\ + c_1 \exp(x \text{Root}[\#1^4 - 2\#1^3 + 3\#1 - 1 \&, 1]) + e^x(x + 1)^2$$

19.18 problem section 9.3, problem 18

Internal problem ID [1515]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 18.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 4y''' + 6y'' - 4y' + 2y = e^{2x}(x^4 + x + 24)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 87

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+6*diff(y(x),x$2)-4*diff(y(x),x)+2*y(x))=exp(2*x)*(24+x
```

$$y(x) = e^{\frac{(2+\sqrt{2})x}{2}} \left(\cos\left(\frac{\sqrt{2}x}{2}\right) c_1 + \sin\left(\frac{\sqrt{2}x}{2}\right) c_2 \right) \\ + \left(\cos\left(\frac{\sqrt{2}x}{2}\right) c_3 + \sin\left(\frac{\sqrt{2}x}{2}\right) c_4 \right) e^{-\frac{(\sqrt{2}-2)x}{2}} \\ + \frac{e^{2x}(x^4 - 8x^3 + 12x^2 + 49x - 62)}{2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 102

```
DSolve[y''''[x]-4*y'''[x]+6*y''[x]-4*y'[x]+2*y[x]==Exp[2*x]*(24+x+x^4),y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{1}{2} e^{x-\frac{x}{\sqrt{2}}} \left(e^{\frac{x}{\sqrt{2}}+x} (x^4 - 8x^3 + 12x^2 + 49x - 62) + 2 \left(c_4 e^{\sqrt{2}x} + c_2 \right) \cos\left(\frac{x}{\sqrt{2}}\right) \right) \\ + 2 \left(c_1 e^{\sqrt{2}x} + c_3 \right) \sin\left(\frac{x}{\sqrt{2}}\right)$$

19.19 problem section 9.3, problem 19

Internal problem ID [1516]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 19.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$2y'''' + 5y''' - 5y' - 2y = 18e^x(2x + 5)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve(2*diff(y(x),x$4)+5*diff(y(x),x$3)+0*diff(y(x),x$2)-5*diff(y(x),x)-2*y(x)=18*exp(x)*(5+2*x),y(x),x)
```

$$y(x) = e^{-2x} \left(e^{\frac{3x}{2}} c_4 + (x^2 + c_1 + 2x) e^{3x} + c_3 e^x + c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 48

```
DSolve[2*y''''[x]+5*y'''[x]+0*y''[x]-5*y'[x]-2*y[x]==18*Exp[x]*(5+2*x),y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-2x} \left(e^{3x} \left(x^2 + 2x - \frac{40}{9} + c_4 \right) + c_1 e^{3x/2} + c_3 e^x + c_2 \right)$$

19.20 problem section 9.3, problem 20

Internal problem ID [1517]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 20.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + y''' - 2y'' - 6y' - 4y = -e^{2x}(15x^2 + 28x + 4)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve(1*diff(y(x),x$4)+1*diff(y(x),x$3)-2*diff(y(x),x$2)-6*diff(y(x),x)-4*y(x))=-exp(2*x)*(4
```

$$y(x) = (\cos(x)c_3 + c_4 \sin(x) + c_1) e^{-x} - \frac{e^{2x}(x^3 - 6c_2 - x)}{6}$$

✓ Solution by Mathematica

Time used: 0.193 (sec). Leaf size: 65

```
DSolve[1*y''''[x]+1*y'''[x]-2*y''[x]-6*y'[x]-4*y[x]==-Exp[2*x]*(4+28*x+15*x^2),y[x],x,Includ
```

$$y(x) \rightarrow \frac{1}{90} e^{-x} (-15e^{3x}x^3 + 15e^{3x}x - 11e^{3x} + 90c_4e^{3x} + 90c_2 \cos(x) + 90c_1 \sin(x) + 90c_3)$$

19.21 problem section 9.3, problem 21

Internal problem ID [1518]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 21.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$2y'''' + y''' - 2y' - y = 3e^{-\frac{x}{2}}(1 - 6x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(2*diff(y(x),x$4)+1*diff(y(x),x$3)-0*diff(y(x),x$2)-2*diff(y(x),x)-1*y(x)=3*exp(-x/2)*
```

$$y(x) = c_3 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_4 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + (4x^2 + c_2 + 4x) e^{-\frac{x}{2}} + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.635 (sec). Leaf size: 63

```
DSolve[2*y''''[x]+1*y'''[x]-0*y''[x]-2*y'[x]-1*y[x]==3*Exp[-x/2]*(1-6*x),y[x],x,IncludeSingu
```

$$y(x) \rightarrow e^{-x/2} \left(4x^2 + 4x + c_4 e^{3x/2} + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) - 8 + c_3 \right)$$

19.22 problem section 9.3, problem 22

Internal problem ID [1519]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 22.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 5y'' + 4y = e^x(-3x^2 + x + 3)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve(1*diff(y(x),x$4)+0*diff(y(x),x$3)-5*diff(y(x),x$2)-0*diff(y(x),x)+4*y(x)=exp(x)*(3+x-
```

$$y(x) = \frac{((x^3 + 6c_1 + x) e^{3x} + 6c_3 e^x + 6c_4 e^{4x} + 6c_2) e^{-2x}}{6}$$

✓ Solution by Mathematica

Time used: 0.111 (sec). Leaf size: 51

```
DSolve[1*y''''[x]+0*y'''[x]-5*y''[x]-0*y'[x]+4*y[x]==Exp[x]*(3+x-3*x^2),y[x],x,IncludeSingul
```

$$y(x) \rightarrow \frac{1}{36} e^x (6x^3 + 6x + 7 + 36c_3) + c_1 e^{-2x} + c_2 e^{-x} + c_4 e^{2x}$$

19.23 problem section 9.3, problem 23

Internal problem ID [1520]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 23.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 2y''' - 3y'' + 4y' + 4y = e^{2x}(18x^2 + 33x + 13)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 38

```
dsolve(1*diff(y(x),x$4)-2*diff(y(x),x$3)-3*diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=exp(2*x)*(13
```

$$y(x) = \frac{(x^4 + x^3 + 6c_4x + x^2 + 6c_2)e^{2x}}{6} + e^{-x}(c_3x + c_1)$$

✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 58

```
DSolve[1*y''''[x]-2*y'''[x]-3*y''[x]+4*y'[x]+4*y[x]==Exp[2*x]*(13+33*x+18*x^2),y[x],x,Includ
```

$$y(x) \rightarrow \frac{1}{54}e^{2x}(9x^4 + 9x^3 + 9x^2 + 18(-1 + 3c_4)x + 10 + 54c_3) + e^{-x}(c_2x + c_1)$$

19.24 problem section 9.3, problem 24

Internal problem ID [1521]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 24.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 3y''' + 4y' = e^{2x}(12x^2 + 26x + 15)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
dsolve(1*diff(y(x),x$4)-3*diff(y(x),x$3)-0*diff(y(x),x$2)+4*diff(y(x),x)+0*y(x)=exp(2*x)*(15
```

$$y(x) = \frac{(2x^4 + 2x^3 + 6x^2 + (6c_3 - 6)x + 6c_2 - 3c_3 + 3)e^{2x}}{12} - e^{-x}c_1 + c_4$$

✓ Solution by Mathematica

Time used: 0.189 (sec). Leaf size: 58

```
DSolve[1*y''''[x]-3*y'''[x]-0*y''[x]+4*y'[x]+0*y[x]==Exp[2*x]*(15+26*x+12*x^2),y[x],x,Includ
```

$$y(x) \rightarrow \frac{1}{12}e^{2x}(2x^4 + 2x^3 + 6x^2 + 6(-2 + c_3)x + 8 + 6c_2 - 3c_3) + c_1(-e^{-x}) + c_4$$

19.25 problem section 9.3, problem 25

Internal problem ID [1522]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 25.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 2y''' + 2y' - y = (x + 1)e^x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 37

```
dsolve(1*diff(y(x),x$4)-2*diff(y(x),x$3)-0*diff(y(x),x$2)+2*diff(y(x),x)-1*y(x)=exp(x)*(1+x)
```

$$y(x) = e^{-x}c_2 + \frac{e^x(x^4 + 48c_4x^2 + 2x^3 + 48c_3x + 48c_1)}{48}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 55

```
DSolve[1*y''''[x]-2*y'''[x]-0*y''[x]+2*y'[x]-1*y[x]==Exp[x]*(1+x),y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \frac{1}{96}e^x(2x^4 + 4x^3 + (-6 + 96c_4)x^2 + (6 + 96c_3)x - 3 + 96c_2) + c_1e^{-x}$$

19.26 problem section 9.3, problem 26

Internal problem ID [1523]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 26.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$2y'''' - 5y''' + 3y'' + y' - y = e^x(11 + 12x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(2*diff(y(x),x$4)-5*diff(y(x),x$3)+3*diff(y(x),x$2)+1*diff(y(x),x)-1*y(x)=exp(x)*(11+12*x),y(x))
```

$$y(x) = c_4 e^{-\frac{x}{2}} + \frac{e^x(x^4 + 6c_3x^2 + x^3 + 6c_2x + 6c_1)}{6}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 58

```
DSolve[2*y''''[x]-5*y'''[x]+3*y''[x]+1*y'[x]-1*y[x]==Exp[x]*(11+12*x),y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^x \left(\frac{x^4}{6} + \frac{x^3}{6} + \left(-\frac{1}{3} + c_4 \right) x^2 + \left(\frac{4}{9} + c_3 \right) x - \frac{8}{27} + c_2 \right) + c_1 e^{-x/2}$$

19.27 problem section 9.3, problem 27

Internal problem ID [1524]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 27.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' + 3y''' + 3y'' + y' = e^{-x}(10x^2 - 24x + 5)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 55

```
dsolve(1*diff(y(x),x$4)+3*diff(y(x),x$3)+3*diff(y(x),x$2)+1*diff(y(x),x)-0*y(x)=exp(-x)*(5-2
```

$$y(x) = \frac{(-x^5 + x^4 - x^3 + (-6c_3 - 3)x^2 + (-6c_2 - 12c_3 - 6)x - 6c_1 - 6c_2 - 12c_3 - 6)e^{-x}}{6} + c_4$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 65

```
DSolve[1*y''''[x]+3*y'''[x]+3*y''[x]+1*y'[x]-0*y[x]==Exp[-x]*(5-24*x+10*x^2),y[x],x,IncludeS
```

$$y(x) \rightarrow \frac{1}{6}e^{-x}(-x^5 + x^4 - x^3 - 3(1 + 2c_3)x^2 - 6(1 + c_2 + 2c_3)x - 6(1 + c_1 + c_2 + 2c_3)) + c_4$$

19.28 problem section 9.3, problem 28

Internal problem ID [1525]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 28.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 7y''' + 18y'' - 20y' + 8y = e^{2x}(-5x^2 - 8x + 3)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 42

```
dsolve(1*diff(y(x),x$4)-7*diff(y(x),x$3)+18*diff(y(x),x$2)-20*diff(y(x),x)+8*y(x)=exp(2*x)*
```

$$y(x) = -\frac{e^x((x^5 - x^4 - 12c_4x^2 - 2x^3 - 12c_3x - 12c_2)e^x - 12c_1)}{12}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 59

```
DSolve[1*y''''[x]-7*y'''[x]+18*y''[x]-20*y'[x]+8*y[x]==Exp[2*x]*(3-8*x-5*x^2),y[x],x,Include
```

$$y(x) \rightarrow \frac{1}{12}e^{2x}(-x^5 + x^4 + 2x^3 + 6(-1 + 2c_4)x^2 + 12(1 + c_3)x + 12(-1 + c_2)) + c_1e^x$$

19.29 problem section 9.3, problem 29

Internal problem ID [1526]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 29.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[_3rd_order, _linear, _nonhomogeneous]`

$$y''' - y'' - 4y' + 4y = e^{-x}((16 + 10x) \cos(x) + (30 - 10x) \sin(x))$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 40

```
dsolve(0*diff(y(x),x$4)+1*diff(y(x),x$3)-1*diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=exp(-x)*((16+10*x)*cos(x)+(30-10*x)*sin(x))
```

$$y(x) = (c_1 e^{3x} + c_3 e^{4x} + ((x + 1) \cos(x) - \sin(x) (-2 + x)) e^x + c_2) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 49

```
DSolve[0*y''''[x]+1*y'''[x]-1*y''[x]-4*y'[x]+4*y[x]==Exp[-x]*((16+10*x)*Cos[x]+(30-10*x)*Sin[x]]
```

$$y(x) \rightarrow e^{-2x}(-e^x(x - 2) \sin(x) + e^x(x + 1) \cos(x) + c_2 e^{3x} + c_3 e^{4x} + c_1)$$

19.30 problem section 9.3, problem 30

Internal problem ID [1527]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 30.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + y'' - 4y' - 4y = e^{-x}((1 - 22x) \cos(2x) - (1 + 6x) \sin(2x))$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
dsolve(1*diff(y(x),x$3)+1*diff(y(x),x$2)-4*diff(y(x),x)-4*y(x)=exp(-x)*((1-22*x)*cos(2*x)-(1
```

$$y(x) = -\left(e^x(x-1) \cos(2x) - (x+1)e^x \sin(2x) - c_3 e^{4x} + \left(-c_2 + \frac{5}{3}\right) e^x - c_1 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 51

```
DSolve[1*y'''[x]+1*y''[x]-4*y'[x]-4*y[x]==Exp[-x]*((1-22*x)*Cos[2*x]-(1+6*x)*Sin[2*x]),y[x],
```

$$y(x) \rightarrow e^{-2x} (e^x(x+1) \sin(2x) - e^x(x-1) \cos(2x) + c_2 e^x + c_3 e^{4x} + c_1)$$

19.31 problem section 9.3, problem 31

Internal problem ID [1528]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 31.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + 2y' - 2y = e^{2x}((-x^2 + 5x + 27) \cos(x) + (9x^2 + 13x + 2) \sin(x))$$

✓ Solution by Maple

Time used: 0.266 (sec). Leaf size: 179

```
dsolve(1*diff(y(x),x$3)-1*diff(y(x),x$2)+2*diff(y(x),x)-2*y(x)=exp(2*x)*((27+5*x-x^2)*cos(1*
```

$$y(x) = \frac{\sqrt{2} \left(\int e^{2x} (\sqrt{2} \cos(\sqrt{2}x) - \sin(\sqrt{2}x)) ((-9x^2 - 13x - 2) \sin(x) + (x^2 - 5x - 27) \cos(x)) dx \right) \cos(\sqrt{2}x)}{6} - \frac{\sqrt{2} \left(\int -e^{2x} (\sqrt{2} \sin(\sqrt{2}x) + \cos(\sqrt{2}x)) ((-9x^2 - 13x - 2) \sin(x) + (x^2 - 5x - 27) \cos(x)) dx \right) \sin(\sqrt{2}x)}{6} + c_2 \cos(\sqrt{2}x) + c_3 \sin(\sqrt{2}x) + \frac{(5(-x^2 + x + 3) \cos(x) + 4(x^2 + \frac{5}{2}x + \frac{7}{4}) \sin(x)) e^{2x}}{3} + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 60

```
DSolve[1*y'''[x]-1*y''[x]+2*y'[x]-2*y[x]==Exp[2*x]*((27+5*x-x^2)*Cos[1*x]+(2+13*x+9*x^2)*Sin
```

$$y(x) \rightarrow e^{2x}((-x^2 + x + 1) \cos(x) + (2x + 1) \sin(x)) + c_3 e^x + c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)$$

19.32 problem section 9.3, problem 32

Internal problem ID [1529]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 32.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 2y'' + y' - 2y = -e^x((4x^2 + 5x + 9) \cos(2x) - (-3x^2 - 5x + 6) \sin(2x))$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 48

```
dsolve(1*diff(y(x),x$3)-2*diff(y(x),x$2)+1*diff(y(x),x)-2*y(x)=-exp(x)*((9+5*x+4*x^2)*cos(2*x)-(-3*x^2-5*x+6)*sin(2*x)),y(x))
```

$$y(x) = \frac{e^x \left(x^2 + \frac{3}{5}x - \frac{27}{25} \right) \sin(2x)}{2} + \frac{(55x + 61) e^x \cos(2x)}{50} + \cos(x) c_1 + \sin(x) c_2 + c_3 e^{2x}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 65

```
DSolve[1*y'''[x]-2*y''[x]+1*y'[x]-2*y[x]==Exp[2*x]*((9+5*x+4*x^2)*Cos[2*x]-(-3*x^2-5*x+6)*Sin[2*x]),y[x]]
```

$$y(x) \rightarrow -\frac{e^{2x}((6760x^2 - 17680x - 29907) \sin(2x) + 2(5915x^2 + 7345x + 3928) \cos(2x))}{43940} + c_3 e^{2x} + c_1 \cos(x) + c_2 \sin(x)$$

19.33 problem section 9.3, problem 33

Internal problem ID [1530]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 33.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[_3rd_order, _linear, _nonhomogeneous]`

$$y''' + 3y'' + 4y' + 12y = 8 \cos(2x) - 16 \sin(2x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(1*diff(y(x),x$3)+3*diff(y(x),x$2)+4*diff(y(x),x)+12*y(x)=8*cos(2*x)-16*sin(2*x),y(x),
```

$$y(x) = \frac{(169c_1 + 104x + 56) \cos(2x)}{169} + \frac{(182x + 169c_3 - 136) \sin(2x)}{169} + e^{-3x}c_2$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 47

```
DSolve[1*y'''[x]+3*y''[x]+4*y'[x]+12*y[x]==8*Cos[2*x]-16*Sin[2*x],y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow \frac{1}{169} (169c_3 e^{-3x} + (104x + 43 + 169c_1) \cos(2x) + (182x - 32 + 169c_2) \sin(2x))$$

19.34 problem section 9.3, problem 34

Internal problem ID [1531]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 34.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + 2y = e^x((20 + 4x) \cos(x) - (12x + 12) \sin(x))$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 37

```
dsolve(1*diff(y(x),x$3)-1*diff(y(x),x$2)+0*diff(y(x),x)+2*y(x)=exp(x)*((20+4*x)*cos(x)-(12+12*x)*sin(x)),y(x),x)
```

$$y(x) = e^{-x}c_1 + e^x \left(\left(x^2 + x + c_2 + \frac{22}{5} \right) \cos(x) + \sin(x) \left(x^2 + 3x + \frac{1}{5} + c_3 \right) \right)$$

✓ Solution by Mathematica

Time used: 0.308 (sec). Leaf size: 60

```
DSolve[1*y'''[x]-1*y''[x]+0*y'[x]+2*y[x]==Exp[x]*((20+4*x)*Cos[x]-(12+12*x)*Sin[x]),y[x],x]
```

$$y(x) \rightarrow \frac{1}{10}e^x(10x^2 + 10x + 23 + 10c_2) \cos(x) + \frac{1}{10}e^x(10x^2 + 30x - 21 + 10c_1) \sin(x) + c_3e^{-x}$$

19.35 problem section 9.3, problem 35

Internal problem ID [1532]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 35.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 7y'' + 20y' - 24y = -e^{2x}((13 - 8x) \cos(2x) - (8 - 4x) \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 48

```
dsolve(1*diff(y(x),x$3)-7*diff(y(x),x$2)+20*diff(y(x),x)-24*y(x)=-exp(2*x)*((13-8*x)*cos(2*x)
```

$$y(x) = \frac{((-20x^2 + 40c_2 + 60x - 83) \cos(2x) + 20 \sin(2x) (x + 2c_3 - \frac{47}{10})) e^{2x}}{40} + c_1 e^{3x}$$

✓ Solution by Mathematica

Time used: 0.955 (sec). Leaf size: 55

```
DSolve[1*y'''[x]-7*y''[x]+20*y'[x]-24*y[x]==-Exp[2*x]*((13-8*x)*Cos[2*x]-(8-4*x)*Sin[2*x]),y
```

$$y(x) \rightarrow \frac{1}{40} e^{2x} ((-20x^2 + 60x + 21 + 40c_2) \cos(2x) + 40c_3 e^x + (20x - 37 + 40c_1) \sin(2x))$$

19.36 problem section 9.3, problem 36

Internal problem ID [1533]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 36.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$y''' - 6y'' + 18y' = -e^{3x}((-3x + 2) \cos(3x) - (3 + 3x) \sin(3x))$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
dsolve(1*diff(y(x),x$3)-6*diff(y(x),x$2)+18*diff(y(x),x)-0*y(x)=-exp(3*x)*((2-3*x)*cos(3*x)-
```

$$y(x) = \frac{((-3x^2 + 6c_1 - 6c_2 - 1) \cos(3x) - 3 \sin(3x) (x - 2c_1 - 2c_2 + \frac{1}{9})) e^{3x}}{36} + c_3$$

✓ Solution by Mathematica

Time used: 3.765 (sec). Leaf size: 57

```
DSolve[1*y'''[x]-6*y''[x]+18*y'[x]-0*y[x]==-Exp[3*x]*((2-3*x)*Cos[3*x]-(3+3*x)*Sin[3*x]),y[x]
```

$$y(x) \rightarrow c_3 - \frac{1}{216} e^{3x} (6(3x^2 + 1 + 6c_1 - 6c_2) \cos(3x) + (18x + 1 - 36c_1 - 36c_2) \sin(3x))$$

19.37 problem section 9.3, problem 37

Internal problem ID [1534]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 37.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 2y''' - 2y'' - 8y' - 8y = e^x(8 \cos(x) + 16 \sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 47

```
dsolve(1*diff(y(x),x$4)+2*diff(y(x),x$3)-2*diff(y(x),x$2)-8*diff(y(x),x)-8*y(x)=exp(x)*(8*cos(x)+16*sin(x)),y(x),x,Incl
```

$$y(x) = -\frac{((7 \sin(x) + \cos(x)) e^{3x} - 10c_3 \cos(x) e^x - 10c_4 \sin(x) e^x - 10c_2 e^{4x} - 10c_1) e^{-2x}}{10}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 56

```
DSolve[1*y''''[x]+2*y'''[x]-2*y''[x]-8*y'[x]-8*y[x]==Exp[x]*(8*Cos[x]+16*Sin[x]),y[x],x,Incl
```

$$y(x) \rightarrow c_3 e^{-2x} + c_4 e^{2x} - \frac{1}{10} e^x (7 \sin(x) + \cos(x)) + c_2 e^{-x} \cos(x) + c_1 e^{-x} \sin(x)$$

19.38 problem section 9.3, problem 38

Internal problem ID [1535]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 38.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 3y''' + 2y'' + 2y' - 4y = e^x(-\sin(2x) + 2\cos(2x))$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(1*diff(y(x),x$4)-3*diff(y(x),x$3)+2*diff(y(x),x$2)+2*diff(y(x),x)-4*y(x)=exp(x)*(2*cos(2*x)-sin(2*x)),y(x),x,inc
```

$$y(x) = e^{-x}c_1 + c_2e^{2x} + e^x\left(\frac{\cos(x)^2}{6} + \left(c_3 - \frac{\sin(x)}{6}\right)\cos(x) + c_4\sin(x) - \frac{1}{12}\right)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 56

```
DSolve[1*y''''[x]-3*y'''[x]+2*y''[x]+2*y'[x]-4*y[x]==Exp[x]*(2*Cos[2*x]-Sin[2*x]),y[x],x,Inc
```

$$y(x) \rightarrow c_3e^{-x} + c_4e^{2x} + \frac{1}{12}e^x(\cos(2x) - \sin(2x)) + c_2e^x \cos(x) + c_1e^x \sin(x)$$

19.39 problem section 9.3, problem 39

Internal problem ID [1536]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 39.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 8y''' + 24y'' - 32y' + 15y = e^{2x}(15x \cos(2x) + 32 \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

```
dsolve(1*diff(y(x),x$4)-8*diff(y(x),x$3)+24*diff(y(x),x$2)-32*diff(y(x),x)+15*y(x)=exp(2*x)*
```

$$y(x) = e^x(2e^x x \cos(x)^2 + c_4 \sin(x) e^x + c_3 \cos(x) e^x - x e^x + c_2 e^{2x} + c_1)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 45

```
DSolve[1*y''''[x]-8*y'''[x]+24*y''[x]-32*y'[x]+15*y[x]==Exp[2*x]*(15*x*Cos[2*x]+32*Sin[2*x])
```

$$y(x) \rightarrow e^x(e^x x \cos(2x) + c_4 e^{2x} + c_2 e^x \cos(x) + c_1 e^x \sin(x) + c_3)$$

19.40 problem section 9.3, problem 40

Internal problem ID [1537]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 40.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 6y''' + 13y'' + 12y' + 4y = e^{-x}((-x + 4) \cos(x) - (x + 5) \sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 43

```
dsolve(1*diff(y(x),x$4)+6*diff(y(x),x$3)+13*diff(y(x),x$2)+12*diff(y(x),x)+4*y(x)=exp(-1*x)*
```

$$y(x) = \frac{((-x - 1) \cos(x) + \sin(x)(-2 + x) + 2c_3x + 2c_2) e^{-x}}{2} + e^{-2x}(c_4x + c_1)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 56

```
DSolve[1*y''''[x]+6*y'''[x]+13*y''[x]+12*y'[x]+4*y[x]==Exp[-1*x]*((4-x)*Cos[x]-(5+x)*Sin[x])
```

$$y(x) \rightarrow \frac{1}{2} e^{-2x} (e^x (x - 2) \sin(x) - e^x (x + 1) \cos(x) + 2(c_2x + c_3e^x + c_4e^x x + c_1))$$

19.41 problem section 9.3, problem 41

Internal problem ID [1538]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 41.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 3y''' + 2y'' - 2y' - 4y = -e^{-x}(\cos(x) - \sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(1*diff(y(x),x$4)+3*diff(y(x),x$3)+2*diff(y(x),x$2)-2*diff(y(x),x)-4*y(x))=-exp(-1*x)*(
```

$$y(x) = \frac{(10c_1 e^{3x} + ((x + 10c_3 + \frac{14}{5}) \cos(x) + 2 \sin(x) (x + 5c_4 - \frac{1}{5})) e^x + 10c_2) e^{-2x}}{10}$$

✓ Solution by Mathematica

Time used: 0.136 (sec). Leaf size: 58

```
DSolve[1*y''''[x]+3*y'''[x]+2*y''[x]-2*y'[x]-4*y[x]==-Exp[-1*x]*(Cos[x]-Sin[x]),y[x],x,Inclu
```

$$y(x) \rightarrow \frac{1}{50} e^{-2x} (50(c_4 e^{3x} + c_3) + e^x (5x + 14 + 50c_2) \cos(x) + e^x (10x - 7 + 50c_1) \sin(x))$$

19.42 problem section 9.3, problem 42

Internal problem ID [1539]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 42.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 5y''' + 13y'' - 19y' + 10y = e^x(\cos(2x) + \sin(2x))$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 48

```
dsolve(1*diff(y(x),x$4)-5*diff(y(x),x$3)+13*diff(y(x),x$2)-19*diff(y(x),x)+10*y(x)=exp(x)*(cos(2*x)+sin(2*x)),y(x),x,In
```

$$y(x) = \frac{3e^x\left(x + \frac{40c_3}{3} + \frac{1}{15}\right)\cos(2x)}{40} - \frac{e^x\left(x - 40c_4 + \frac{39}{10}\right)\sin(2x)}{40} + c_2e^{2x} + \frac{e^x(8c_1 + 1)}{8}$$

✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 53

```
DSolve[1*y''''[x]-5*y'''[x]+13*y''[x]-19*y'[x]+10*y[x]==Exp[x]*(Cos[2*x]+Sin[2*x]),y[x],x,In
```

$$y(x) \rightarrow \frac{1}{400}e^x(400(c_4e^x + c_3) + (30x - 13 + 400c_2)\cos(2x) - 2(5x + 17 - 200c_1)\sin(2x))$$

19.43 problem section 9.3, problem 43

Internal problem ID [1540]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 43.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 8y''' + 32y'' + 64y' + 39y = e^{-2x}((4 - 15x) \cos(3x) - (4 + 15x) \sin(3x))$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 57

```
dsolve(1*diff(y(x),x$4)+8*diff(y(x),x$3)+32*diff(y(x),x$2)+64*diff(y(x),x)+39*y(x)=exp(-2*x)
```

$$y(x) = \frac{((-30x^2 + 240c_3 + 30x + 11) \cos(3x) + 30(x^2 + x + 8c_4 - \frac{11}{30}) \sin(3x)) e^{-2x}}{240} + c_1 e^{-3x} + e^{-x} c_2$$

✓ Solution by Mathematica

Time used: 0.841 (sec). Leaf size: 73

```
DSolve[1*y''''[x]+8*y'''[x]+32*y''[x]+64*y'[x]+39*y[x]==Exp[-2*x]*((4-15*x)*Cos[3*x]-(4+15*x)
```

$$y(x) \rightarrow \frac{1}{720} e^{-3x} (-5e^x (18x^2 - 18x - 5 - 144c_2) \cos(3x) + e^x (90x^2 + 90x - 41 + 720c_1) \sin(3x) + 720(c_4 e^{2x} + c_3))$$

19.44 problem section 9.3, problem 44

Internal problem ID [1541]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 44.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 5y''' + 13y'' - 19y' + 10y = e^x((8x + 7) \cos(2x) + (8 - 4x) \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(1*diff(y(x),x$4)-5*diff(y(x),x$3)+13*diff(y(x),x$2)-19*diff(y(x),x)+10*y(x)=exp(x)*((8*x+7)*cos(2*x)+(8-4*x)*sin(2*x)))
```

$$y(x) = -\frac{e^x(x^2 + x - 4c_4 + \frac{23}{4}) \sin(2x)}{4} + \frac{(4c_3 + 3)e^x \cos(2x)}{4} + c_2 e^{2x} + \frac{e^x(2c_1 + 7)}{2}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 172

```
DSolve[1*y''''[x]-5*y'''[x]+13*y''[x]-19*y'[x]-10*y[x]==Exp[x]*((7+8*x)*Cos[2*x]+(8-4*x)*Sin[2*x]),y[x]]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \exp(x \text{Root}[\#1^4 - 5\#1^3 + 13\#1^2 - 19\#1 - 10\&, 1]) \\ & + c_3 \exp(x \text{Root}[\#1^4 - 5\#1^3 + 13\#1^2 - 19\#1 - 10\&, 3]) \\ & + c_4 \exp(x \text{Root}[\#1^4 - 5\#1^3 + 13\#1^2 - 19\#1 - 10\&, 4]) \\ & + c_2 \exp(x \text{Root}[\#1^4 - 5\#1^3 + 13\#1^2 - 19\#1 - 10\&, 2]) \\ & - \frac{1}{100} e^x (-20x \sin(2x) + 64 \sin(2x) + 40x \cos(2x) + 67 \cos(2x)) \end{aligned}$$

19.45 problem section 9.3, problem 45

Internal problem ID [1542]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 45.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 4y''' + 8y'' + 8y' + 4y = -2e^x(\cos(x) - \sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve(1*diff(y(x),x$4)+4*diff(y(x),x$3)+8*diff(y(x),x$2)+8*diff(y(x),x)+4*y(x)=-2*exp(x)*(cos(x)-sin(x)),y(x),x,In
```

$$y(x) = ((c_3x + c_1) \cos(x) + \sin(x)(c_4x + c_2)) e^{-x} - \frac{(\sin(x) + \cos(x)) e^x}{16}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 54

```
DSolve[1*y''''[x]+4*y'''[x]+8*y''[x]+8*y'[x]+4*y[x]==-2*Exp[x]*(Cos[1*x]-Sin[1*x]),y[x],x,In
```

$$y(x) \rightarrow \frac{1}{16} e^{-x} ((-e^{2x} + 16(c_4x + c_3)) \cos(x) - (e^{2x} - 16(c_2x + c_1)) \sin(x))$$

19.46 problem section 9.3, problem 46

Internal problem ID [1543]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 46.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 8y''' + 32y'' - 64y' + 64y = e^{2x}(\cos(2x) - \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 52

```
dsolve(1*diff(y(x),x$4)-8*diff(y(x),x$3)+32*diff(y(x),x$2)-64*diff(y(x),x)+64*y(x)=exp(2*x)*
```

$$y(x) = \frac{e^{2x} \left((x^2 + (-32c_3 - \frac{3}{2})x - 32c_1 - \frac{41}{4}) \cos(2x) - \sin(2x) (x^2 + (32c_4 + \frac{1}{2})x + 32c_2 - \frac{517}{18}) \right)}{32}$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 65

```
DSolve[1*y''''[x]-8*y'''[x]+32*y''[x]-64*y'[x]+64*y[x]==Exp[2*x]*(Cos[2*x]-Sin[2*x]),y[x],x,
```

$$y(x) \rightarrow \frac{1}{256} e^{2x} \left((-8x^2 + 4(1 + 64c_4)x + 5 + 256c_3) \cos(2x) + (8x^2 + 8(1 + 32c_2)x - 1 + 256c_1) \sin(2x) \right)$$

19.47 problem section 9.3, problem 47

Internal problem ID [1544]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 47.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 8y''' + 26y'' - 40y' + 25y = e^{2x}(3 \cos(x) - (3x + 1) \sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(1*diff(y(x),x$4)-8*diff(y(x),x$3)+26*diff(y(x),x$2)-40*diff(y(x),x)+25*y(x)=exp(2*x)*
```

$$y(x) = \frac{e^{2x}((x^3 + x^2 + (8c_4 + 3)x + 8c_2 - 2) \sin(x) + 8 \cos(x) ((c_3 + \frac{1}{4})x + c_1 + \frac{9}{16}))}{8}$$

✓ Solution by Mathematica

Time used: 0.239 (sec). Leaf size: 60

```
DSolve[1*y''''[x]-8*y'''[x]+26*y''[x]-40*y'[x]+25*y[x]==Exp[2*x]*(3*Cos[1*x]- (1+3*x)*Sin[1*x
```

$$y(x) \rightarrow \frac{1}{16} e^{2x}((2x^3 + 2x^2 + (9 + 16c_2)x - 1 + 16c_1) \sin(x) + (2(1 + 8c_4)x + 3 + 16c_3) \cos(x))$$

19.48 problem section 9.3, problem 48

Internal problem ID [1545]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 48.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 4y'' + 5y' - 2y = e^{2x} - 4e^x - 2\cos(x) + 4\sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(1*diff(y(x),x$3)-4*diff(y(x),x$2)+5*diff(y(x),x)-2*y(x)=exp(2*x)-4*exp(x)-2*cos(x)+4*
```

$$y(x) = (x + c_2 - 2) e^{2x} + (2x^2 + (c_3 + 4)x + c_1 + 4) e^x - \cos(x)$$

✓ Solution by Mathematica

Time used: 0.373 (sec). Leaf size: 38

```
DSolve[1*y'''[x]-4*y''[x]+5*y'[x]-2*y[x]==Exp[2*x]-4*Exp[x]-2*Cos[x]+4*Sin[x],y[x],x,Include
```

$$y(x) \rightarrow -\cos(x) + e^x(2x^2 + (4 + c_2)x + e^x(x - 2 + c_3) + 4 + c_1)$$

19.49 problem section 9.3, problem 49

Internal problem ID [1546]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 49.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' + y' - y = 5e^{2x} + 2e^x - 4\cos(x) + 4\sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve(1*diff(y(x),x$3)-1*diff(y(x),x$2)+1*diff(y(x),x)-1*y(x)=5*exp(2*x)+2*exp(x)-4*cos(x)+4*sin(x),y(x),x)
```

$$y(x) = e^{2x} + (2x + c_1 + 2) \cos(x) + (c_2 + x - 1) e^x + (c_3 - 2) \sin(x)$$

✓ Solution by Mathematica

Time used: 0.282 (sec). Leaf size: 35

```
DSolve[1*y'''[x]-1*y''[x]+1*y'[x]-1*y[x]==5*Exp[2*x]+2*Exp[x]-4*Cos[x]+4*Sin[x],y[x],x,IncludeSingularFunctions->True]
```

$$y(x) \rightarrow e^x(x + e^x - 1 + c_3) + (2x + 1 + c_1) \cos(x) + (-2 + c_2) \sin(x)$$

19.50 problem section 9.3, problem 50

Internal problem ID [1547]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 50.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$y''' - y' = -2 - 2x + 4e^x - 6e^{-x} + 96e^{3x}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(1*diff(y(x),x$3)-0*diff(y(x),x$2)-1*diff(y(x),x)-0*y(x)=-2*(1+x)+4*exp(x)-6*exp(-x)+96*exp(3*x),y(x),x,Inc
```

$$y(x) = \frac{(-6x - 2c_2 - 9)e^{-x}}{2} + 4e^{3x} + (2x - 3 + c_1)e^x + x^2 + 2x + c_3$$

✓ Solution by Mathematica

Time used: 1.399 (sec). Leaf size: 49

```
DSolve[1*y'''[x]-0*y''[x]-1*y'[x]-0*y[x]==-2*(1+x)+4*Exp[x]-6*Exp[-x]+96*Exp[3*x],y[x],x,Inc
```

$$y(x) \rightarrow x(x + 2) + 4e^{3x} + e^x(2x - 3 + c_1) - \frac{1}{2}e^{-x}(6x + 9 + 2c_2) + c_3$$

19.51 problem section 9.3, problem 51

Internal problem ID [1548]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 51.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 4y'' + 9y' - 10y = 10e^{2x} + 20e^x \sin(2x) - 10$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
dsolve(1*diff(y(x),x$3)-4*diff(y(x),x$2)+9*diff(y(x),x)-10*y(x)=10*exp(2*x)+20*exp(x)*sin(2*x)-10,y(x),x,IncludeS
```

$$y(x) = 1 + e^x \left(x - \frac{8}{5} + c_2 \right) \cos(2x) + \frac{(-4 + 10x + 5c_1)e^{2x}}{5} - 2 \left(x - \frac{c_3}{2} + \frac{13}{20} \right) e^x \sin(2x)$$

✓ Solution by Mathematica

Time used: 0.992 (sec). Leaf size: 72

```
DSolve[1*y'''[x]-4*y''[x]+9*y'[x]-10*y[x]==10*Exp[2*x]+20*Exp[x]*Sin[2*x]-10,y[x],x,IncludeS
```

$$y(x) \rightarrow 2e^{2x}x - \frac{4e^{2x}}{5} + c_3e^{2x} + \frac{1}{10}e^x(10x - 11 + 10c_2) \cos(2x) - \frac{1}{20}e^x(40x + 21 - 20c_1) \sin(2x) + 1$$

19.52 problem section 9.3, problem 52

Internal problem ID [1549]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 52.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 3y'' + 3y' + y = 12e^{-x} + 9\cos(2x) - 13\sin(2x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(1*diff(y(x),x$3)+3*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=12*exp(-x)+9*cos(2*x)-13*sin(2*x),y(x),x)
```

$$y(x) = (c_3x^2 + 2x^3 + c_2x + c_1)e^{-x} - \cos(2x) + \sin(2x)$$

✓ Solution by Mathematica

Time used: 0.295 (sec). Leaf size: 46

```
DSolve[1*y'''[x]+3*y''[x]+3*y'[x]+1*y[x]==12*Exp[-x]+9*Cos[2*x]-13*Sin[2*x],y[x],x,IncludeS
```

$$y(x) \rightarrow e^{-x}(2x^3 + c_3x^2 + e^x \sin(2x) - e^x \cos(2x) + c_2x + c_1)$$

19.53 problem section 9.3, problem 53

Internal problem ID [1550]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 53.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + y'' - y' - y = 4e^{-x}(1 - 6x) - 2x \cos(x) + 2(x + 1) \sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(1*diff(y(x),x$3)+1*diff(y(x),x$2)-1*diff(y(x),x)-1*y(x)=4*exp(-x)*(1-6*x)-2*x*cos(x)+
```

$$y(x) = (2x^3 + 2x^2 + (c_3 + 2)x + c_2 + 1)e^{-x} + x \cos(x) + e^x c_1 - 2 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.627 (sec). Leaf size: 54

```
DSolve[1*y'''[x]+1*y''[x]-1*y'[x]-1*y[x]==4*Exp[-x]*(1-6*x)-2*x*Cos[x]+2*(1+x)*Sin[x],y[x],x
```

$$y(x) \rightarrow e^{-x}(2x^3 + 2x^2 + 2x - 2e^x \sin(x) + e^x x \cos(x) + c_2 x + c_3 e^{2x} + 1 + c_1)$$

19.54 problem section 9.3, problem 54

Internal problem ID [1551]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 54.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 5y'' + 4y = -12e^x + 6e^{-x} + 10\cos(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

```
dsolve(diff(y(x),x$4)-0*diff(y(x),x$3)-5*diff(y(x),x$2)-0*diff(y(x),x)+4*y(x))=-12*exp(x)+6*exp(-x)+10*cos(x),y(x),x,inc
```

$$y(x) = e^{-2x} \left(\frac{(6x + 3c_1 + 1)e^{3x}}{3} + \cos(x)e^{2x} + c_4e^{4x} + \left(x + c_3 - \frac{1}{6} \right) e^x + c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.19 (sec). Leaf size: 58

```
DSolve[y''''[x]-0*y'''[x]-5*y''[x]-0*y'[x]+4*y[x]==-12*Exp[x]+6*Exp[-x]+10*Cos[x],y[x],x,Inc
```

$$y(x) \rightarrow \cos(x) + c_1e^{-2x} + \frac{1}{6}e^{-x}(6x - 1 + 6c_2) + \frac{1}{3}e^x(6x + 1 + 3c_3) + c_4e^{2x}$$

19.55 problem section 9.3, problem 55

Internal problem ID [1552]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 55.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 4y''' + 11y'' - 14y' + 10y = -e^x(\sin(x) + 2\cos(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+11*diff(y(x),x$2)-14*diff(y(x),x)+10*y(x)=-exp(x)*(sin(x)+2*cos(2*x)),y(x),x,Inc
```

$$y(x) = \frac{e^x \left((6c_3 + \frac{7}{3}) \cos(2x) + (x + 6c_4) \sin(2x) + (x + 6c_1) \cos(x) + 6(\frac{1}{9} + c_2) \sin(x) \right)}{6}$$

✓ Solution by Mathematica

Time used: 0.151 (sec). Leaf size: 53

```
DSolve[y''''[x]-4*y'''[x]+11*y''[x]-14*y'[x]+10*y[x]==-Exp[x]*(Sin[x]+2*Cos[2*x]),y[x],x,Inc
```

$$y(x) \rightarrow \frac{1}{36} e^x \left((11 + 36c_2) \cos(2x) + (1 + 36c_3) \sin(x) + 6 \cos(x) (x + 2(x + 6c_1) \sin(x) + 6c_4) \right)$$

19.56 problem section 9.3, problem 56

Internal problem ID [1553]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 56.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 2y''' - 3y'' - 4y' + 4y = 2(x + 1)e^x + e^{-2x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)-3*diff(y(x),x$2)-4*diff(y(x),x)+4*y(x))=2*exp(x)*(1+x)
```

$$y(x) = \frac{e^{-2x} \left((x^3 + x^2 + (27c_3 - 2)x + 27c_1 + \frac{10}{9}) e^{3x} + \frac{3x^2}{2} + (27c_4 + 2)x + 27c_2 + 1 \right)}{27}$$

✓ Solution by Mathematica

Time used: 0.239 (sec). Leaf size: 66

```
DSolve[y''''[x]+2*y'''[x]-3*y''[x]-4*y'[x]+4*y[x]==2*Exp[x]*(1+x)+Exp[-2*x],y[x],x,IncludeSi
```

$$y(x) \rightarrow \frac{1}{54} e^{-2x} (3x^2 + (4 + 54c_2)x + 2 + 54c_1) + \frac{1}{243} e^x (9x^3 + 9x^2 + 9(-2 + 27c_4)x + 10 + 243c_3)$$

19.57 problem section 9.3, problem 57

Internal problem ID [1554]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 57.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 4y = \sinh(x) \cos(x) - \cosh(x) \sin(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
dsolve(diff(y(x),x$4)+0*diff(y(x),x$3)-0*diff(y(x),x$2)-0*diff(y(x),x)+4*y(x))=sinh(x)*cos(x)
```

$$y(x) = \frac{((-4x + 64c_4 - 3) \sin(x) + \cos(x) (64c_3 - 3)) e^{-x}}{64} + \frac{e^x \left((x + 16c_2 - \frac{3}{4}) \sin(x) + \cos(x) (16c_1 + \frac{3}{4}) \right)}{16}$$

✓ Solution by Mathematica

Time used: 0.676 (sec). Leaf size: 63

```
DSolve[y''''[x]+0*y'''[x]-0*y''[x]-0*y'[x]+4*y[x]==Sinh[x]*Cos[x]-Cosh[x]*Sin[x],y[x],x,Incl
```

$$y(x) \rightarrow \frac{1}{64} e^{-x} \left((3 + 64c_4) e^{2x} - 3 + 64c_1 \right) \cos(x) + (-4x + e^{2x} (4x - 3 + 64c_3) - 3 + 64c_2) \sin(x)$$

19.58 problem section 9.3, problem 58

Internal problem ID [1555]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 58.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 5y''' + 9y'' + 7y' + 2y = e^{-x}(30 + 24x) - e^{-2x}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$4)+5*diff(y(x),x$3)+9*diff(y(x),x$2)+7*diff(y(x),x)+2*y(x))=exp(-x)*(30+24*x)-exp(-2*x),y(x))
```

$$y(x) = (x^4 + x^3 + (c_4 - 3)x^2 + (c_3 + 6)x + c_2 - 6)e^{-x} + e^{-2x}(x + c_1 + 3)$$

✓ Solution by Mathematica

Time used: 0.274 (sec). Leaf size: 44

```
DSolve[y''''[x]+5*y'''[x]+9*y''[x]+7*y'[x]+2*y[x]==Exp[-x]*(30+24*x)-Exp[-2*x],y[x],x,IncludeSingularFunctions->True]
```

$$y(x) \rightarrow e^{-2x}(e^x(x^4 + x^3 + (-3 + c_4)x^2 + (6 + c_3)x - 6 + c_2) + x + 3 + c_1)$$

19.59 problem section 9.3, problem 59

Internal problem ID [1556]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 59.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 4y''' + 7y'' - 6y' + 2y = e^x(12x - 2\cos(x) + 2\sin(x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+7*diff(y(x),x$2)-6*diff(y(x),x)+2*y(x))=exp(x)*(12*x-2*cos(x)+2*sin(x)),y(x),x,In
```

$$y(x) = e^x((x + c_3 + 3)\cos(x) + (c_4 + x - 2)\sin(x) + 2x^3 + (c_2 - 12)x + c_1)$$

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 40

```
DSolve[y''''[x]-4*y'''[x]+7*y''[x]-6*y'[x]+2*y[x]==Exp[x]*(12*x-2*Cos[x]+2*Sin[x]),y[x],x,In
```

$$y(x) \rightarrow e^x(2x^3 - 12x + c_4x + (x + 3 + c_2)\cos(x) + (x - 2 + c_1)\sin(x) + c_3)$$

19.60 problem section 9.3, problem 60

Internal problem ID [1557]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 60.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' - y' + y = e^{2x}(10 + 3x)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$3)-1*diff(y(x),x$2)-1*diff(y(x),x)+1*y(x)=exp(2*x)*(10+3*x),y(x), singsol
```

$$y(x) = (x + 1)e^{2x} + e^{-x}c_2 + e^x(c_3x + c_1)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 34

```
DSolve[y'''[x]-1*y''[x]-1*y'[x]+1*y[x]==Exp[2*x]*(10+3*x),y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow e^{2x}(x + 1) + c_1e^{-x} + e^x(c_3x + c_2)$$

19.61 problem section 9.3, problem 61

Internal problem ID [1558]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 61.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + y'' - 2y = -e^{3x}(17x^2 + 67x + 9)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(diff(y(x),x$3)+1*diff(y(x),x$2)-0*diff(y(x),x)-2*y(x)=-exp(3*x)*(9+67*x+17*x^2),y(x),
```

$$y(x) = -\frac{e^{-x}((x^2 + 2x - 2)e^{4x} - 2c_1e^{2x} - 2c_2 \cos(x) - 2c_3 \sin(x))}{2}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 49

```
DSolve[y'''[x]+1*y''[x]-0*y'[x]-2*y[x]==-Exp[3*x]*(9+67*x+17*x^2),y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow -\frac{1}{2}e^{3x}(x^2 + 2x - 2) + c_3e^x + c_2e^{-x} \cos(x) + c_1e^{-x} \sin(x)$$

19.62 problem section 9.3, problem 62

Internal problem ID [1559]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 62.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 6y'' + 11y' - 6y = e^{2x}(-3x^2 - 4x + 5)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+11*diff(y(x),x)-6*y(x)=exp(2*x)*(5-4*x-3*x^2),y(x), s
```

$$y(x) = e^x(c_3e^{2x} + (x^3 + 2x^2 + c_2 + x)e^x + c_1)$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 37

```
DSolve[y'''[x]-6*y''[x]+11*y'[x]-6*y[x]==Exp[2*x]*(5-4*x-3*x^2),y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow e^x(e^x(x^3 + 2x^2 + x + 4 + c_2) + c_3e^{2x} + c_1)$$

19.63 problem section 9.3, problem 63

Internal problem ID [1560]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 63.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$y''' + 2y'' + y' = -2e^{-x}(6x^2 - 18x + 7)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)+2*diff(y(x),x$2)+1*diff(y(x),x)-0*y(x)=-2*exp(-x)*(7-18*x+6*x^2),y(x),
```

$$y(x) = (x^4 - 2x^3 + x^2 + (-c_1 + 2)x - c_1 - c_2 + 2)e^{-x} + c_3$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 42

```
DSolve[y'''[x]+2*y''[x]+1*y'[x]-0*y[x]==-2*Exp[-x]*(7-18*x+6*x^2),y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow e^{-x}(x^4 - 2x^3 + x^2 - (-2 + c_2)x + 2 - c_1 - c_2) + c_3$$

19.64 problem section 9.3, problem 64

Internal problem ID [1561]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 64.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 3y'' + 3y' - y = (x + 1)e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+3*diff(y(x),x)-y(x)=exp(x)*(1+x),y(x), singsol=all)
```

$$y(x) = \frac{e^x(x^4 + 4x^3 + (24c_3 + 3)x^2 + 24c_2x + 24c_1)}{24}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 37

```
DSolve[y'''[x]-3*y''[x]+3*y'[x]-1*y[x]==Exp[x]*(1+x),y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{24}e^x(x^4 + 4x^3 + 24c_3x^2 + 24c_2x + 24c_1)$$

19.65 problem section 9.3, problem 65

Internal problem ID [1562]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 65.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 2y'' + y = -e^{-x}(3x^2 - 9x + 4)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$4)-0*diff(y(x),x$3)-2*diff(y(x),x$2)+0*diff(y(x),x)+y(x)=-exp(-x)*(4-9*x+
```

$$y(x) = \frac{(-x^4 + 2x^3 + 16c_4x + x^2 + 16c_2)e^{-x}}{16} + e^x(c_3x + c_1)$$

✓ Solution by Mathematica

Time used: 0.113 (sec). Leaf size: 61

```
DSolve[y''''[x]-0*y'''[x]-2*y''[x]+0*y'[x]+1*y[x]==-Exp[-x]*(4-9*x+3*x^2),y[x],x,IncludeSing
```

$$y(x) \rightarrow \frac{1}{32}e^{-x}(-2x^4 + 4x^3 + 2x^2 + x(32c_4e^{2x} - 2 + 32c_2) + 32c_3e^{2x} - 3 + 32c_1)$$

19.66 problem section 9.3, problem 66

Internal problem ID [1563]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 66.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 2y'' - y' - 2y = e^{-2x}((23 - 2x) \cos(x) + (8 - 9x) \sin(x))$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$3)+2*diff(y(x),x$2)-diff(y(x),x)-2*y(x)=exp(-2*x)*((23-2*x)*cos(x)+(8-9*x)*sin(x)),y(x),x,Incl
```

$$y(x) = \frac{(2c_1e^{3x} + (2 + x) \cos(x) + (-4x + 3) \sin(x) + 2c_3e^x + 2c_2) e^{-2x}}{2}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 48

```
DSolve[y'''[x]+2*y''[x]-y'[x]-2*y[x]==Exp[-2*x]*((23-2*x)*Cos[x]+(8-9*x)*Sin[x]),y[x],x,Incl
```

$$y(x) \rightarrow \frac{1}{2}e^{-2x}((3 - 4x) \sin(x) + (x + 2) \cos(x) + 2(c_2e^x + c_3e^{3x} + c_1))$$

19.67 problem section 9.3, problem 67

Internal problem ID [1564]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 67.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 3y''' + 4y'' - 2y' = e^x((28 + 6x) \cos(2x) + (11 - 12x) \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$4)-3*diff(y(x),x$3)+4*diff(y(x),x$2)-2*diff(y(x),x)-0*y(x))=exp(x)*((28+6*
```

$$y(x) = \frac{((-4x \sin(x) + c_2 - c_3) \cos(x) + (c_2 + c_3) \sin(x) + 2c_1 + 3) e^x}{2} + c_4$$

✓ Solution by Mathematica

Time used: 0.715 (sec). Leaf size: 43

```
DSolve[y''''[x]-3*y'''[x]+4*y''[x]-2*y'[x]-0*y[x]==Exp[x]*((28+6*x)*Cos[2*x]+(11-12*x)*Sin[2*
```

$$y(x) \rightarrow \frac{1}{2} e^x ((c_1 + c_2) \sin(x) + \cos(x) (-4x \sin(x) - c_1 + c_2) + 2c_3) + c_4$$

19.68 problem section 9.3, problem 68

Internal problem ID [1565]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 68.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 4y''' + 14y'' - 20y' + 25y = e^x((6x + 2) \cos(2x) + 3 \sin(2x))$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 46

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+14*diff(y(x),x$2)-20*diff(y(x),x)+25*y(x)=exp(x)*((2+
```

$$y(x) = \frac{e^x \left((x^3 + x^2 + (-16c_3 + \frac{63}{2})x - 16c_1 - \frac{41}{4}) \cos(2x) - 16 \left((c_4 + \frac{1}{32})x + c_2 + \frac{1111}{192} \right) \sin(2x) \right)}{16}$$

✓ Solution by Mathematica

Time used: 0.674 (sec). Leaf size: 62

```
DSolve[y''''[x]-4*y'''[x]+14*y''[x]-20*y'[x]+25*y[x]==Exp[x]*((2+6*x)*Cos[2*x]+3*Sin[2*x]),y
```

$$y(x) \rightarrow \frac{1}{256} e^x \left((-16x^3 - 16x^2 + (6 + 256c_4)x + 6 + 256c_3) \cos(2x) + (8(3 + 32c_2)x + 3 + 256c_1) \sin(2x) \right)$$

19.69 problem section 9.3, problem 69

Internal problem ID [1566]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 69.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - 2y'' - 5y' + 6y = 2e^x(1 - 6x)$$

With initial conditions

$$[y(0) = 2, y'(0) = 7, y''(0) = 9]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$3)-2*diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=2*exp(x)*(1-6*x),y(0) = 2, D(y
```

$$y(x) = (e^{5x} + x^2e^{3x} + 2e^{3x} - 1)e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 27

```
DSolve[{y'''[x]-2*y''[x]-5*y'[x]+6*y[x]==2*Exp[x]*(1-6*x),{y[0]==2,y'[0]==7,y''[0]==9}},y[x]
```

$$y(x) \rightarrow e^x(x^2 + 2) - e^{-2x} + e^{3x}$$

19.70 problem section 9.3, problem 70

Internal problem ID [1567]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 70.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' - y'' - y' + y = -e^{-x}(4 - 8x)$$

With initial conditions

$$[y(0) = 2, y'(0) = 0, y''(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$3)-1*diff(y(x),x$2)-1*diff(y(x),x)+1*y(x)=-exp(-x)*(4-8*x),y(0) = 2, D(y
```

$$y(x) = (x^2 + x + 1)e^{-x} - (x - 1)e^x$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 27

```
DSolve[{y'''[x]-1*y''[x]-1*y'[x]+1*y[x]==-Exp[-x]*(4-8*x),{y[0]==2,y'[0]==0,y''[0]==0}},y[x]
```

$$y(x) \rightarrow e^{-x}(x^2 + x - e^{2x}(x - 1) + 1)$$

19.71 problem section 9.3, problem 71

Internal problem ID [1568]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 71.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$4y''' - 3y' - y = e^{-\frac{x}{2}}(-3x + 2)$$

With initial conditions

$$[y(0) = -1, y'(0) = 15, y''(0) = -17]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 22

```
dsolve([4*diff(y(x),x$3)-0*diff(y(x),x$2)-3*diff(y(x),x)-1*y(x)=exp(-x/2)*(2-3*x),y(0) = -1,
```

$$y(x) = \frac{(x^3 + 192x)e^{-\frac{x}{2}}}{12} - e^x$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 35

```
DSolve[{4*y'''[x]-0*y''[x]-3*y'[x]-1*y[x]==Exp[-x/2]*(2-3*x),{y[0]==-1,y'[0]==15,y''[0]==-17}},y,
```

$$y(x) \rightarrow \frac{1}{36}e^{-x/2}(3x^3 + 24x + 8e^{3x/2} + 64)$$

19.72 problem section 9.3, problem 72

Internal problem ID [1569]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 72.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' + 2y''' + 2y'' + 2y' + y = e^{-x}(20 - 12x)$$

With initial conditions

$$[y(0) = 3, y'(0) = -4, y''(0) = 7, y'''(0) = -22]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 31

```
dsolve([diff(y(x),x$4)+2*diff(y(x),x$3)+2*diff(y(x),x$2)+2*diff(y(x),x)+1*y(x)=exp(-x)*(20-12*x),{y(0)=3,y'(0)=-4,y''(0)=7,y'''(0)=-22})
```

$$y(x) = (-x^3 + 2x^2 - x + 2) e^{-x} + \cos(x) - \sin(x)$$

✓ Solution by Mathematica

Time used: 0.085 (sec). Leaf size: 39

```
DSolve[{y''''[x]+2*y'''[x]+2*y''[x]+2*y'[x]+1*y[x]==Exp[-x]*(20-12*x)},{y[0]==3,y'[0]==-4,y''[0]==7,y'''[0]==-22}]
```

$$y(x) \rightarrow e^{-x}(-x^3 + 2x^2 - x - e^x \sin(x) + e^x \cos(x) + 2)$$

19.73 problem section 9.3, problem 73

Internal problem ID [1570]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 73.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _linear, _nonhomogeneous]]`

$$y''' + 2y'' + y' + 2y = 30 \cos(x) - 10 \sin(x)$$

With initial conditions

$$[y(0) = 3, y'(0) = -4, y''(0) = 16]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 25

```
dsolve([0*diff(y(x),x$4)+1*diff(y(x),x$3)+2*diff(y(x),x$2)+1*diff(y(x),x)+2*y(x)=30*cos(x)-10*sin(x),y(0)=3,y'(0)=-4,y''(0)=16])
```

$$y(x) = e^{-2x} + (2 - x) \cos(x) + (7x - 1) \sin(x)$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 26

```
DSolve[{0*y''''[x]+1*y'''[x]+2*y''[x]+1*y'[x]+2*y[x]==30*Cos[x]-10*Sin[x],{y[0]==3,y'[0]==-4,y''[0]==16}],y,x
```

$$y(x) \rightarrow e^{-2x} + (7x - 1) \sin(x) - ((x - 2) \cos(x))$$

19.74 problem section 9.3, problem 74

Internal problem ID [1571]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.3. Undetermined Coefficients for Higher Order Equations. Page 495

Problem number: section 9.3, problem 74.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_y]]`

$$y'''' - 3y''' + 5y'' - 2y' = -2e^x(\cos(x) - \sin(x))$$

With initial conditions

$$[y(0) = 2, y'(0) = 0, y''(0) = -1, y'''(0) = -5]$$

✓ Solution by Maple

Time used: 4.203 (sec). Leaf size: 1300

```
dsolve([1*diff(y(x),x$4)-3*diff(y(x),x$3)+5*diff(y(x),x$2)-2*diff(y(x),x)+0*y(x)=-2*exp(x)*
```

Expression too large to display

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 3484

```
DSolve[{1*y''''[x]-3*y'''[x]+5*y''[x]-2*y'[x]+0*y[x]==-2*Exp[x]*(Cos[x]-Sin[x]),{y[0]==2,y'
```

Too large to display

20 Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

20.1	problem section 9.4, problem 3	778
20.2	problem section 9.4, problem 8	779
20.3	problem section 9.4, problem 11	780
20.4	problem section 9.4, problem 14	781
20.5	problem section 9.4, problem 16	782
20.6	problem section 9.4, problem 18	783
20.7	problem section 9.4, problem 22	784
20.8	problem section 9.4, problem 23	785
20.9	problem section 9.4, problem 25	786
20.10	problem section 9.4, problem 27	787
20.11	problem section 9.4, problem 30	788
20.12	problem section 9.4, problem 32	789
20.13	problem section 9.4, problem 33	790
20.14	problem section 9.4, problem 35	791
20.15	problem section 9.4, problem 36	792
20.16	problem section 9.4, problem 39	793
20.17	problem section 9.4, problem 41	794

20.1 problem section 9.4, problem 3

Internal problem ID [1572]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 3.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 3x^2 y'' + 6y'x - 6y = 2x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=2*x,y(x), singsol=all
```

$$y(x) = \frac{x(2c_3x^2 + 2c_2x + 2\ln(x) + 2c_1 + 3)}{2}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 25

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==2*x,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow x \left(c_3 x^2 + \log(x) + c_2 x + \frac{3}{2} + c_1 \right)$$

20.2 problem section 9.4, problem 8

Internal problem ID [1573]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 8.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$4x^3y''' + 4x^2y'' - 5y'x + 2y = 30x^2$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(4*x^3*diff(y(x),x$3)+4*x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+2*y(x)=30*x^2,y(x), singularities = false)
```

$$y(x) = \frac{(c_1 + 2 \ln(x) - \frac{32}{15}) x^{\frac{5}{2}} + c_3 x + c_2}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 38

```
DSolve[4*x^3*y'''[x]+4*x^2*y''[x]-5*x*y'[x]+2*y[x]==30*x^2,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow 2x^2 \log(x) + \frac{(-\frac{32}{15} + c_3) x^{5/2} + c_2 x + c_1}{\sqrt{x}}$$

20.3 problem section 9.4, problem 11

Internal problem ID [1574]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 11.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _nonhomogeneous]]`

$$x^3 y''' + x^2 y'' - 2y'x + 2y = x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{2x^3 \ln(x) + 6c_3 x^3 + 6c_2 x^2 + c_1}{6x}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 35

```
DSolve[x^3*y'''[x]+x^2*y''[x]-2*x*y'[x]+2*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}x^2 \log(x) + \left(-\frac{4}{9} + c_3\right)x^2 + c_2x + \frac{c_1}{x}$$

20.4 problem section 9.4, problem 14

Internal problem ID [1575]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 14.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$16x^4y'''' + 96x^3y'''' + 72x^2y'' - 24y'x + 9y = 96x^{\frac{5}{2}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(16*x^4*diff(y(x),x$4)+96*x^3*diff(y(x),x$3)+72*x^2*diff(y(x),x$2)-24*x*diff(y(x),x)+9*y(x),x)=96*x^(5/2),y(x),x,Inc
```

$$y(x) = \frac{4c_4x^3 + x^4 + 4c_3x^2 + 4c_2x + 4c_1}{4x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 41

```
DSolve[16*x^4*y''''[x]+96*x^3*y''''[x]+72*x^2*y''[x]-24*x*y'[x]+9*y[x]==96*x^(5/2),y[x],x,Inc
```

$$y(x) \rightarrow \frac{x^4 + 4c_4x^3 + 4c_3x^2 + 4c_2x + 4c_1}{4x^{3/2}}$$

20.5 problem section 9.4, problem 16

Internal problem ID [1576]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 16.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _with_linear_symmetries]]`

$$x^4 y'''' - 4x^3 y''' + 12x^2 y'' - 24y'x + 24y = x^4$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(x^4*diff(y(x),x$4)-4*x^3*diff(y(x),x$3)+12*x^2*diff(y(x),x$2)-24*x*diff(y(x),x)+24*y(x),x)=x^4,y(x),x)
```

$$y(x) = \left(\frac{x^3 \ln(x)}{6} + \left(c_4 - \frac{11}{36} \right) x^3 + c_3 x^2 + c_2 x + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 40

```
DSolve[x^4*y''''[x]-4*x^3*y'''[x]+12*x^2*y''[x]-24*x*y'[x]+24*y[x]==x^4,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{6} x^4 \log(x) + x \left(\left(-\frac{11}{36} + c_4 \right) x^3 + c_3 x^2 + c_2 x + c_1 \right)$$

20.6 problem section 9.4, problem 18

Internal problem ID [1577]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 18.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$x^4 y'''' + 6x^3 y''' + 2x^2 y'' - 4y'x + 4y = 12x^2$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+4*y(x)=
```

$$y(x) = \frac{12x^4 \ln(x) + (12c_2 - 15)x^4 + 12c_3x^3 + 2c_1x + 12c_4}{12x^2}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 38

```
DSolve[x^4*y''''[x]+6*x^3*y'''[x]+2*x^2*y''[x]-4*x*y'[x]+4*y[x]==12*x^2,y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{x^4 \log(x) + \left(-\frac{19}{12} + c_4\right)x^4 + c_3x^3 + c_2x + c_1}{x^2}$$

20.7 problem section 9.4, problem 22

Internal problem ID [1578]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 22.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 2x^2 y'' + 3y'x - 3y = 4x$$

With initial conditions

$$[y(1) = 4, y'(1) = 4, y''(1) = 2]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 23

```
dsolve([x^3*diff(y(x),x$3)-2*x^2*diff(y(x),x$2)+3*x*diff(y(x),x)-3*y(x)=4*x,y(1) = 4, D(y)(1) = 4, D(y)(1) = 2],y(x),x)
```

$$y(x) = x(-\ln(x)^2 + x^2 - 2\ln(x) + 3)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 22

```
DSolve[{x^3*y'''[x]-2*x^2*y''[x]+3*x*y'[x]-3*y[x]==4*x,{y[1]==4,y'[1]==4,y''[1]==2}},y[x],x
```

$$y(x) \rightarrow x(x^2 - \log^2(x) - 2\log(x) + 3)$$

20.8 problem section 9.4, problem 23

Internal problem ID [1579]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 23.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 5x^2 y'' + 14y'x - 18y = x^3$$

With initial conditions

$$[y(1) = 0, y'(1) = 1, y''(1) = 7]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 25

```
dsolve([x^3*diff(y(x),x$3)-5*x^2*diff(y(x),x$2)+14*x*diff(y(x),x)-18*y(x)=x^3,y(1) = 0, D(y
```

$$y(x) = \frac{x^2(\ln(x)^2 x + 4x \ln(x) - 2x + 2)}{2}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 28

```
DSolve[{x^3*y'''[x]-5*x^2*y''[x]+14*x*y'[x]-18*y[x]==x^3,{y[1]==0,y'[1]==1,y''[1]==7}},y[x],
```

$$y(x) \rightarrow \frac{1}{2}x^2(-2x + x \log^2(x) + 4x \log(x) + 2)$$

20.9 problem section 9.4, problem 25

Internal problem ID [1580]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 25.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _with_linear_symmetries]]`

$$x^3 y''' - 6x^2 y'' + 16y'x - 16y = 9x^4$$

With initial conditions

$$[y(1) = 2, y'(1) = 1, y''(1) = 5]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([x^3*diff(y(x),x$3)-6*x^2*diff(y(x),x$2)+16*x*diff(y(x),x)-16*y(x)=9*x^4,y(1) = 2, D
```

$$y(x) = -x^4 + \frac{3 \ln(x)^2 x^4}{2} + 2x^4 \ln(x) + 3x$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 32

```
DSolve[{x^3*y'''[x]-6*x^2*y''[x]+16*x*y'[x]-16*y[x]==9*x^4,{y[1]==2,y'[1]==1,y''[1]==5}},y[x
```

$$y(x) \rightarrow -x^4 + \frac{3}{2}x^4 \log^2(x) + 2x^4 \log(x) + 3x$$

20.10 problem section 9.4, problem 27

Internal problem ID [1581]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 27.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _nonhomogeneous]]`

$$x^3 y''' + x^2 y'' - 2y'x + 2y = x(x + 1)$$

With initial conditions

$$\left[y(-1) = -6, y'(-1) = \frac{43}{6}, y''(-1) = -\frac{5}{2} \right]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 30

```
dsolve([x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=x*(x+1),y(-1) = -6, D
```

$$y(x) = \frac{x(-2i\pi x + 2x \ln(x) + 3i\pi - 3 \ln(x) - 12x + 24)}{6}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 34

```
DSolve[{x^3*y'''[x]+x^2*y''[x]-2*x*y'[x]+2*y[x]==x*(x+1),{y[-1]==-6,y'[-1]==43/6,y''[-1]==-5
```

$$y(x) \rightarrow \frac{1}{6}x(-2i\pi x - 12x + (2x - 3) \log(x) + 3i\pi + 24)$$

20.11 problem section 9.4, problem 30

Internal problem ID [1582]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 30.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$x^4 y'''' + 3x^3 y''' - x^2 y'' + 2y'x - 2y = 9x^2$$

With initial conditions

$$[y(1) = -7, y'(1) = -11, y''(1) = -5, y'''(1) = 6]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 14

```
dsolve([x^4*diff(y(x),x$4)+3*x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=9
```

$$y(x) = x^2(-7 + 3 \ln(x))$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 15

```
DSolve[{x^4*y''''[x]+3*x^3*y'''[x]-x^2*y''[x]+2*x*y'[x]-2*y[x]==9*x^2,{y[1]==-7,y'[1]==-11,y
```

$$y(x) \rightarrow x^2(3 \log(x) - 7)$$

20.12 problem section 9.4, problem 32

Internal problem ID [1583]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 32.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$4x^4y'''' + 24x^3y'''' + 23x^2y'' - y'x + y = 6x$$

With initial conditions

$$\left[y(1) = 2, y'(1) = 0, y''(1) = 4, y'''(1) = -\frac{37}{4} \right]$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 27

```
dsolve([4*x^4*diff(y(x),x$4)+24*x^3*diff(y(x),x$3)+23*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)]=6*x,
```

$$y(x) = \frac{\ln(x)x^{\frac{5}{2}} - x^2 + x^{\frac{5}{2}} + \sqrt{x} + x}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 26

```
DSolve[{4*x^4*y''''[x]+24*x^3*y''''[x]+23*x^2*y''[x]-x*y'[x]+y[x]==6*x,{y[1]==2,y'[1]==0,y''[1]==4,y'''[1]==-37/4}],y[x],x]
```

$$y(x) \rightarrow x - \sqrt{x} + \frac{1}{\sqrt{x}} + \frac{1}{x} + x \log(x)$$

20.13 problem section 9.4, problem 33

Internal problem ID [1584]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 33.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$x^4 y'''' + 5x^3 y''' - 3x^2 y'' - 6y'x + 6y = 40x^3$$

With initial conditions

$$[y(-1) = -1, y'(-1) = -7, y''(-1) = -1, y'''(-1) = -31]$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 30

```
dsolve([x^4*diff(y(x),x$4)+5*x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)-6*x*diff(y(x),x)+6*y(x)
```

$$y(x) = \frac{x^5 \ln(x) - 1 + (-i\pi - 2)x^5 + x^3 + x}{x^2}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 32

```
DSolve[{x^4*y''''[x]+5*x^3*y'''[x]-3*x^2*y''[x]-6*x*y'[x]+6*y[x]==40*x^3,{y[-1]==-1,y'[-1]==
```

$$y(x) \rightarrow \frac{(-2 - i\pi)x^5 + x^5 \log(x) + x^3 + x - 1}{x^2}$$

20.14 problem section 9.4, problem 35

Internal problem ID [1585]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 35.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[_3rd_order, _linear, _nonhomogeneous]`

$$y''' + 2y'' - y' - 2y = F(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 62

```
dsolve(diff(y(x),x$3)+2*diff(y(x),x$2)-diff(y(x),x)-2*y(x)=F(x),y(x), singsol=all)
```

$$y(x) = \frac{\left(\int e^{-x} F(x) dx\right) e^{3x} + 6c_1 e^{3x} - 3\left(\int e^x F(x) dx\right) e^x + 6c_3 e^x + 2\left(\int F(x) e^{2x} dx\right) + 6c_2}{6} e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 96

```
DSolve[y''''[x]+2*y'''[x]-y''[x]-2*y[x]==f[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(\int_1^x \frac{1}{3} e^{2K[1]} f(K[1]) dK[1] + e^x \int_1^x -\frac{1}{2} e^{K[2]} f(K[2]) dK[2] + e^{3x} \int_1^x \frac{1}{6} e^{-K[3]} f(K[3]) dK[3] + c_2 e^x + c_3 e^{3x} + c_1 \right)$$

20.15 problem section 9.4, problem 36

Internal problem ID [1586]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 36.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _nonhomogeneous]]`

$$x^3 y''' + x^2 y'' - 2y'x + 2y = F(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=F(x),y(x), singsol=all)
```

$$y(x) = \left(c_3 + \int \frac{c_2 + \int \frac{c_1 + \int \frac{F(x) dx}{x^3} dx}{x^2} dx}{x^2} dx \right) x^2$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 82

```
DSolve[x^3*y'''[x]+x^2*y''[x]-2*x*y'[x]+2*y[x]==f[x],y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{x^3 \int_1^x \frac{f(K[3])}{3K[3]^3} dK[3] + x^2 \int_1^x -\frac{f(K[2])}{2K[2]^2} dK[2] + \int_1^x \frac{1}{6} f(K[1]) dK[1] + c_3 x^3 + c_2 x^2 + c_1}{x}$$

20.16 problem section 9.4, problem 39

Internal problem ID [1587]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 39.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _linear, _nonhomogeneous]]`

$$y'''' - 5y'' + 4y = F(x)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 84

```
dsolve(diff(y(x),x$4)-5*diff(y(x),x$2)+4*y(x)=F(x),y(x), singsol=all)
```

$$y(x) = \frac{\left(\left(\int e^{-x} F(x) dx \right) e^{3x} - 6c_1 e^{3x} - \frac{\left(\int F(x) e^{-2x} dx \right) e^{4x}}{2} - 6c_4 e^{4x} - \left(\int e^x F(x) dx \right) e^x - 6c_3 e^x + \frac{\left(\int F(x) e^{2x} dx \right)}{2} \right)}{6}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 130

```
DSolve[y''''[x]-5*y''[x]+4*y[x]==f[x],y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-2x} \left(\int_1^x -\frac{1}{12} e^{2K[1]} f(K[1]) dK[1] + e^x \int_1^x \frac{1}{6} e^{K[2]} f(K[2]) dK[2] + e^{3x} \int_1^x -\frac{1}{6} e^{-K[3]} f(K[3]) dK[3] + e^{4x} \int_1^x \frac{1}{12} e^{-2K[4]} f(K[4]) dK[4] + c_2 e^x + c_3 e^{3x} + c_4 e^{4x} + c_1 \right)$$

20.17 problem section 9.4, problem 41

Internal problem ID [1588]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 9 Introduction to Linear Higher Order Equations. Section 9.4. Variation of Parameters for Higher Order Equations. Page 503

Problem number: section 9.4, problem 41.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _exact, _linear, _nonhomogeneous]]`

$$x^4 y'''' + 6x^3 y''' + 2x^2 y'' - 4y'x + 4y = F(x)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(x^4*diff(y(x),x$4)+6*x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+4*y(x)=
```

$$y(x) = \frac{c_4 + \int \left(2c_2 x + c_3 + \int \int \frac{c_1 + \int \frac{F(x) dx}{x^4} dx dx \right) x^2 dx}{x^2}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 104

```
DSolve[x^4*y''''[x]+6*x^3*y'''[x]+2*x^2*y''[x]-4*x*y'[x]+4*y[x]==f[x],y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{x^4 \int_1^x \frac{f(K[4])}{12K[4]^3} dK[4] + x^3 \int_1^x -\frac{f(K[3])}{6K[3]^2} dK[3] + x \int_1^x \frac{1}{6} f(K[2]) dK[2] + \int_1^x -\frac{1}{12} f(K[1]) K[1] dK[1] + c_4 x^4 + c_5}{x^2}$$

21 Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

21.1	problem section 10.4, problem 1	796
21.2	problem section 10.4, problem 2	797
21.3	problem section 10.4, problem 3	798
21.4	problem section 10.4, problem 4	799
21.5	problem section 10.4, problem 5	800
21.6	problem section 10.4, problem 6	801
21.7	problem section 10.4, problem 7	802
21.8	problem section 10.4, problem 8	803
21.9	problem section 10.4, problem 9	804
21.10	problem section 10.4, problem 10	805
21.11	problem section 10.4, problem 11	807
21.12	problem section 10.4, problem 12	808
21.13	problem section 10.4, problem 13	809
21.14	problem section 10.4, problem 14	811
21.15	problem section 10.4, problem 15	812

21.1 problem section 10.4, problem 1

Internal problem ID [1589]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 1.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = y_1(t) + 2y_2(t)$$

$$y_2'(t) = 2y_1(t) + y_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(y__1(t),t)=y__1(t)+2*y__2(t),diff(y__2(t),t)=2*y__1(t)+1*y__2(t)],singsol=all)
```

$$y_1(t) = c_1 e^{3t} + c_2 e^{-t}$$

$$y_2(t) = c_1 e^{3t} - c_2 e^{-t}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 68

```
DSolve[{y1'[t]==y1[t]+2*y2[t],y2'[t]==2*y1[t]+y2[t]},{y1[t],y2[t]},t,IncludeSingularSolution
```

$$y_1(t) \rightarrow \frac{1}{2} e^{-t} (c_1 (e^{4t} + 1) + c_2 (e^{4t} - 1))$$

$$y_2(t) \rightarrow \frac{1}{2} e^{-t} (c_1 (e^{4t} - 1) + c_2 (e^{4t} + 1))$$

21.2 problem section 10.4, problem 2

Internal problem ID [1590]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 2.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -\frac{5y_1(t)}{4} + \frac{3y_2(t)}{4} \\y_2'(t) &= \frac{3y_1(t)}{4} - \frac{5y_2(t)}{4}\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(y__1(t),t)=-5/4*y__1(t)+3/4*y__2(t),diff(y__2(t),t)=3/4*y__1(t)-5/4*y__2(t)],si
```

$$\begin{aligned}y_1(t) &= c_1 e^{-2t} + c_2 e^{-\frac{t}{2}} \\y_2(t) &= -c_1 e^{-2t} + c_2 e^{-\frac{t}{2}}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 76

```
DSolve[{y1'[t]==-5/4*y1[t]+3/4*y2[t],y2'[t]==3/4*y1[t]-5/4*y2[t]},{y1[t],y2[t]},t,IncludeSim
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{2}e^{-2t}(c_1(e^{3t/2} + 1) + c_2(e^{3t/2} - 1)) \\y_2(t) &\rightarrow \frac{1}{2}e^{-2t}(c_1(e^{3t/2} - 1) + c_2(e^{3t/2} + 1))\end{aligned}$$

21.3 problem section 10.4, problem 3

Internal problem ID [1591]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -\frac{4y_1(t)}{5} + \frac{3y_2(t)}{5} \\y_2'(t) &= -\frac{2y_1(t)}{5} - \frac{11y_2(t)}{5}\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(y__1(t),t)=-4/5*y__1(t)+3/5*y__2(t),diff(y__2(t),t)=-2/5*y__1(t)-11/5*y__2(t)],
```

$$\begin{aligned}y_1(t) &= e^{-t}c_1 + c_2e^{-2t} \\y_2(t) &= -\frac{e^{-t}c_1}{3} - 2c_2e^{-2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 65

```
DSolve[{y1'[t]==-4/5*y1[t]+3/5*y2[t],y2'[t]==-2/5*y1[t]-11/5*y2[t]},{y1[t],y2[t]},t,IncludeS
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{5}e^{-2t}(c_1(6e^t - 1) + 3c_2(e^t - 1)) \\y_2(t) &\rightarrow \frac{1}{5}e^{-2t}(-2c_1(e^t - 1) - c_2(e^t - 6))\end{aligned}$$

21.4 problem section 10.4, problem 4

Internal problem ID [1592]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_1(t) - 4y_2(t) \\y_2'(t) &= -y_1(t) - y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve([diff(y__1(t),t)=-1*y__1(t)-4*y__2(t),diff(y__2(t),t)=-1*y__1(t)-1*y__2(t)],singsol=a
```

$$\begin{aligned}y_1(t) &= c_1 e^t + c_2 e^{-3t} \\y_2(t) &= -\frac{c_1 e^t}{2} + \frac{c_2 e^{-3t}}{2}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 71

```
DSolve[{y1'[t]==-1*y1[t]-4*y2[t],y2'[t]==-1*y1[t]-1*y2[t]},{y1[t],y2[t]},t,IncludeSingularSo
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{2}e^{-3t}(c_1(e^{4t} + 1) - 2c_2(e^{4t} - 1)) \\y_2(t) &\rightarrow \frac{1}{4}e^{-3t}(2c_2(e^{4t} + 1) - c_1(e^{4t} - 1))\end{aligned}$$

21.5 problem section 10.4, problem 5

Internal problem ID [1593]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 5.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 2y_1(t) - 4y_2(t)$$

$$y_2'(t) = -y_1(t) - y_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(y__1(t),t)=2*y__1(t)-4*y__2(t),diff(y__2(t),t)=-1*y__1(t)-1*y__2(t)],singsol=al
```

$$y_1(t) = c_1 e^{3t} + c_2 e^{-2t}$$

$$y_2(t) = -\frac{c_1 e^{3t}}{4} + c_2 e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 72

```
DSolve[{y1'[t]==2*y1[t]-4*y2[t],y2'[t]==-1*y1[t]-1*y2[t]},{y1[t],y2[t]},t,IncludeSingularSol
```

$$y_1(t) \rightarrow \frac{1}{5} e^{-2t} (c_1 (4e^{5t} + 1) - 4c_2 (e^{5t} - 1))$$

$$y_2(t) \rightarrow \frac{1}{5} e^{-2t} (c_2 (e^{5t} + 4) - c_1 (e^{5t} - 1))$$

21.6 problem section 10.4, problem 6

Internal problem ID [1594]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 6.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 4y_1(t) - 3y_2(t)$$

$$y_2'(t) = 2y_1(t) - y_2(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve([diff(y__1(t),t)=4*y__1(t)-3*y__2(t),diff(y__2(t),t)=2*y__1(t)-1*y__2(t)],singsol=all
```

$$y_1(t) = c_1 e^t + c_2 e^{2t}$$

$$y_2(t) = c_1 e^t + \frac{2c_2 e^{2t}}{3}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 56

```
DSolve[{y1'[t]==4*y1[t]-3*y2[t],y2'[t]==2*y1[t]-1*y2[t]},{y1[t],y2[t]},t,IncludeSingularSolu
```

$$y_1(t) \rightarrow e^t (c_1 (3e^t - 2) - 3c_2 (e^t - 1))$$

$$y_2(t) \rightarrow e^t (2c_1 (e^t - 1) + c_2 (3 - 2e^t))$$

21.7 problem section 10.4, problem 7

Internal problem ID [1595]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 7.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -6y_1(t) - 3y_2(t)$$

$$y_2'(t) = y_1(t) - 2y_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve([diff(y__1(t),t)=-6*y__1(t)-3*y__2(t),diff(y__2(t),t)=1*y__1(t)-2*y__2(t)],singsol=all)
```

$$y_1(t) = c_1 e^{-3t} + c_2 e^{-5t}$$

$$y_2(t) = -c_1 e^{-3t} - \frac{c_2 e^{-5t}}{3}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 72

```
DSolve[{y1'[t]==-6*y1[t]-3*y2[t],y2'[t]==1*y1[t]-2*y2[t]},{y1[t],y2[t]},t,IncludeSingularSol
```

$$y_1(t) \rightarrow \frac{1}{2} e^{-5t} (-(c_1 (e^{2t} - 3)) - 3c_2 (e^{2t} - 1))$$

$$y_2(t) \rightarrow \frac{1}{2} e^{-5t} (c_1 (e^{2t} - 1) + c_2 (3e^{2t} - 1))$$

21.8 problem section 10.4, problem 8

Internal problem ID [1596]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 8.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= y_1(t) - y_2(t) - 2y_3(t) \\y_2'(t) &= y_1(t) - 2y_2(t) - 3y_3(t) \\y_3'(t) &= -4y_1(t) + y_2(t) - y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 72

```
dsolve([diff(y__1(t),t)=1*y__1(t)-1*y__2(t)-2*y__3(t),diff(y__2(t),t)=1*y__1(t)-2*y__2(t)-3*y__3(t),diff(y__3(t),t)=-4*y__1(t)+y__2(t)-y__3(t))
```

$$\begin{aligned}y_1(t) &= e^{-t}c_1 + c_2e^{2t} + c_3e^{-3t} \\y_2(t) &= 4e^{-t}c_1 + c_2e^{2t} + 2c_3e^{-3t} \\y_3(t) &= -e^{-t}c_1 - c_2e^{2t} + c_3e^{-3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 102

```
DSolve[{y1'[t]==1*y1[t]-1*y2[t]-2*y3[t],y2'[t]==1*y1[t]-2*y2[t]-3*y3[t],y3'[t]==-4*y1[t]+1*y2[t]-y3[t]}
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{576}e^{-3t}(c_1(63 - 128e^t) + c_2(64e^t - 27)) \\y_2(t) &\rightarrow \frac{1}{864}e^{-3t}(c_2(224e^t - 81) - 7c_1(64e^t - 27)) \\y_3(t) &\rightarrow \frac{e^{-3t}(c_1(189 - 128e^t) + c_2(64e^t - 81))}{1728}\end{aligned}$$

21.9 problem section 10.4, problem 9

Internal problem ID [1597]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 9.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -6y_1(t) - 4y_2(t) - 8y_3(t)$$

$$y_2'(t) = -4y_1(t) - 4y_3(t)$$

$$y_3'(t) = -8y_1(t) - 4y_2(t) - 6y_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 67

```
dsolve([diff(y__1(t),t)=-6*y__1(t)-4*y__2(t)-8*y__3(t),diff(y__2(t),t)=-4*y__1(t)-0*y__2(t)-
```

$$y_1(t) = 2c_2e^{-16t} + 2c_3e^{2t} + c_1e^{2t}$$

$$y_2(t) = c_2e^{-16t} + c_3e^{2t}$$

$$y_3(t) = 2c_2e^{-16t} - \frac{5c_3e^{2t}}{2} - c_1e^{2t}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 116

```
DSolve[{y1'[t]==-6*y1[t]-4*y2[t]-8*y3[t],y2'[t]==-4*y1[t]-0*y2[t]-4*y3[t],y1'[t]==-8*y1[t]-4
```

$$y_1(t) \rightarrow \frac{e^{-16t}(c_1(8 - 4913e^{18t}) + 2c_2(4913e^{18t} + 1))}{44217}$$

$$y_2(t) \rightarrow \frac{e^{-16t}(4c_1(4913e^{18t} + 1) + c_2(1 - 39304e^{18t}))}{44217}$$

$$y_3(t) \rightarrow \frac{e^{-16t}(c_1(8 - 4913e^{18t}) + 2c_2(4913e^{18t} + 1))}{44217}$$

21.10 problem section 10.4, problem 10

Internal problem ID [1598]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 10.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 3y_1(t) + 5y_2(t) + 8y_3(t)$$

$$y_2'(t) = y_1(t) - y_2(t) - 2y_3(t)$$

$$y_3'(t) = -y_1(t) - y_2(t) - y_3(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 61

```
dsolve([diff(y__1(t),t)=3*y__1(t)+5*y__2(t)+8*y__3(t),diff(y__2(t),t)=1*y__1(t)-1*y__2(t)-2*
```

$$y_1(t) = c_1 e^t + c_2 e^{-2t} + c_3 e^{2t}$$

$$y_2(t) = 2c_1 e^t - c_2 e^{-2t} + \frac{5c_3 e^{2t}}{7}$$

$$y_3(t) = -\frac{3c_1 e^t}{2} - \frac{4c_3 e^{2t}}{7}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 193

```
DSolve[{y1'[t]==3*y1[t]+5*y2[t]+8*y3[t],y2'[t]==1*y1[t]-1*y2[t]-2*y3[t],y1'[t]==-1*y1[t]-1*y
```

$$y_1(t) \rightarrow \frac{e^{-t/9} \left(\sqrt{35}(2c_2 - 121c_1) \sin\left(\frac{\sqrt{35}t}{9}\right) - 7(74c_1 + 53c_2) \cos\left(\frac{\sqrt{35}t}{9}\right) \right)}{1575}$$
$$y_2(t) \rightarrow \frac{e^{-t/9} \left(7(901c_1 + 202c_2) \cos\left(\frac{\sqrt{35}t}{9}\right) - \sqrt{35}(34c_1 + 379c_2) \sin\left(\frac{\sqrt{35}t}{9}\right) \right)}{4725}$$
$$y_3(t) \rightarrow \frac{e^{-t/9} \left(2\sqrt{35}(92c_1 + 125c_2) \sin\left(\frac{\sqrt{35}t}{9}\right) - 14(251c_1 + 32c_2) \cos\left(\frac{\sqrt{35}t}{9}\right) \right)}{4725}$$

21.11 problem section 10.4, problem 11

Internal problem ID [1599]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 11.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= y_1(t) - y_2(t) + 2y_3(t) \\y_2'(t) &= 12y_1(t) - 4y_2(t) + 10y_3(t) \\y_3'(t) &= -6y_1(t) + y_2(t) - 7y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 73

```
dsolve([diff(y__1(t),t)=1*y__1(t)-1*y__2(t)+2*y__3(t),diff(y__2(t),t)=12*y__1(t)-4*y__2(t)+10*y__3(t),diff(y__3(t),t)=-6*y__1(t)+y__2(t)-7*y__3(t))
```

$$\begin{aligned}y_1(t) &= c_1e^{-2t} + c_2e^{-3t} + c_3e^{-5t} \\y_2(t) &= c_1e^{-2t} + 2c_2e^{-3t} + 3c_3e^{-5t} \\y_3(t) &= -c_1e^{-2t} - c_2e^{-3t} - \frac{3c_3e^{-5t}}{2}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 193

```
DSolve[{y1'[t]==1*y1[t]-1*y2[t]+2*y3[t],y2'[t]==12*y1[t]-4*y2[t]+10*y3[t],y1'[t]==-6*y1[t]+10*y2[t]-7*y3[t]}
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{e^{-7t/6} \left(71(77c_1 - 109c_2) \cos\left(\frac{\sqrt{71}t}{6}\right) + \sqrt{71}(143c_2 - 2479c_1) \sin\left(\frac{\sqrt{71}t}{6}\right) \right)}{340800} \\y_2(t) &\rightarrow \frac{e^{-7t/6} \left(71(2071c_1 - 407c_2) \cos\left(\frac{\sqrt{71}t}{6}\right) - \sqrt{71}(2717c_1 + 5411c_2) \sin\left(\frac{\sqrt{71}t}{6}\right) \right)}{852000} \\y_3(t) &\rightarrow \frac{e^{-7t/6} \left(639(23c_1 + 9c_2) \cos\left(\frac{\sqrt{71}t}{6}\right) + 3\sqrt{71}(937c_1 - 329c_2) \sin\left(\frac{\sqrt{71}t}{6}\right) \right)}{568000}\end{aligned}$$

21.12 problem section 10.4, problem 12

Internal problem ID [1600]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 12.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 4y_1(t) - y_2(t) - 4y_3(t) \\y_2'(t) &= 4y_1(t) - 3y_2(t) - 2y_3(t) \\y_3'(t) &= y_1(t) - y_2(t) - y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 71

```
dsolve([diff(y__1(t),t)=4*y__1(t)-1*y__2(t)-4*y__3(t),diff(y__2(t),t)=4*y__1(t)-3*y__2(t)-2*y__3(t),diff(y__3(t),t)=y__1(t)-y__2(t)-y__3(t)),y__1(0)=1,y__2(0)=1,y__3(0)=1)
```

$$\begin{aligned}y_1(t) &= c_1 e^{3t} + c_2 e^{-t} + c_3 e^{-2t} \\y_2(t) &= \frac{7c_1 e^{3t}}{11} + c_2 e^{-t} + 2c_3 e^{-2t} \\y_3(t) &= \frac{c_1 e^{3t}}{11} + c_2 e^{-t} + c_3 e^{-2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 102

```
DSolve[{y1'[t]==4*y1[t]-1*y2[t]-4*y3[t],y2'[t]==4*y1[t]-3*y2[t]-2*y3[t],y3'[t]==1*y1[t]-1*y2[t]-y3[t],y1[0]==1,y2[0]==1,y3[0]==1},y1,y2,y3,t]
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{216} e^{-2t} (c_1 (54e^t - 8) + c_2 (8 - 27e^t)) \\y_2(t) &\rightarrow \frac{1}{216} e^{-2t} (2c_1 (27e^t - 8) + c_2 (16 - 27e^t)) \\y_3(t) &\rightarrow \frac{1}{216} e^{-2t} (c_1 (54e^t - 8) + c_2 (8 - 27e^t))\end{aligned}$$

21.13 problem section 10.4, problem 13

Internal problem ID [1601]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 13.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -2y_1(t) + 2y_2(t) - 6y_3(t)$$

$$y_2'(t) = 2y_1(t) + 6y_2(t) + 2y_3(t)$$

$$y_3'(t) = -2y_1(t) - 2y_2(t) + 2y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 66

```
dsolve([diff(y__1(t),t)=-2*y__1(t)+2*y__2(t)-6*y__3(t),diff(y__2(t),t)=2*y__1(t)+6*y__2(t)+2
```

$$y_1(t) = c_1 e^{4t} + c_2 e^{-4t} + c_3 e^{6t}$$

$$y_2(t) = -\frac{c_2 e^{-4t}}{4} + c_3 e^{6t}$$

$$y_3(t) = -c_1 e^{4t} + \frac{c_2 e^{-4t}}{4} - c_3 e^{6t}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 257

```
DSolve[{y1'[t]==-2*y1[t]+2*y2[t]-6*y3[t],y2'[t]==2*y1[t]+6*y2[t]+2*y3[t],y1'[t]==-2*y1[t]-2*
```

$y_1(t) \rightarrow$

$$\frac{e^{-\frac{1}{2}(\sqrt{73}-5)t} \left(2c_1 \left((841\sqrt{73} - 7227) e^{\sqrt{73}t} - 7227 - 841\sqrt{73} \right) + c_2 \left((171\sqrt{73} - 1825) e^{\sqrt{73}t} - 1825 - \right. \right.}{598016}$$

$y_2(t)$

$$\rightarrow \frac{e^{-\frac{1}{2}(\sqrt{73}-5)t} \left(c_1 \left((342\sqrt{73} - 3650) e^{\sqrt{73}t} - 3650 - 342\sqrt{73} \right) - c_2 \left((1971 + 143\sqrt{73}) e^{\sqrt{73}t} + 1971 - 143 \right. \right.}{598016}$$

$y_3(t)$

$$\rightarrow \frac{e^{-\frac{1}{2}(\sqrt{73}-5)t} \left(c_1 \left((342\sqrt{73} - 3650) e^{\sqrt{73}t} - 3650 - 342\sqrt{73} \right) - c_2 \left((1971 + 143\sqrt{73}) e^{\sqrt{73}t} + 1971 - 143 \right. \right.}{1196032}$$

21.14 problem section 10.4, problem 14

Internal problem ID [1602]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 14.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 3y_1(t) + 2y_2(t) - 2y_3(t) \\y_2'(t) &= -2y_1(t) + 7y_2(t) - 2y_3(t) \\y_3'(t) &= -10y_1(t) + 10y_2(t) - 5y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 57

```
dsolve([diff(y__1(t),t)=3*y__1(t)+2*y__2(t)-2*y__3(t),diff(y__2(t),t)=-2*y__1(t)+7*y__2(t)-2*y__3(t),diff(y__3(t),t)=-10*y__1(t)+10*y__2(t)-5*y__3(t))
```

$$\begin{aligned}y_1(t) &= c_2 e^{5t} + c_3 e^{-5t} \\y_2(t) &= c_2 e^{5t} + c_3 e^{-5t} + e^{5t} c_1 \\y_3(t) &= 5c_3 e^{-5t} + e^{5t} c_1\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 126

```
DSolve[{y1'[t]==3*y1[t]+2*y2[t]-2*y3[t],y2'[t]==-2*y1[t]+7*y2[t]-2*y3[t],y1'[t]==-10*y1[t]+10*y2[t]-5*y3[t]}
```

$$\begin{aligned}y_1(t) &\rightarrow -\frac{e^{5t}(c_1(432e^{10t/3} - 1331) + c_2(1331 - 216e^{10t/3}))}{85184} \\y_2(t) &\rightarrow -\frac{e^{5t}(c_1(216e^{10t/3} - 1331) + c_2(1331 - 108e^{10t/3}))}{42592} \\y_3(t) &\rightarrow \frac{e^{5t}(c_1(720e^{10t/3} + 1331) - c_2(360e^{10t/3} + 1331))}{85184}\end{aligned}$$

21.15 problem section 10.4, problem 15

Internal problem ID [1603]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.4, constant coefficient homogeneous system. Page 540

Problem number: section 10.4, problem 15.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 3y_1(t) + y_2(t) - y_3(t) \\y_2'(t) &= 3y_1(t) + 5y_2(t) + y_3(t) \\y_3'(t) &= -6y_1(t) + 2y_2(t) + 4y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 53

```
dsolve([diff(y__1(t),t)=3*y__1(t)+1*y__2(t)-1*y__3(t),diff(y__2(t),t)=3*y__1(t)+5*y__2(t)+1*y__3(t),diff(y__3(t),t)=-6*y__1(t)+2*y__2(t)+4*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$\begin{aligned}y_1(t) &= c_2 + c_3 e^{6t} \\y_2(t) &= -c_2 - c_3 e^{6t} + e^{6t} c_1 \\y_3(t) &= -4c_3 e^{6t} + 2c_2 + e^{6t} c_1\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 93

```
DSolve[{y1'[t]==3*y1[t]+1*y2[t]-1*y3[t],y2'[t]==3*y1[t]+5*y2[t]+1*y3[t],y3'[t]==-6*y1[t]+2*y2[t]+4*y3[t]},y1[t],y2[t],y3[t]]
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{625}(-c_1(e^{6t} - 500) - c_2(e^{6t} + 125)) \\y_2(t) &\rightarrow \frac{1}{625}(c_2(125 - 4e^{6t}) - 4c_1(e^{6t} + 125)) \\y_3(t) &\rightarrow \frac{1}{625}(-c_1(e^{6t} - 1000) - c_2(e^{6t} + 250))\end{aligned}$$

22 Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

22.1	problem section 10.5, problem 1	814
22.2	problem section 10.5, problem 2	815
22.3	problem section 10.5, problem 3	816
22.4	problem section 10.5, problem 4	817
22.5	problem section 10.5, problem 5	818
22.6	problem section 10.5, problem 6	819
22.7	problem section 10.5, problem 7	820
22.8	problem section 10.5, problem 8	821
22.9	problem section 10.5, problem 9	822
22.10	problem section 10.5, problem 10	823
22.11	problem section 10.5, problem 11	824
22.12	problem section 10.5, problem 12	825
22.13	problem section 10.5, problem 13	826
22.14	problem section 10.5, problem 14	827
22.15	problem section 10.5, problem 15	828
22.16	problem section 10.5, problem 16	829
22.17	problem section 10.5, problem 17	830
22.18	problem section 10.5, problem 18	831
22.19	problem section 10.5, problem 19	832
22.20	problem section 10.5, problem 20	833
22.21	problem section 10.5, problem 21	835
22.22	problem section 10.5, problem 22	836
22.23	problem section 10.5, problem 23	837
22.24	problem section 10.5, problem 24	838
22.25	problem section 10.5, problem 25	839
22.26	problem section 10.5, problem 26	840
22.27	problem section 10.5, problem 27	841
22.28	problem section 10.5, problem 28	842
22.29	problem section 10.5, problem 29	843
22.30	problem section 10.5, problem 30	844
22.31	problem section 10.5, problem 31	845
22.32	problem section 10.5, problem 32	846

22.1 problem section 10.5, problem 1

Internal problem ID [1604]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 1.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 3y_1(t) + 4y_2(t) \\y_2'(t) &= -y_1(t) + 7y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve([diff(y__1(t),t)=3*y__1(t)+4*y__2(t),diff(y__2(t),t)=-1*y__1(t)+7*y__2(t)],singsol=al
```

$$\begin{aligned}y_1(t) &= e^{5t}(c_2t + c_1) \\y_2(t) &= \frac{e^{5t}(2c_2t + 2c_1 + c_2)}{4}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{y1'[t]==3*y1[t]+4*y2[t],y2'[t]==-1*y1[t]+7*y2[t]},{y1[t],y2[t]},t,IncludeSingularSol
```

$$\begin{aligned}y_1(t) &\rightarrow e^{5t}(-2c_1t + 4c_2t + c_1) \\y_2(t) &\rightarrow e^{5t}(c_1(-t) + 2c_2t + c_2)\end{aligned}$$

22.2 problem section 10.5, problem 2

Internal problem ID [1605]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 2.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_2(t) \\ y_2'(t) &= y_1(t) - 2y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve([diff(y__1(t),t)=0*y__1(t)-1*y__2(t),diff(y__2(t),t)=1*y__1(t)-2*y__2(t)],singsol=all
```

$$\begin{aligned}y_1(t) &= e^{-t}(c_2t + c_1) \\ y_2(t) &= e^{-t}(c_2t + c_1 - c_2)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 44

```
DSolve[{y1'[t]==0*y1[t]-1*y2[t],y2'[t]==1*y1[t]-2*y2[t]},{y1[t],y2[t]},t,IncludeSingularSolu
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-t}(c_1(t + 1) - c_2t) \\ y_2(t) &\rightarrow e^{-t}((c_1 - c_2)t + c_2)\end{aligned}$$

22.3 problem section 10.5, problem 3

Internal problem ID [1606]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 3.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -7y_1(t) + 4y_2(t)$$

$$y_2'(t) = -y_1(t) - 11y_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(y__1(t),t)=-7*y__1(t)+4*y__2(t),diff(y__2(t),t)=-1*y__1(t)-11*y__2(t)],singsol=
```

$$y_1(t) = e^{-9t}(c_2t + c_1)$$

$$y_2(t) = -\frac{e^{-9t}(2c_2t + 2c_1 - c_2)}{4}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{y1'[t]==-7*y1[t]+4*y2[t],y2'[t]==-1*y1[t]-11*y2[t]},{y1[t],y2[t]},t,IncludeSingularS
```

$$y1(t) \rightarrow e^{-9t}(2c_1t + 4c_2t + c_1)$$

$$y2(t) \rightarrow e^{-9t}(c_2 - (c_1 + 2c_2)t)$$

22.4 problem section 10.5, problem 4

Internal problem ID [1607]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 3y_1(t) + y_2(t) \\y_2'(t) &= -y_1(t) + y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve([diff(y__1(t),t)=3*y__1(t)+1*y__2(t),diff(y__2(t),t)=-1*y__1(t)+1*y__2(t)],singsol=al
```

$$\begin{aligned}y_1(t) &= e^{2t}(c_2t + c_1) \\y_2(t) &= -e^{2t}(c_2t + c_1 - c_2)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 42

```
DSolve[{y1'[t]==3*y1[t]+1*y2[t],y2'[t]==-1*y1[t]+1*y2[t]},{y1[t],y2[t]},t,IncludeSingularSol
```

$$\begin{aligned}y_1(t) &\rightarrow e^{2t}(c_1(t + 1) + c_2t) \\y_2(t) &\rightarrow e^{2t}(c_2 - (c_1 + c_2)t)\end{aligned}$$

22.5 problem section 10.5, problem 5

Internal problem ID [1608]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 5.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 4y_1(t) + 12y_2(t) \\y_2'(t) &= -3y_1(t) - 8y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(y__1(t),t)=4*y__1(t)+12*y__2(t),diff(y__2(t),t)=-3*y__1(t)-8*y__2(t)],singsol=a
```

$$\begin{aligned}y_1(t) &= e^{-2t}(c_2t + c_1) \\y_2(t) &= -\frac{e^{-2t}(6c_2t + 6c_1 - c_2)}{12}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{y1'[t]==4*y1[t]+12*y2[t],y2'[t]==-3*y1[t]-8*y2[t]},{y1[t],y2[t]},t,IncludeSingularSo
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-2t}(6c_1t + 12c_2t + c_1) \\y_2(t) &\rightarrow e^{-2t}(c_2 - 3(c_1 + 2c_2)t)\end{aligned}$$

22.6 problem section 10.5, problem 6

Internal problem ID [1609]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 6.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -10y_1(t) + 9y_2(t) \\y_2'(t) &= -4y_1(t) + 2y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve([diff(y__1(t),t)=-10*y__1(t)+9*y__2(t),diff(y__2(t),t)=-4*y__1(t)+2*y__2(t)],singsol=
```

$$\begin{aligned}y_1(t) &= e^{-4t}(c_2t + c_1) \\y_2(t) &= \frac{e^{-4t}(6c_2t + 6c_1 + c_2)}{9}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{y1'[t]==-10*y1[t]+9*y2[t],y2'[t]==-4*y1[t]+2*y2[t]},{y1[t],y2[t]},t,IncludeSingularS
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-4t}(-6c_1t + 9c_2t + c_1) \\y_2(t) &\rightarrow e^{-4t}(-4c_1t + 6c_2t + c_2)\end{aligned}$$

22.7 problem section 10.5, problem 7

Internal problem ID [1610]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 7.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -13y_1(t) + 16y_2(t)$$

$$y_2'(t) = -9y_1(t) + 11y_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve([diff(y__1(t),t)=-13*y__1(t)+16*y__2(t),diff(y__2(t),t)=-9*y__1(t)+11*y__2(t)],singso
```

$$y_1(t) = e^{-t}(c_2t + c_1)$$

$$y_2(t) = \frac{e^{-t}(12c_2t + 12c_1 + c_2)}{16}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{y1'[t]==-13*y1[t]+16*y2[t],y2'[t]==-9*y1[t]+11*y2[t]},{y1[t],y2[t]},t,IncludeSingula
```

$$y_1(t) \rightarrow e^{-t}(-12c_1t + 16c_2t + c_1)$$

$$y_2(t) \rightarrow e^{-t}(-9c_1t + 12c_2t + c_2)$$

22.8 problem section 10.5, problem 8

Internal problem ID [1611]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 8.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 2y_2(t) + y_3(t) \\y_2'(t) &= -4y_1(t) + 6y_2(t) + y_3(t) \\y_3'(t) &= 4y_2(t) + 2y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 71

```
dsolve([diff(y__1(t),t)=0*y__1(t)+2*y__2(t)+1*y__3(t),diff(y__2(t),t)=-4*y__1(t)+6*y__2(t)+1*y__3(t),diff(y__3(t),t)=4*y__2(t)+2*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$\begin{aligned}y_1(t) &= \frac{c_2 e^{4t}}{2} + \frac{c_3 e^{4t} t}{2} - \frac{c_1}{2} \\y_2(t) &= \frac{c_2 e^{4t}}{2} + \frac{c_3 e^{4t} t}{2} + \frac{c_3 e^{4t}}{4} - \frac{c_1}{2} \\y_3(t) &= c_1 + c_2 e^{4t} + c_3 e^{4t} t\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 131

```
DSolve[{y1'[t]==0*y1[t]+2*y2[t]+1*y3[t],y2'[t]==-4*y1[t]+6*y2[t]+1*y3[t],y3'[t]==0*y1[t]+4*y2[t]+2*y3[t]},y1[t],y2[t],y3[t]]
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{4}(c_1(e^{4t}(2-8t)+2)+e^{4t}(8c_2t+c_3)-c_3) \\y_2(t) &\rightarrow \frac{1}{4}(-2c_1(e^{4t}(4t+1)-1)+e^{4t}(c_2(8t+4)+c_3)-c_3) \\y_3(t) &\rightarrow c_1(e^{4t}(1-4t)-1)+\frac{1}{2}(e^{4t}(8c_2t+c_3)+c_3)\end{aligned}$$

22.9 problem section 10.5, problem 9

Internal problem ID [1612]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 9.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= \frac{y_1(t)}{3} + \frac{y_2(t)}{3} - y_3(t) \\y_2'(t) &= -\frac{4y_1(t)}{3} - \frac{4y_2(t)}{3} + y_3(t) \\y_3'(t) &= -\frac{2y_1(t)}{3} + \frac{y_2(t)}{3}\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 75

```
dsolve([diff(y__1(t),t)=1/3*y__1(t)+1/3*y__2(t)-1*y__3(t),diff(y__2(t),t)=-4/3*y__1(t)-4/3*y
```

$$\begin{aligned}y_1(t) &= -c_1e^t + c_2e^{-t} + c_3e^{-t}t \\y_2(t) &= c_1e^t - c_2e^{-t} - c_3e^{-t}t + 3c_3e^{-t} \\y_3(t) &= c_1e^t + c_2e^{-t} + c_3e^{-t}t\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 136

```
DSolve[{y1'[t]==1/3*y1[t]+1/3*y2[t]-1*y3[t],y2'[t]==-4/3*y1[t]-4/3*y2[t]+1*y3[t],y3'[t]==-2/
```

$$\begin{aligned}y_1(t) &\rightarrow \frac{1}{6}e^{-t}(c_1(2t + 3e^{2t} + 3) + 2c_2t - 3c_3(e^{2t} - 1)) \\y_2(t) &\rightarrow \frac{1}{6}e^{-t}(c_1(-2t - 3e^{2t} + 3) - 2c_2(t - 3) + 3c_3(e^{2t} - 1)) \\y_3(t) &\rightarrow \frac{1}{6}e^{-t}(c_1(2t - 3e^{2t} + 3) + 2c_2t + 3c_3(e^{2t} + 1))\end{aligned}$$

22.10 problem section 10.5, problem 10

Internal problem ID [1613]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 10.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_1(t) + y_2(t) - y_3(t) \\y_2'(t) &= -2y_1(t) + 2y_3(t) \\y_3'(t) &= -y_1(t) + 3y_2(t) - y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 63

```
dsolve([diff(y__1(t),t)=-1*y__1(t)+1*y__2(t)-1*y__3(t),diff(y__2(t),t)=-2*y__1(t)+0*y__2(t)+
```

$$\begin{aligned}y_1(t) &= (2c_2t + c_1)e^{-2t} \\y_2(t) &= c_2e^{-2t} + c_3e^{2t} \\y_3(t) &= 2c_2e^{-2t}t + c_3e^{2t} + c_1e^{-2t} - c_2e^{-2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 125

```
DSolve[{y1'[t]==-1*y1[t]+1*y2[t]-1*y3[t],y2'[t]==-2*y1[t]+0*y2[t]+2*y3[t],y3'[t]==-1*y1[t]+3
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-2t}(c_1(t+1) + (c_2 - c_3)t) \\y_2(t) &\rightarrow \frac{1}{2}e^{-2t}(-(c_1(e^{4t} - 1)) + c_2(e^{4t} + 1) + c_3(e^{4t} - 1)) \\y_3(t) &\rightarrow \frac{1}{2}e^{-2t}(c_1(2t - e^{4t} + 1) + c_2(2t + e^{4t} - 1) + c_3(-2t + e^{4t} + 1))\end{aligned}$$

22.11 problem section 10.5, problem 11

Internal problem ID [1614]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 11.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 4y_1(t) - 2y_2(t) - 2y_3(t)$$

$$y_2'(t) = -2y_1(t) + 3y_2(t) - y_3(t)$$

$$y_3'(t) = 2y_1(t) - y_2(t) + 3y_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 69

```
dsolve([diff(y__1(t),t)=4*y__1(t)-2*y__2(t)-2*y__3(t),diff(y__2(t),t)=-2*y__1(t)+3*y__2(t)-1*y__3(t),diff(y__3(t),t)=2*y__1(t)-y__2(t)+3*y__3(t))
```

$$y_1(t) = c_2 e^{4t} + c_3 e^{2t}$$

$$y_2(t) = -2c_2 e^{4t} t + c_1 e^{4t} + \frac{3c_3 e^{2t}}{2}$$

$$y_3(t) = 2c_2 e^{4t} t - c_1 e^{4t} - \frac{c_3 e^{2t}}{2}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 126

```
DSolve[{y1'[t]==4*y1[t]-2*y2[t]-2*y3[t],y2'[t]==-2*y1[t]+3*y2[t]-1*y3[t],y3'[t]==2*y1[t]-1*y2[t]+3*y3[t]}
```

$$y_1(t) \rightarrow e^{2t}((c_1 - c_2 - c_3)e^{2t} + c_2 + c_3)$$

$$y_2(t) \rightarrow \frac{1}{2}(3(c_2 + c_3)e^{2t} - e^{4t}(4(c_1 - c_2 - c_3)t + c_2 + 3c_3))$$

$$y_3(t) \rightarrow -\frac{1}{2}e^{2t}(-e^{2t}(4(c_1 - c_2 - c_3)t + c_2 + 3c_3) + c_2 + c_3)$$

22.12 problem section 10.5, problem 12

Internal problem ID [1615]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 12.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 6y_1(t) - 5y_2(t) + 3y_3(t)$$

$$y_2'(t) = 2y_1(t) - y_2(t) + 3y_3(t)$$

$$y_3'(t) = 2y_1(t) + y_2(t) + y_3(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 86

```
dsolve([diff(y__1(t),t)=6*y__1(t)-5*y__2(t)+3*y__3(t),diff(y__2(t),t)=2*y__1(t)-1*y__2(t)+3*
```

$$y_1(t) = c_1 e^{-2t} + c_2 e^{4t} + c_3 e^{4t} t$$

$$y_2(t) = c_1 e^{-2t} + c_2 e^{4t} + c_3 e^{4t} t - \frac{c_3 e^{4t}}{2}$$

$$y_3(t) = -c_1 e^{-2t} + c_2 e^{4t} + c_3 e^{4t} t - \frac{c_3 e^{4t}}{2}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 127

```
DSolve[{y1'[t]==6*y1[t]-5*y2[t]+3*y3[t],y2'[t]==2*y1[t]-1*y2[t]+3*y3[t],y3'[t]==2*y1[t]+1*y2
```

$$y_1(t) \rightarrow \frac{1}{2} e^{-2t} (e^{6t} (c_1 (4t + 2) - c_2 (4t + 1) + c_3) + c_2 - c_3)$$

$$y_2(t) \rightarrow \frac{1}{2} e^{-2t} (e^{6t} (4(c_1 - c_2)t + c_2 + c_3) + c_2 - c_3)$$

$$y_3(t) \rightarrow \frac{1}{2} e^{-2t} (e^{6t} (4(c_1 - c_2)t + c_2 + c_3) - c_2 + c_3)$$

22.13 problem section 10.5, problem 13

Internal problem ID [1616]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 13.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -11y_1(t) + 8y_2(t) \\y_2'(t) &= -2y_1(t) - 3y_2(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = 6, y_2(0) = 2]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve([diff(y__1(t),t) = -11*y__1(t)+8*y__2(t), diff(y__2(t),t) = -2*y__1(t)-3*y__2(t), y__
```

$$\begin{aligned}y_1(t) &= e^{-7t}(-8t + 6) \\y_2(t) &= \frac{e^{-7t}(-32t + 16)}{8}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[{y1'[t]==-11*y1[t]+8*y2[t],y2'[t]==-2*y1[t]-3*y2[t]},{y1[0]==6,y2[0]==2},{y1[t],y2[t]
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-7t}(6 - 8t) \\y_2(t) &\rightarrow e^{-7t}(2 - 4t)\end{aligned}$$

22.14 problem section 10.5, problem 14

Internal problem ID [1617]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 14.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 15y_1(t) - 9y_2(t)$$

$$y_2'(t) = 16y_1(t) - 9y_2(t)$$

With initial conditions

$$[y_1(0) = 5, y_2(0) = 8]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve([diff(y__1(t),t) = 15*y__1(t)-9*y__2(t), diff(y__2(t),t) = 16*y__1(t)-9*y__2(t), y__1
```

$$y_1(t) = e^{3t}(-12t + 5)$$

$$y_2(t) = \frac{e^{3t}(-144t + 72)}{9}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[{y1'[t]==15*y1[t]-9*y2[t],y2'[t]==16*y1[t]-9*y2[t]},{y1[0]==5,y2[0]==8},{y1[t],y2[t]}
```

$$y1(t) \rightarrow e^{3t}(5 - 12t)$$

$$y2(t) \rightarrow -8e^{3t}(2t - 1)$$

22.15 problem section 10.5, problem 15

Internal problem ID [1618]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 15.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -3y_1(t) - 4y_2(t) \\y_2'(t) &= y_1(t) - 7y_2(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = 2, y_2(0) = 3]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve([diff(y__1(t),t) = -3*y__1(t)-4*y__2(t), diff(y__2(t),t) = y__1(t)-7*y__2(t), y__1(0)
```

$$\begin{aligned}y_1(t) &= e^{-5t}(-8t + 2) \\y_2(t) &= \frac{e^{-5t}(-16t + 12)}{4}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[{y1'[t]==-3*y1[t]-4*y2[t],y2'[t]==1*y1[t]-7*y2[t]},{y1[0]==2,y2[0]==3},{y1[t],y2[t]},
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-5t}(2 - 8t) \\y_2(t) &\rightarrow e^{-5t}(3 - 4t)\end{aligned}$$

22.16 problem section 10.5, problem 16

Internal problem ID [1619]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 16.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -7y_1(t) + 24y_2(t) \\y_2'(t) &= -6y_1(t) + 17y_2(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = 3, y_2(0) = 1]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve([diff(y__1(t),t) = -7*y__1(t)+24*y__2(t), diff(y__2(t),t) = -6*y__1(t)+17*y__2(t), y_
```

$$\begin{aligned}y_1(t) &= e^{5t}(-12t + 3) \\y_2(t) &= \frac{e^{5t}(-144t + 24)}{24}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[{y1'[t]==-7*y1[t]+24*y2[t],y2'[t]==-6*y1[t]+17*y2[t]},{y1[0]==3,y2[0]==1},{y1[t],y2[t]
```

$$\begin{aligned}y1(t) &\rightarrow -3e^{5t}(4t - 1) \\y2(t) &\rightarrow e^{5t}(1 - 6t)\end{aligned}$$

22.17 problem section 10.5, problem 17

Internal problem ID [1620]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 17.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -7y_1(t) + 3y_2(t)$$

$$y_2'(t) = -3y_1(t) - y_2(t)$$

With initial conditions

$$[y_1(0) = 0, y_2(0) = 2]$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve([diff(y__1(t),t) = -7*y__1(t)+3*y__2(t), diff(y__2(t),t) = -3*y__1(t)-y__2(t), y__1(0)=0, y__2(0)=2])
```

$$y_1(t) = 6e^{-4t}$$

$$y_2(t) = \frac{e^{-4t}(18t + 6)}{3}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 27

```
DSolve[{y1'[t]==-7*y1[t]+3*y2[t],y2'[t]==-3*y1[t]-1*y2[t]},{y1[0]==0,y2[0]==2},{y1[t],y2[t]}
```

$$y1(t) \rightarrow 6e^{-4t}$$

$$y2(t) \rightarrow e^{-4t}(6t + 2)$$

22.18 problem section 10.5, problem 18

Internal problem ID [1621]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 18.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_1(t) + y_2(t) \\y_2'(t) &= y_1(t) - y_2(t) - 2y_3(t) \\y_3'(t) &= -y_1(t) - y_2(t) - y_3(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = 6, y_2(0) = 5, y_3(0) = -7]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 57

```
dsolve([diff(y__1(t),t) = -y__1(t)+y__2(t), diff(y__2(t),t) = y__1(t)-y__2(t)-2*y__3(t), dif
```

$$\begin{aligned}y_1(t) &= 4e^t + 2e^{-2t} - e^{-2t}t \\y_2(t) &= 8e^t - 3e^{-2t} + e^{-2t}t \\y_3(t) &= -6e^t - e^{-2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 58

```
DSolve[{y1'[t]==-1*y1[t]+1*y2[t]+0*y3[t],y2'[t]==1*y1[t]-1*y2[t]-2*y3[t],y3'[t]==-1*y1[t]-1*
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-2t}(-t + 4e^{3t} + 2) \\y_2(t) &\rightarrow e^{-2t}(t + 8e^{3t} - 3) \\y_3(t) &\rightarrow -e^{-2t} - 6e^t\end{aligned}$$

22.19 problem section 10.5, problem 19

Internal problem ID [1622]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 19.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -2y_1(t) + 2y_2(t) + y_3(t) \\y_2'(t) &= -2y_1(t) + 2y_2(t) + y_3(t) \\y_3'(t) &= -3y_1(t) + 3y_2(t) + 2y_3(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = -6, y_2(0) = -2, y_3(0) = 0]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

```
dsolve([diff(y__1(t),t) = -2*y__1(t)+2*y__2(t)+y__3(t), diff(y__2(t),t) = -2*y__1(t)+2*y__2(t)
```

$$\begin{aligned}y_1(t) &= -9 + 2t + 3e^{2t} \\y_2(t) &= 3e^{2t} - 5 + 2t \\y_3(t) &= -6 + 6e^{2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 44

```
DSolve[{y1'[t]==-2*y1[t]+2*y2[t]+1*y3[t],y2'[t]==-2*y1[t]+2*y2[t]+1*y3[t],y3'[t]==-3*y1[t]+3
```

$$\begin{aligned}y_1(t) &\rightarrow 2t + 3e^{2t} - 9 \\y_2(t) &\rightarrow 2t + 3e^{2t} - 5 \\y_3(t) &\rightarrow 6(e^{2t} - 1)\end{aligned}$$

22.20 problem section 10.5, problem 20

Internal problem ID [1623]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 20.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -7y_1(t) - 4y_2(t) + 4y_3(t)$$

$$y_2'(t) = y_1(t) + y_3(t)$$

$$y_3'(t) = -9y_1(t) - 5y_2(t) + 6y_3(t)$$

With initial conditions

$$[y_1(0) = -6, y_2(0) = 9, y_3(0) = -1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 80

```
dsolve([diff(y__1(t),t) = -7*y__1(t)-4*y__2(t)+4*y__3(t), diff(y__2(t),t) = y__1(t)+y__3(t),
```

$$y_1(t) = -\frac{4e^{-t}\left(\frac{13\sin(2t)}{2} + \frac{39\cos(2t)}{2}\right)}{13}$$
$$y_2(t) = \frac{9e^t}{2} - \frac{7e^{-t}\sin(2t)}{2} + \frac{9e^{-t}\cos(2t)}{2}$$
$$y_3(t) = \frac{9e^t}{2} - \frac{7e^{-t}\sin(2t)}{2} - \frac{11e^{-t}\cos(2t)}{2}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 50

```
DSolve[{y1'[t]==-7*y1[t]-4*y2[t]+4*y3[t],y2'[t]==-1*y1[t]-0*y2[t]+1*y3[t],y3'[t]==-9*y1[t]-5
```

$$y_1(t) \rightarrow -2e^{-3t} - 4e^t$$

$$y_2(t) \rightarrow e^t(9 - 4t)$$

$$y_3(t) \rightarrow e^t(1 - 4t) - 2e^{-3t}$$

22.21 problem section 10.5, problem 21

Internal problem ID [1624]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 21.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_1(t) - 4y_2(t) - y_3(t) \\y_2'(t) &= 3y_1(t) + 6y_2(t) + y_3(t) \\y_3'(t) &= -3y_1(t) - 2y_2(t) + 3y_3(t)\end{aligned}$$

With initial conditions

$$[y_1(0) = -2, y_2(0) = 1, y_3(0) = 3]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 63

```
dsolve([diff(y__1(t),t) = -y__1(t)-4*y__2(t)-y__3(t), diff(y__2(t),t) = 3*y__1(t)+6*y__2(t)+
```

$$\begin{aligned}y_1(t) &= -2e^{4t} + 3e^{2t} \\y_2(t) &= 2e^{4t} - 3e^{2t}t - e^{2t} \\y_3(t) &= 2e^{4t} + 3e^{2t}t + e^{2t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 63

```
DSolve[{y1'[t]==-1*y1[t]-4*y2[t]-1*y3[t],y2'[t]==3*y1[t]+6*y2[t]+1*y3[t],y3'[t]==-3*y1[t]-2*
```

$$\begin{aligned}y_1(t) &\rightarrow 3e^{2t}t - 2e^{4t} \\y_2(t) &\rightarrow e^{2t}(-3t + 2e^{2t} - 1) \\y_3(t) &\rightarrow e^{2t}(3t + 2e^{2t} + 1)\end{aligned}$$

22.22 problem section 10.5, problem 22

Internal problem ID [1625]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 22.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 4y_1(t) - 8y_2(t) - 4y_3(t)$$

$$y_2'(t) = -3y_1(t) - y_2(t) - 4y_3(t)$$

$$y_3'(t) = y_1(t) - y_2(t) + 9y_3(t)$$

With initial conditions

$$[y_1(0) = -4, y_2(0) = 1, y_3(0) = -3]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 62

```
dsolve([diff(y__1(t),t) = 4*y__1(t)-8*y__2(t)-4*y__3(t), diff(y__2(t),t) = -3*y__1(t)-y__2(t)
```

$$y_1(t) = -\frac{50e^{7t}}{11} - \frac{164e^{-4t}}{143} + \frac{22e^{9t}}{13}$$
$$y_2(t) = \frac{5e^{7t}}{11} - \frac{164e^{-4t}}{143} + \frac{22e^{9t}}{13}$$
$$y_3(t) = \frac{5e^{7t}}{2} - \frac{11e^{9t}}{2}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 57

```
DSolve[{y1'[t]==4*y1[t]-8*y2[t]-4*y3[t],y2'[t]==-3*y1[t]-1*y2[t]-3*y3[t],y3'[t]==1*y1[t]-1*y
```

$$y_1(t) \rightarrow e^{8t}(8t - 3) - e^{-4t}$$

$$y_2(t) \rightarrow 2e^{8t} - e^{-4t}$$

$$y_3(t) \rightarrow -e^{8t}(8t + 3)$$

22.23 problem section 10.5, problem 23

Internal problem ID [1626]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 23.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -5y_1(t) - y_2(t) + 11y_3(t)$$

$$y_2'(t) = -7y_1(t) + y_2(t) + 13y_3(t)$$

$$y_3'(t) = -4y_1(t) + 8y_3(t)$$

With initial conditions

$$[y_1(0) = 0, y_2(0) = 2, y_3(0) = 2]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve([diff(y__1(t),t) = -5*y__1(t)-y__2(t)+11*y__3(t), diff(y__2(t),t) = -7*y__1(t)+y__2(t)
```

$$y_1(t) = -3 + 3e^{4t} + 8t$$

$$y_2(t) = 6e^{4t} - 4 + 4t$$

$$y_3(t) = -1 + 4t + 3e^{4t}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 47

```
DSolve[{y1'[t]==-5*y1[t]-1*y2[t]+11*y3[t],y2'[t]==-7*y1[t]+1*y2[t]+13*y3[t],y3'[t]==-4*y1[t]
```

$$y1(t) \rightarrow 8t + 3e^{4t} - 3$$

$$y2(t) \rightarrow 4t + 6e^{4t} - 4$$

$$y3(t) \rightarrow 4t + 3e^{4t} - 1$$

22.24 problem section 10.5, problem 24

Internal problem ID [1627]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 24.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 5y_1(t) - y_2(t) + y_3(t) \\y_2'(t) &= -y_1(t) + 9y_2(t) - 3y_3(t) \\y_3'(t) &= -2y_1(t) + 2y_2(t) + 4y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 65

```
dsolve([diff(y__1(t),t)=5*y__1(t)-1*y__2(t)+1*y__3(t),diff(y__2(t),t)=-1*y__1(t)+9*y__2(t)-3*y__3(t),diff(y__3(t),t)=-2*y__1(t)+2*y__2(t)+4*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$\begin{aligned}y_1(t) &= e^{6t}(c_3t + c_2) \\y_2(t) &= (-2c_3t^2 - 4c_2t - 3c_3t + c_1) e^{6t} \\y_3(t) &= e^{6t}(-2c_3t^2 - 4c_2t - 2c_3t + c_1 + c_2 + c_3)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 106

```
DSolve[{y1'[t]==5*y1[t]-1*y2[t]+1*y3[t],y2'[t]==-1*y1[t]+9*y2[t]-3*y3[t],y3'[t]==-2*y1[t]+2*y2[t]+4*y3[t]},y1[t],y2[t],y3[t],t]
```

$$\begin{aligned}y_1(t) &\rightarrow -e^{6t}(c_1(t-1) + (c_2 - c_3)t) \\y_2(t) &\rightarrow e^{6t}(2(c_1 + c_2 - c_3)t^2 - (c_1 - 3c_2 + 3c_3)t + c_2) \\y_3(t) &\rightarrow e^{6t}(2(c_1 + c_2 - c_3)t^2 - 2(c_1 - c_2 + c_3)t + c_3)\end{aligned}$$

22.25 problem section 10.5, problem 25

Internal problem ID [1628]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 25.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = y_1(t) + 10y_2(t) - 12y_3(t)$$

$$y_2'(t) = 2y_1(t) + 2y_2(t) + 3y_3(t)$$

$$y_3'(t) = 2y_1(t) - y_2(t) + 6y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 86

```
dsolve([diff(y__1(t),t)=1*y__1(t)+10*y__2(t)-12*y__3(t),diff(y__2(t),t)=2*y__1(t)+2*y__2(t)+
```

$$y_1(t) = e^{3t}(c_3t^2 + c_2t + c_1)$$
$$y_2(t) = -\frac{e^{3t}(6c_3t^2 + 6c_2t + 6c_3t + 6c_1 + 3c_2 + 4c_3)}{6}$$
$$y_3(t) = -\frac{e^{3t}(18c_3t^2 + 18c_2t + 18c_3t + 18c_1 + 9c_2 + 10c_3)}{18}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 118

```
DSolve[{y1'[t]==1*y1[t]+10*y2[t]-12*y3[t],y2'[t]==2*y1[t]+2*y2[t]+3*y3[t],y3'[t]==2*y1[t]-1*
```

$$y_1(t) \rightarrow -e^{3t}(c_1(2t - 1) + c_2t(9t - 10) + 3c_3(4 - 3t)t)$$
$$y_2(t) \rightarrow e^{3t}(9(c_2 - c_3)t^2 + (2c_1 - c_2 + 3c_3)t + c_2)$$
$$y_3(t) \rightarrow e^{3t}(9(c_2 - c_3)t^2 + (2c_1 - c_2 + 3c_3)t + c_3)$$

22.26 problem section 10.5, problem 26

Internal problem ID [1629]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 26.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -6y_1(t) - 4y_2(t) - 4y_3(t)$$

$$y_2'(t) = 2y_1(t) - y_2(t) + y_3(t)$$

$$y_3'(t) = 2y_1(t) + 3y_2(t) + y_3(t)$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 73

```
dsolve([diff(y__1(t),t)=-6*y__1(t)-4*y__2(t)-4*y__3(t),diff(y__2(t),t)=2*y__1(t)-1*y__2(t)+1
```

$$y_1(t) = e^{-2t}(c_3t + c_2)$$

$$y_2(t) = \frac{(2c_3t^2 + 4c_2t - c_3t + 4c_1)e^{-2t}}{4}$$

$$y_3(t) = -\frac{e^{-2t}(2c_3t^2 + 4c_2t + 3c_3t + 4c_1 + 4c_2 + c_3)}{4}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 100

```
DSolve[{y1'[t]==-6*y1[t]-4*y2[t]-4*y3[t],y2'[t]==2*y1[t]-1*y2[t]+1*y3[t],y3'[t]==2*y1[t]+3*y
```

$$y_1(t) \rightarrow e^{-2t}(c_1(1 - 4t) - 4(c_2 + c_3)t)$$

$$y_2(t) \rightarrow e^{-2t}(-2(c_1 + c_2 + c_3)t^2 + (2c_1 + c_2 + c_3)t + c_2)$$

$$y_3(t) \rightarrow e^{-2t}(2(c_1 + c_2 + c_3)t^2 + 2c_1t + 3(c_2 + c_3)t + c_3)$$

22.27 problem section 10.5, problem 27

Internal problem ID [1630]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 27.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 2y_2(t) - 2y_3(t) \\y_2'(t) &= -y_1(t) + 5y_2(t) - 3y_3(t) \\y_3'(t) &= y_1(t) + y_2(t) + y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 74

```
dsolve([diff(y__1(t),t)=0*y__1(t)+2*y__2(t)-2*y__3(t),diff(y__2(t),t)=-1*y__1(t)+5*y__2(t)-3
```

$$\begin{aligned}y_1(t) &= e^{2t}(c_3t + c_2) \\y_2(t) &= \frac{(2c_3t^2 + 4c_2t + 3c_3t + 2c_1)e^{2t}}{2} \\y_3(t) &= \frac{e^{2t}(2c_3t^2 + 4c_2t + c_3t + 2c_1 - 2c_2 - c_3)}{2}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 108

```
DSolve[{y1'[t]==0*y1[t]+2*y2[t]-2*y3[t],y2'[t]==-1*y1[t]+5*y2[t]-3*y3[t],y3'[t]==1*y1[t]+1*y
```

$$\begin{aligned}y_1(t) &\rightarrow -e^{2t}(c_1(2t - 1) + 2(c_3 - c_2)t) \\y_2(t) &\rightarrow e^{2t}(-2(c_1 - c_2 + c_3)t^2 - (c_1 - 3c_2 + 3c_3)t + c_2) \\y_3(t) &\rightarrow e^{2t}(-2(c_1 - c_2 + c_3)t^2 + (c_1 + c_2 - c_3)t + c_3)\end{aligned}$$

22.28 problem section 10.5, problem 28

Internal problem ID [1631]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 28.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -2y_1(t) - 12y_2(t) + 10y_3(t)$$

$$y_2'(t) = 2y_1(t) - 24y_2(t) + 11y_3(t)$$

$$y_3'(t) = 2y_1(t) - 24y_2(t) + 8y_3(t)$$

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 84

```
dsolve([diff(y__1(t),t)=-2*y__1(t)-12*y__2(t)+10*y__3(t),diff(y__2(t),t)=2*y__1(t)-24*y__2(t)
```

$$y_1(t) = e^{-6t}(c_3t^2 + c_2t + c_1)$$
$$y_2(t) = -\frac{e^{-6t}(18c_3t^2 + 18c_2t - 24c_3t + 18c_1 - 12c_2 + 5c_3)}{36}$$
$$y_3(t) = -\frac{e^{-6t}(6c_3t^2 + 6c_2t - 6c_3t + 6c_1 - 3c_2 + c_3)}{6}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 131

```
DSolve[{y1'[t]==-2*y1[t]-12*y2[t]+10*y3[t],y2'[t]==2*y1[t]-24*y2[t]+11*y3[t],y3'[t]==2*y1[t]
```

$$y_1(t) \rightarrow e^{-6t}(c_1(6t^2 + 4t + 1) + 2t(c_3(12t + 5) - 6c_2(3t + 1)))$$
$$y_2(t) \rightarrow e^{-6t}(-3(c_1 - 6c_2 + 4c_3)t^2 + (2c_1 - 18c_2 + 11c_3)t + c_2)$$
$$y_3(t) \rightarrow e^{-6t}(-6(c_1 - 6c_2 + 4c_3)t^2 + 2(c_1 - 12c_2 + 7c_3)t + c_3)$$

22.29 problem section 10.5, problem 29

Internal problem ID [1632]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 29.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -y_1(t) - 12y_2(t) + 8y_3(t)$$

$$y_2'(t) = y_1(t) - 9y_2(t) + 4y_3(t)$$

$$y_3'(t) = y_1(t) - 6y_2(t) + y_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 53

```
dsolve([diff(y__1(t),t)=-1*y__1(t)-12*y__2(t)+8*y__3(t),diff(y__2(t),t)=1*y__1(t)-9*y__2(t)+
```

$$y_1(t) = e^{-3t}(c_3t + c_2)$$

$$y_2(t) = \frac{e^{-3t}(c_3t + 2c_1 + c_2)}{2}$$

$$y_3(t) = \frac{e^{-3t}(4c_3t + 12c_1 + 4c_2 + c_3)}{8}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 79

```
DSolve[{y1'[t]==-1*y1[t]-12*y2[t]+8*y3[t],y2'[t]==1*y1[t]-9*y2[t]+4*y3[t],y3'[t]==1*y1[t]-6*
```

$$y1(t) \rightarrow e^{-3t}(2c_1t - 12c_2t + 8c_3t + c_1)$$

$$y2(t) \rightarrow e^{-3t}((c_1 - 6c_2 + 4c_3)t + c_2)$$

$$y3(t) \rightarrow e^{-3t}((c_1 - 6c_2 + 4c_3)t + c_3)$$

22.30 problem section 10.5, problem 30

Internal problem ID [1633]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 30.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -4y_1(t) - y_3(t) \\y_2'(t) &= -y_1(t) - 3y_2(t) - y_3(t) \\y_3'(t) &= y_1(t) - 2y_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve([diff(y__1(t),t)=-4*y__1(t)-0*y__2(t)-1*y__3(t),diff(y__2(t),t)=-1*y__1(t)-3*y__2(t)-
```

$$\begin{aligned}y_1(t) &= e^{-3t}(c_3t + c_2) \\y_2(t) &= e^{-3t}(c_3t + c_1 + c_2) \\y_3(t) &= -e^{-3t}(c_3t + c_2 + c_3)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 63

```
DSolve[{y1'[t]==-4*y1[t]-0*y2[t]-1*y3[t],y2'[t]==-1*y1[t]-3*y2[t]-1*y3[t],y3'[t]==1*y1[t]-0*
```

$$\begin{aligned}y_1(t) &\rightarrow e^{-3t}(c_1(-t) - c_3t + c_1) \\y_2(t) &\rightarrow e^{-3t}(c_2 - (c_1 + c_3)t) \\y_3(t) &\rightarrow e^{-3t}((c_1 + c_3)t + c_3)\end{aligned}$$

22.31 problem section 10.5, problem 31

Internal problem ID [1634]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 31.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -3y_1(t) - 3y_2(t) + 4y_3(t)$$

$$y_2'(t) = 4y_1(t) + 5y_2(t) - 8y_3(t)$$

$$y_3'(t) = 2y_1(t) + 3y_2(t) - 5y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

```
dsolve([diff(y__1(t),t)=-3*y__1(t)-3*y__2(t)+4*y__3(t),diff(y__2(t),t)=4*y__1(t)+5*y__2(t)-8*y__3(t),diff(y__3(t),t)=2*y__1(t)+3*y__2(t)-5*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$y_1(t) = e^{-t}(c_3t + c_2)$$
$$y_2(t) = e^{-t}(-2c_3t + c_1 - 2c_2)$$
$$y_3(t) = \frac{e^{-t}(-4c_3t + 3c_1 - 4c_2 + c_3)}{4}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 83

```
DSolve[{y1'[t]==-3*y1[t]-3*y2[t]+4*y3[t],y2'[t]==4*y1[t]+5*y2[t]-8*y3[t],y3'[t]==2*y1[t]+3*y2[t]-5*y3[t]},y1[t],y2[t],y3[t]]
```

$$y_1(t) \rightarrow e^{-t}(-2c_1t - 3c_2t + 4c_3t + c_1)$$
$$y_2(t) \rightarrow e^{-t}((4c_1 + 6c_2 - 8c_3)t + c_2)$$
$$y_3(t) \rightarrow e^{-t}((2c_1 + 3c_2 - 4c_3)t + c_3)$$

22.32 problem section 10.5, problem 32

Internal problem ID [1635]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.5, constant coefficient homogeneous system II. Page 555

Problem number: section 10.5, problem 32.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -3y_1(t) - y_2(t)$$

$$y_2'(t) = y_1(t) - y_2(t)$$

$$y_3'(t) = -y_1(t) - y_2(t) - 2y_3(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve([diff(y__1(t),t)=-3*y__1(t)-1*y__2(t)+0*y__3(t),diff(y__2(t),t)=1*y__1(t)-1*y__2(t)+0
```

$$y_1(t) = e^{-2t}(c_3t + c_2)$$

$$y_2(t) = -e^{-2t}(c_3t + c_2 + c_3)$$

$$y_3(t) = e^{-2t}(c_3t + c_1 + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 63

```
DSolve[{y1'[t]==-3*y1[t]-1*y2[t]+0*y3[t],y2'[t]==1*y1[t]-1*y2[t]+0*y3[t],y3'[t]==-1*y1[t]-1*
```

$$y_1(t) \rightarrow e^{-2t}(c_1(-t) - c_2t + c_1)$$

$$y_2(t) \rightarrow e^{-2t}((c_1 + c_2)t + c_2)$$

$$y_3(t) \rightarrow e^{-2t}(c_3 - (c_1 + c_2)t)$$

23 Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

23.1	problem section 10.6, problem 1	848
23.2	problem section 10.6, problem 2	849
23.3	problem section 10.6, problem 3	850
23.4	problem section 10.6, problem 4	851
23.5	problem section 10.6, problem 5	852
23.6	problem section 10.6, problem 6	854
23.7	problem section 10.6, problem 7	855
23.8	problem section 10.6, problem 8	856

23.1 problem section 10.6, problem 1

Internal problem ID [1636]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 1.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= -y_1(t) + 2y_2(t) \\y_2'(t) &= -5y_1(t) + 5y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 48

```
dsolve([diff(y__1(t),t)=-1*y__1(t)+2*y__2(t),diff(y__2(t),t)=-5*y__1(t)+5*y__2(t)],singsol=a
```

$$\begin{aligned}y_1(t) &= e^{2t}(c_1 \sin(t) + c_2 \cos(t)) \\y_2(t) &= \frac{e^{2t}(3c_1 \sin(t) - c_2 \sin(t) + c_1 \cos(t) + 3c_2 \cos(t))}{2}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 55

```
DSolve[{y1'[t]==-1*y1[t]+2*y2[t],y2'[t]==-5*y1[t]+5*y2[t]},{y1[t],y2[t]},t,IncludeSingularSo
```

$$\begin{aligned}y_1(t) &\rightarrow e^{2t}(c_1 \cos(t) + (2c_2 - 3c_1) \sin(t)) \\y_2(t) &\rightarrow e^{2t}(c_2(3 \sin(t) + \cos(t)) - 5c_1 \sin(t))\end{aligned}$$

23.2 problem section 10.6, problem 2

Internal problem ID [1637]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 2.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -11y_1(t) + 4y_2(t)$$

$$y_2'(t) = -26y_1(t) + 9y_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 60

```
dsolve([diff(y__1(t),t)=-11*y__1(t)+4*y__2(t),diff(y__2(t),t)=-26*y__1(t)+9*y__2(t)],singsol
```

$$y_1(t) = e^{-t}(c_1 \sin(2t) + c_2 \cos(2t))$$
$$y_2(t) = \frac{e^{-t}(5c_1 \sin(2t) - c_2 \sin(2t) + c_1 \cos(2t) + 5c_2 \cos(2t))}{2}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 64

```
DSolve[{y1'[t]==-11*y1[t]+4*y2[t],y2'[t]==-26*y1[t]+9*y2[t]},{y1[t],y2[t]},t,IncludeSingular
```

$$y_1(t) \rightarrow e^{-t}(c_1 \cos(2t) + (2c_2 - 5c_1) \sin(2t))$$
$$y_2(t) \rightarrow e^{-t}(c_2 \cos(2t) + (5c_2 - 13c_1) \sin(2t))$$

23.3 problem section 10.6, problem 3

Internal problem ID [1638]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 3.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = y_1(t) + 2y_2(t)$$

$$y_2'(t) = -4y_1(t) + 5y_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 57

```
dsolve([diff(y__1(t),t)=1*y__1(t)+2*y__2(t),diff(y__2(t),t)=-4*y__1(t)+5*y__2(t)],singsol=all)
```

$$y_1(t) = e^{3t}(c_1 \sin(2t) + c_2 \cos(2t))$$

$$y_2(t) = e^{3t}(c_1 \sin(2t) - c_2 \sin(2t) + c_1 \cos(2t) + c_2 \cos(2t))$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 60

```
DSolve[{y1'[t]==1*y1[t]+2*y2[t],y2'[t]==-4*y1[t]+5*y2[t]},{y1[t],y2[t]},t,IncludeSingularSolutions->True]
```

$$y_1(t) \rightarrow e^{3t}(c_1 \cos(2t) + (c_2 - c_1) \sin(2t))$$

$$y_2(t) \rightarrow e^{3t}(c_2 \cos(2t) + (c_2 - 2c_1) \sin(2t))$$

23.4 problem section 10.6, problem 4

Internal problem ID [1639]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 4.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}y_1'(t) &= 5y_1(t) - 6y_2(t) \\y_2'(t) &= 3y_1(t) - y_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 58

```
dsolve([diff(y__1(t),t)=5*y__1(t)-6*y__2(t),diff(y__2(t),t)=3*y__1(t)-1*y__2(t)],singsol=all
```

$$\begin{aligned}y_1(t) &= e^{2t}(c_1 \sin(3t) + c_2 \cos(3t)) \\y_2(t) &= \frac{e^{2t}(c_1 \sin(3t) + c_2 \sin(3t) - c_1 \cos(3t) + c_2 \cos(3t))}{2}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 60

```
DSolve[{y1'[t]==5*y1[t]-6*y2[t],y2'[t]==3*y1[t]-1*y2[t]},{y1[t],y2[t]},t,IncludeSingularSolu
```

$$\begin{aligned}y_1(t) &\rightarrow e^{2t}(c_1 \cos(3t) + (c_1 - 2c_2) \sin(3t)) \\y_2(t) &\rightarrow e^{2t}(c_2 \cos(3t) + (c_1 - c_2) \sin(3t))\end{aligned}$$

23.5 problem section 10.6, problem 5

Internal problem ID [1640]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 5.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -3y_1(t) - 3y_2(t) + y_3(t)$$

$$y_2'(t) = 2y_2(t) + 2y_3(t)$$

$$y_3'(t) = 5y_1(t) + y_2(t) + y_3(t)$$

✓ Solution by Maple

Time used: 0.282 (sec). Leaf size: 1975

`dsolve([diff(y__1(t),t)=-3*y__1(t)-3*y__2(t)+1*y__3(t),diff(y__2(t),t)=0*y__1(t)+2*y__2(t)+2`

Expression too large to display

$$\begin{aligned}
 y_2(t) = & e^{\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{6}+7\right)t}{(540+6\sqrt{6042})^{\frac{1}{3}}}} \sin\left(\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{6}-42\right)t\sqrt{3}36^{\frac{1}{3}}}{36(90+\sqrt{6042})^{\frac{1}{3}}}\right) c_2 \\
 & + e^{\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{6}+7\right)t}{(540+6\sqrt{6042})^{\frac{1}{3}}}} \cos\left(\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{6}-42\right)t\sqrt{3}36^{\frac{1}{3}}}{36(90+\sqrt{6042})^{\frac{1}{3}}}\right) c_3 \\
 y_3(t) = & c_1 e^{-\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{3}+42\right)t}{3(540+6\sqrt{6042})^{\frac{1}{3}}}} \\
 & - 2c_1 \left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{3}+42\right) e^{-\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{3}+42\right)t}{3(540+6\sqrt{6042})^{\frac{1}{3}}}} + c_2 \left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{3}+42\right) e^{\frac{\left(\frac{(540+6\sqrt{6042})^{\frac{2}{3}}}{6}+42\right)t}{6(540+6\sqrt{6042})^{\frac{1}{3}}}} \sin
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 187

`DSolve[{y1'[t]==3*y1[t]-3*y2[t]+1*y3[t],y2'[t]==0*y1[t]+2*y2[t]+2*y3[t],y3'[t]==5*y1[t]+1*y2`

$$\begin{aligned}
 y_1(t) & \rightarrow \frac{1}{4}e^{-2t}((3c_1 - c_2 + c_3)e^{6t} \cos(2t) + (c_1 - 3c_2 - c_3)e^{6t} \sin(2t) + c_1 + c_2 - c_3) \\
 y_2(t) & \rightarrow \frac{1}{4}e^{-2t}(-(c_1 - 3c_2 - c_3)e^{6t} \cos(2t) + (3c_1 - c_2 + c_3)e^{6t} \sin(2t) + c_1 + c_2 - c_3) \\
 y_3(t) & \rightarrow \frac{1}{2}e^{-2t}((c_1 + c_2 + c_3)e^{6t} \cos(2t) + 2(c_1 - c_2)e^{6t} \sin(2t) - c_1 - c_2 + c_3)
 \end{aligned}$$

23.6 problem section 10.6, problem 6

Internal problem ID [1641]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 6.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -3y_1(t) + 3y_2(t) + y_3(t)$$

$$y_2'(t) = y_1(t) - 5y_2(t) - 3y_3(t)$$

$$y_3'(t) = -3y_1(t) + 7y_2(t) + 3y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 115

```
dsolve([diff(y__1(t),t)=-3*y__1(t)+3*y__2(t)+1*y__3(t),diff(y__2(t),t)=1*y__1(t)-5*y__2(t)-3*y__3(t),diff(y__3(t),t)=-3*y__1(t)+7*y__2(t)+3*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$y_1(t) = e^{-t}c_1 + c_2e^{-2t} \sin(2t) + c_3e^{-2t} \cos(2t)$$

$$y_2(t) = e^{-t}c_1 + c_2e^{-2t} \cos(2t) - c_3e^{-2t} \sin(2t)$$

$$y_3(t) = -e^{-t}c_1 + c_2e^{-2t} \sin(2t) - c_2e^{-2t} \cos(2t) + c_3e^{-2t} \cos(2t) + c_3e^{-2t} \sin(2t)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 158

```
DSolve[{y1'[t]==-3*y1[t]+3*y2[t]+1*y3[t],y2'[t]==1*y1[t]-5*y2[t]-3*y3[t],y3'[t]==-3*y1[t]+7*y2[t]+3*y3[t]},y1[t],y2[t],y3[t]]
```

$$y_1(t) \rightarrow e^{-2t}((c_1 - c_2 - c_3)e^t + (c_2 + c_3) \cos(2t) + (-c_1 + 2c_2 + c_3) \sin(2t))$$

$$y_2(t) \rightarrow e^{-2t}((c_1 - c_2 - c_3)e^t + (-c_1 + 2c_2 + c_3) \cos(2t) - (c_2 + c_3) \sin(2t))$$

$$y_3(t) \rightarrow e^{-2t}((-c_1 + c_2 + c_3)e^t + (c_1 - c_2) \cos(2t) + (-c_1 + 3c_2 + 2c_3) \sin(2t))$$

23.7 problem section 10.6, problem 7

Internal problem ID [1642]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 7.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = 2y_1(t) + y_2(t) - y_3(t)$$

$$y_2'(t) = y_2(t) + y_3(t)$$

$$y_3'(t) = y_1(t) + y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 70

```
dsolve([diff(y__1(t),t)=2*y__1(t)+1*y__2(t)-1*y__3(t),diff(y__2(t),t)=0*y__1(t)+1*y__2(t)+1*
```

$$y_1(t) = c_1 e^{2t} + c_2 e^t \cos(t) - c_3 e^t \sin(t)$$

$$y_2(t) = c_1 e^{2t} - c_2 e^t \cos(t) + c_3 e^t \sin(t)$$

$$y_3(t) = c_1 e^{2t} + c_2 e^t \sin(t) + c_3 e^t \cos(t)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 129

```
DSolve[{y1'[t]==2*y1[t]+1*y2[t]-1*y3[t],y2'[t]==0*y1[t]+1*y2[t]+1*y3[t],y3'[t]==1*y1[t]+0*y2
```

$$y_1(t) \rightarrow \frac{1}{2} e^t (-2c_3 \sin(t) + c_2 (e^t + \sin(t) - \cos(t)) + c_1 (e^t + \sin(t) + \cos(t)))$$

$$y_2(t) \rightarrow \frac{1}{2} e^t ((c_1 + c_2) e^t + (c_2 - c_1) \cos(t) - (c_1 + c_2 - 2c_3) \sin(t))$$

$$y_3(t) \rightarrow \frac{1}{2} e^t ((c_1 + c_2) e^t - (c_1 + c_2 - 2c_3) \cos(t) + (c_1 - c_2) \sin(t))$$

23.8 problem section 10.6, problem 8

Internal problem ID [1643]

Book: Elementary differential equations with boundary value problems. William F. Trench. Brooks/Cole 2001

Section: Chapter 10 Linear system of Differential equations. Section 10.6, constant coefficient homogeneous system III. Page 566

Problem number: section 10.6, problem 8.

ODE order: 1.

ODE degree: 1.

Solve

$$y_1'(t) = -3y_1(t) + y_2(t) - 3y_3(t)$$

$$y_2'(t) = 4y_1(t) - y_2(t) + 2y_3(t)$$

$$y_3'(t) = 4y_1(t) - 2y_2(t) + 3y_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 133

```
dsolve([diff(y__1(t),t)=-3*y__1(t)+1*y__2(t)-3*y__3(t),diff(y__2(t),t)=4*y__1(t)-1*y__2(t)+2*y__3(t),diff(y__3(t),t)=4*y__1(t)-2*y__2(t)+3*y__3(t)),y__1(t),y__2(t),y__3(t))
```

$$y_1(t) = c_1 e^t + c_2 e^{-t} \sin(2t) + c_3 e^{-t} \cos(2t)$$

$$y_2(t) = c_1 e^t - c_2 e^{-t} \sin(2t) - c_2 e^{-t} \cos(2t) - c_3 e^{-t} \cos(2t) + c_3 e^{-t} \sin(2t)$$

$$y_3(t) = -c_1 e^t - c_2 e^{-t} \sin(2t) - c_2 e^{-t} \cos(2t) - c_3 e^{-t} \cos(2t) + c_3 e^{-t} \sin(2t)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 163

```
DSolve[{y1'[t]==-3*y1[t]+1*y2[t]-3*y3[t],y2'[t]==4*y1[t]-1*y2[t]+2*y3[t],y3'[t]==4*y1[t]-2*y2[t]+3*y3[t]},y1[t],y2[t],y3[t]]
```

$$y_1(t) \rightarrow \frac{1}{2} e^{-t} ((c_2 - c_3) e^{2t} + (2c_1 - c_2 + c_3) \cos(2t) - 2(c_1 + c_3) \sin(2t))$$

$$y_2(t) \rightarrow \frac{1}{2} e^{-t} ((c_2 - c_3) e^{2t} + (c_2 + c_3) \cos(2t) + (4c_1 - c_2 + 3c_3) \sin(2t))$$

$$y_3(t) \rightarrow \frac{1}{2} e^{-t} ((c_3 - c_2) e^{2t} + (c_2 + c_3) \cos(2t) + (4c_1 - c_2 + 3c_3) \sin(2t))$$